UNITED STATES ENVIRONMENTAL PROTECTION AGENCY



U.S. ENVIRONMENTAL PROTECTION AGEN
Office of Pesticide Programs
Registration Division (H7505C)
401 MM St., S.W.
Washington, D.C. 20460

EPA Reg. Date of Number: Issuance:

52287-3

MAR 1 8 1994

Product:

NOTICE OF PESTICIDE:

x
Registration

Reregistration

(under FIFRA, as amended)

Name of Pesticide

Term of Issuance:

Conditional

Ronstar 0.70 with

Fertilizer

Name and Address of Registrant (include ZIP Code):

H. R. McLane, Inc.

7210 Red Road Suite 206

Miami, FL 33143

in labeling in substance Note: Changes differing from that accepted by the connection with this registration must be submitted to and accepted Division prior to use of the label in commerce. Registration refer to the above EPA registration on this product always correspondence number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

- 1. Submit and/or cite all data required for registration/ reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
- 2. Change the label by revising the EPA Registration Number to read, "EPA Reg. No. 52287-49".
- 3. Submit two copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFAA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

	COHCURRENCES								
SYMBOL	dignature of	THE PART OF THE	Official:			ete:			
SURNAM	MODERAL								
DATE	3/18/14·	************		••••••••	************		••••••••		

EPA Form 1320-1A (1/90)

Printed on Recycled Paper

OFFICIAL FILE COPY

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Joanne I. Miller Product Manager (23) Fungicide-Herbicide Branch Registration Division BEST WINITHDER DOLL !

RONSTAR® 0.70 with FERT

for sale to, use & Storage by TURF AND LANDSCAPE PERSONNEL ONLY 50 lb. Bag Treats up to 7,700 Sq.Ft.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial need production, or for research purposes.

FOR PARKS, GOLF COURSES, ATHLETIC FIELDS, AND OTHER RECREATIONAL AND ORNAMENTAL TURFGRASS AREAS.

—Not For Use On Home Lewns—

Controls: Crabgrass, Annual Bluegrass, Goosegrass, Florida Pusiey, Oxalls, Stinging Nettle, Carpetgrass, Pigweed and Other Weeds Listed

(hadiszon; [2-let-buhl-4-(2.4-dichloro-5-leopropoxychenyl) &-1,3,4-oxediszolin-5-one] INERT INGREDIENTS: Total 100.0%

Contains RONSTARD Herbittle & Reghtund Trace Mait of Co.

KEEP OUT OF REACH OF CHILDREN WARNING

VVARNING

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED* Call a bityelctan or Policin Control Center immediately. Induce vorniling by giving victin 1 or 2 glasses of water and touching back of throat with finger. On not induce vomiting or give anything by mouth a an unconscious or comutaing person. [E-th] EYES: Firsh eyes with plenty of water, Gat medical attention if iritiation persons. [E-th] House IF ON SKIN: Yeah with prenty of ecop and water, Gat medical attention if iritiation persons. [E-th] PRECAUTIONAR PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS

HAZARIDS TO HUMANS AND DOMESTIC ANMALS.

"RNING: Causes are and skin irrustion. Avoid contact with eth, a or clothing. Thotoughly wesh face and hands with scop and water or exposure anno before earing or smolding. Wear protective cohing including long-steered whit and gives when handling, Leunder exposure do before existing or smolding. Wear protective cohing including long-steered whit and gives when handling, Leunder exposure do the person of the swift of the protective cohing including long-steered whit and gives when handling, Leunder exposure do to be a swift of the protective cohing including long-steered whit and gives when handling in the mean high water match to not apply when weather conditions flavor diff from steep treated. Do not apply where runof is kitch to occur. Do not contaminate water when disposing of equipment washwaters. Apply only as specified on this table. Do not slow people (offer than the application) or pets on trastment area during application. Do not enter treatment area until digit has earlied.

DIRECTIONS FOR USE

...

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent, with its lebeling.

STORAGE AND DISPOSAL

Do not contestinate water, food, or feed by storage or disposal storage. Store product in its original beg, in a cool, dry, locked place out of reach of children, in case of spill, sweep up, and use according to directions or dispose of ser follows.

PESTICED DISPOSAL Pesticide waster are toxic. Improper disposal of excess pessicide, or insette is a violation of Federal law. If these wasters cannot be disposed of by use according to label instruming waster specially your State Pesticids or Environmental Control Agency, with flazardous Waste representative at the marcest EPA regioner office for guidence. CONTAINER DISPOSAL Compiletely empty bag into application equipment. Then dispose of empty bag into application equipment. Then dispose of empty bag its a sankery landition by inchesiation, or, it stowed by State and local authorises, by burning, it nned stay out of smoke,

GUARANTEED FERTILIZER ANALYSIS GUAKANY FEEU PERTINGGETS -- 198 Hilliad pithopon Ammendmedi Pringen Pringen 12 5 Walso Schiple Pithopon Pringen 12 5 Walso Schiple Pithopon Principal Pringen 275 Amiliado Pringen 275 Schiple Pithopon 275 Schiple Pithopon Schiple Pithopon Description Amilianous Sulling Under noce as Mr. S'S from on Fo 1,0% Chimina et

When applied according to Directions this product can be used safely on the following established turigrass varieties: BERMUDAGRASS, PERENNIAL BLUEGRASS, PERENNIAL RYEGRASS VARIETIES, ST. AUGUSTINE GRASS VARIETIES, TALL FESCUE, and 20YSIA-ORASS lawne and grammental turit, it may be used for Apartment and Condornhum complexes, Parks, Golf Courses, Altiella Fields, Cemetales and all other companies and recreational turigrass ereas

EPA Rep to 52207-0 EPA EST. NO 52207-FL-1

A PRE-EMERGENT WEED CONTROL PLUS FERTILIZER FOR PERENNIAL BLUEGRASS, PERENNIAL RYEGRASS ST. AUGUSTINEGRASS, BERMUDAGRAS TALL FESCUE and ZOYSIAGRASS.

TALL FESCUE BIND ZOYSIAGRASS.

Do not use on find Fescue, Sentatives yestelles, Dichondra, or Canilbade plais Sereure damage will occur, to not use on Flotin greates of less due to verying cultural practices used.

Egr. St. Augustinegrass application raises above the 50 lbs. per 5,20 eq. T. will temporesty cincture and damage, the louf it a higher raise I necessary (such as 50 lbs./1,850 eq.ft. will femporesty concerned and damage, theridas Pursity or Carpet to control General managements. Springe, Pre Annus, Findes Pursity or Carpet grass, but demage can be superied to the \$1 Augustine, therefore applicates, but demage can be solerated and when good cultura practices are claimed in order to revive the demaged but device and finded to revive the demaged but device and control of the control can be supplying the strategy average to the product contact with flowing and strute except as recommende on the Orinstrenial use Supplying that I about a per product. For newly seeded issues do not spept with preduct until after second of the year Spring pre-immergency weed control can then be supplied. Respecting treated Avers; Delay new seeding until 4 month electrical and control and then be second.

CONTROLS GRASSY WEEDS and BROADLEAF WEEDS

united Bleegrass (Pos Annue), Barmard Orass, Crabgrass, Fall Panicum, Floride Pustey, Goose grass, Green Fusiel, Oxafe, Stinging Nettle, Carpet grass, Ploweed.
HOW TO APPLY

FIUW IU A? FLT

For best weed control take the area free of leaves and thatch and mo
if necessary so that granules will reach the soa surface. Measure in
area to be treated and uniformly apply the required amount using
fettiger spreader (see Spreader Satings chart). Do not exply to tur
when it is vest. Do not ever-lep spreader applications, on SI Augusine
grass eithes discoloration "streaking" thay temporarily occur fro
"double rate". After applications if rain is not expected within 24 hour
aprintile the bested turk with 1/4 to 1/2 leich of water to get mission
united.

HOW MUCH TO USE

HOW MUCH TO USE

FOR CRABGRASS

Apply 50 bs // 700 Sq Ft. when furthes been in a yearly manieranc program (equal to 2 hs. All herbicide per acre.) Apply 50 bs. to tree to 3 hs. All herbicide per acre.) Apply 50 bs. to tree to 3 hs. All herbicide per acre. Apply in early spring 1 to 2 weeks prio to weed seed germination. This product is a pre-emergence herbicide it will not control emerged end growing weeds; only germinating wee seeds. Lists applications will control enly those weed seeds their germination and early those weeds seeds their germination and early those weeds acred to travely germinate until fall spring or early summer Apply 50 bs. to teat 3,850 Sq Ft. (4 bs. All/scre) and apply 1 to 2 week prior to used germination. See Chart for When To Apply. See Us Precautions for St. Augustine grave.

Apply 50 bs. to teat 3,850 Sq Ft. (6 bs. All/scre) and apply 1 to 2 week prior to used germination.

Apply 50 bs. to teat 3,850 Sq Ft. (6 bs. All/scre) and apply 1 to 2 week prior to used germination.

Apply 50 bs. to teat 3,850 Sq Ft. (6 bs. All/scre) and apply 1 to 2 week prior to seed germination. The prior to the seture institute per acre. Apply in late summer to late or in early spin 1 to 2 weeks before expected weed seed germination. Vall not control annual bluegrass present at time of application, therefore only germination general and provide and

<u> </u>	\$PF	PAUSE SETT	INGS	. T				
Spreader		7,785 5g FI	5,200 Sq FI	3.000 Bq.FL				
L 1	Predual_	284_L/Acre	38A L/Acre					
Cyclene	SO De.	4 1/2	5 1/2	6 1/2				
Sec. 9 835	30 ba	3 1/2	4	41/2				
HOTE: Spreader role varies with walking speed								
Check paraget								

APPROPRIATE TIMES TO APPLY THIS PRE-EM	ERGSNI HLIKUKJUL
THE PUR YEAR AME YAM BEEN MAL	THE PART OF THE
1	3
3	فسندنس
22	4

1 For CTLABGRASS and other Grosses and Broadest Weeds permissing in the Spring in Northern fort.
2 For GOOSEGRASS and other Late Germinaling Arctual Crosses and Broades Weeds seames PROSTRATE SPURGE. 28. OCCOSEGRASS in Smothern Turl 2. For POA ANMALA In Medition Turl. 28. For POA ANITUA Is Seathern Turl 4. For Fost-Winter-Spring, confeet of ANNUAL IVEEDS in Southern Turl. NOTICE: Steps and used seatment of this analisal othy of use, storage and/or hands of the product not in Societations; with the forms of this label (d 97287-3 sin503-10-84)

NET WEIGHT

HARREL EPA Rea No 52267-G (8.82267-5 pills 03-10-84)

HARRELL'S, INC. PO BOX 807 - LAKELAND FL 33802

NET WEIGHT 50 POUNDS

ACCEPTED
with COMMENTS
In EPA Letter Dated

Under the Federal Investicide. Fundicide and Rodenticide Act as unended for the posticide regimered under EPA Res. No.

BEST AVAILABLE COPY