



# RONSTAR® 0.67 with FERTILIZER

**FOR SALE TO, USE & STORAGE BY  
TURF AND LANDSCAPE PERSONNEL ONLY  
50 lb. Bag Treats up to 7,260 Sq.Ft.**

*Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.*

FOR PARKS, GOLF COURSES, ATHLETIC FIELDS, AND OTHER RECREATIONAL AND ORNAMENTAL TURFGRASS AREAS.

---Not For Use On Home Lawns---

**Controls: Crabgrass, Annual Bluegrass, Goosegrass, Florida Pusley, Oxalis, Stinging Nettle, Carpetgrass, Pigweed and Other Weeds Listed**

**ACTIVE INGREDIENT:**  
Oxadiazon; [2-tert-butyl-4-(2,4-dichloro-5-isopropoxyphenyl) Δ<sup>2</sup>-1,3,4-oxadiazolin-5-one] 0.67%  
**INERT INGREDIENTS:** 99.33%  
Total 100.00%

Contains RONSTAR® Herbicide a Registered Trade Mark of Rhone Poulenc Inc.

**A PRE-EMERGENT WEED CONTROL PLUS FERTILIZER FOR PERENNIAL BLUEGRASS, PERENNIAL RYEGRASS, ST. AUGUSTINEGRASS, BERMUDAGRASS, TALL FESCUE and ZOYSIAGRASS.**

### USE PRECAUTIONS

**Do not use on Red Fescue, Bentgrass varieties, Dichondra, or Centipede grass because damage will occur.** Do not use on Putting greens or "tees" due to varying cultural practices used.  
**For St. Augustinegrass** application rates above the 50 lbs. per 4,840 sq.ft. will temporarily discolor and damage the turf. If a higher rate is necessary (such as 50 lbs./3,630 sq.ft. or 50 lbs./4,174 sq.ft.) in order to control Goosegrass, Spurge, Poa Annua, Florida Pusley or Carpetgrass, turf damage can be expected to the St. Augustine, therefore apply only when that damage can be tolerated and when good cultural practices are planned in order to revive the damaged turf.  
**Avoid product contact with flowers and shrubs** except as recommended on the Ornamental Use Supplemental Labeling available for this product. **For newly seeded lawns** do not apply this product until after the second or third cutting. For best lawn care, place new grass seed in the fall of the year. Spring pre-emergence weed control can then be applied. **Reseeding treated Areas:** Delay new seeding until 4 months after treatment.

## KEEP OUT OF REACH OF CHILDREN WARNING

### STATEMENTS OF PRACTICAL TREATMENT

**IF SWALLOWED:** Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. **IF IN EYES:** Flush eyes with plenty of water. Get medical attention if irritation persists. **IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS  
PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**WARNING:** Causes eye and skin irritation. Avoid contact with skin, eyes or clothing. Thoroughly wash face and hands with soap and water after exposure and before eating or smoking. Wear protective clothing including long-sleeved shirt and gloves when handling. Launder exposed clothing before reuse. Do not allow domestic animals to graze treated areas. Avoid contamination of Feed and foodstuffs.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters. Apply only as specified on this label. Do not allow people (other than the applicator) or pets on treatment area during application. Do not enter treatment area until dust has settled.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **STORAGE:** Store product in its original bag, in a cool, dry, locked place out of reach of children. In case of spill, sweep up, and use according to directions or dispose of as follows.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

### GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N) - Variable  
Nitrate Nitrogen Variable  
Ammoniacal Nitrogen Variable  
Water Soluble Nitrogen and/or Urea Nitrogen Variable  
Water Insoluble Nitrogen Variable  
Available Phosphoric Acid (P<sub>2</sub>O<sub>5</sub>) - Variable  
Soluble Potash (K<sub>2</sub>O) - Variable  
Derived from Ammonium Sulfate, Ureaform, Muriate of Potash Manganese as Mn .5% Iron as Fe 1.0% Chlorine <1

## CONTROLS GRASSY WEEDS and BROADLEAF WEEDS

Annual Bluegrass (Poa Annua), Barnyard Grass, Crabgrass, Fall Panicum, Florida Pusley, Goose grass, Green Foxtail, Oxalis, Stinging Nettle, Carpet grass, Pigweed.

### HOW TO APPLY

For best weed control rake the area free of leaves and thatch and mow if necessary so that granules will reach the soil surface. Measure the area to be treated and uniformly apply the required amount using a fertilizer spreader (see Spreader Settings chart). Do not apply to turf when it is wet. Do not over-lap spreader applications, on St. Augustinegrass since discoloration "streaking" may temporarily occur from "double rate". After applications if rain is not expected within 24 hours sprinkle the treated turf with 1/4 to 1/2 inch of water to get maximum weed control.

### HOW MUCH TO USE

#### FOR CRABGRASS

Apply 50 lbs./7,260 sq.ft. when turf has been in a yearly maintenance program (equal to 2 lbs. A.I. herbicide per acre.) Apply 50 lbs. to treat 4,840 sq.ft. of lawn if it has been heavily infested with Crabgrass; equal to 3 lb. A.I. herbicide per acre. Apply in early spring 1 to 2 weeks prior to weed seed germination. This product is a pre-emergence herbicide. It will not control emerged and growing weeds; only germinating weed seeds. Late applications will control only those weed seeds that germinate after the application. See "When To Apply" Chart.

#### GOOSEGRASS and PROSTRATE SPURGE

These weeds do not usually germinate until late spring or early summer. Apply 50 lbs. to treat 3,630 sq.ft. (4 lbs. A.I./acre) and apply 1 to 2 weeks prior to seed germination. See Chart for When To Apply. See Use Precautions for St. Augustine grass

#### FOR ANNUAL BLUEGRASS (POA ANNUA)

Apply 50 lbs. per 3,630 to 4,170 sq.ft. of Lawn; equal to 3.5 to 4 lbs. active herbicide per acre. Apply in late summer to fall or in early spring 1 to 2 weeks before expected weed seed germination. Will not control annual bluegrass present at time of application; therefore only germinating seeds. Refer to chart for proper timing in your locale. Annual BARNYARDGRASS and PROSTRATE SPURGE may also be controlled by this same rate. See Use Precautions for St. Augustine grass

#### FOR PIGWEED, FLORIDA PUSLEY, OXALIS, STINGING NETTLE AND CARPETGRASS

Where these weeds are a problem apply 50 lbs. to treat 3,630 to 4,174 sq.ft. of Lawn; (equal to 3.5 to 4 lbs. Active Herbicide per acre) in early spring or 1 to 2 weeks before expected weed seed germination. Will not control of emerged and growing weeds; only effective against germinating weed seeds. See Use Precautions for St. Augustine grass


### SPREADER SETTINGS

Spreader	Amt. of Product	7,260 sq.ft. 2#A.I./Acre	4,840 sq.ft. 3#A.I./Acre	3,630 sq.ft. 4#A.I./Acre
Cyclone	50 lbs.	XX	XX	XX
Scotts #35	50 lbs.	XX	XX	XX

NOTE: Spreader rate varies with walking speed. Check spreader over known area.

### APPROPRIATE TIMES TO APPLY THIS PRE-EMERGENT HERBICIDE

JAN. FEB. MAR. APR. MAY JUNE JULY AUG. SEPT. OCT. NOV. DEC.



- For CRABGRASS and other Grasses and Broadleaf Weeds germinating in the Spring in Northern turf.
  - For GOOSEGRASS and other Late Germinating Annual Grasses and Broadleaf Weeds such as PROSTRATE SPURGE. 2a. GOOSEGRASS in Southern Turf.
  - For POA ANNUA in Northern Turf. 3a. For POA ANNUA in Southern Turf.
  - For Fall-Winter-Spring control of ANNUAL WEEDS in Southern Turf.
- NOTICE: Buyer and user assume all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label. (52-3amd.pm502-23-96 reduce al to 67)

When applied according to Directions this product can be used safely on the following established turfgrass varieties: BERMUDAGRASS, PERENNIAL BLUEGRASS, PERENNIAL RYEGRASS VARIETIES, ST. AUGUSTINE GRASS VARIETIES, TALL FESCUE, and ZOYSIAGRASS lawns and ornamental turf. It may be used for Apartment and Condominium complexes, Parks, Golf Courses, Athletic Fields, Cemeteries and all other ornamental and recreational turfgrass areas.

EPA Reg. No. 52287-3 EPA EST. NO. 52287-FL-1

# HARRELL'S, INC.

P.O. BOX 807 - LAKELAND FL 33802

NET WEIGHT  
50 POUNDS

