16

JUN 25 2003

Mr. Robert F. Bischoff Agent for HARRELL'S INC. Total Turf Consulting LLC P.O. Box 7928 Hilton Head, SC 29938

Dear Mr. Bischoff:

This is in response to your request dated May 30, 2003 to change the primary brand name for **EPA Registration 52287-3** from Ronstar 0.67 With Fertilizer to:

TURF FERTILIZER WITH RONSTAR® 0.67

Your request for the name change is accepted and the change will be made to the Agency's records. All future correspondence on this product should reference the new name.

Sincerely yours,

Sherada D. Hobgood Notifications Review Coordinator


Registration Division (7505C)

SEPA	Environmenta	United States I Protection Agency Ington, DC 20460	roim additive	Registration Amendment Other	OPP Identifier Number		
		Application for Pesti	cide - Section	1			
Company/Product Nur 2287-3	nber		A Product Manager NNE I. MILLER	Proposed Classification			
Company/Product (Na RONSTAR 0.67 WI		PM# 23		-			
ARRELL'S, INC. . O. BOX 807 AKELAND, FL 33	Applicant (Include ZIP Co	(b)(i) to: EPA	•	nilar or identical in o	th FIFRA Section 3(c)(3) composition and labeling CATION 5 2003		
		Section	- 11				
Notification - Expl planation: Use addi OTIFICATION OF PRIM	esponse to Agency letter ain below. tional page(s) if necessar IARY BRAND NAME CHA	y. (For section I and Section II NGE PER PR NOTICE 98-10 A //ISIONS/GUIDANCE OF PR NO	Agency letter de "Me Too" Applic Other - Explain b	ation. elow. RELATIVE TO PR NO	TICE 2001-6. THIS		
ATEMENT OF FORMI ATEMENT TO EPA. I	JLA OF THIS PRODUCT. FURTHER UNDERSTANI EQUIREMENTS OF 40 CF	52.46, AND NO OTHER CHANG I UNDERSTAND THAT IT IS A D THAT IF THIS NOTIFICATION FR 156.10 AND 40 CFR 152.46.	VIOLATION OF 18 U. IS NOT CONSISTEN THIS PRODUCT MAY	S.C. SEC.1001 TO WI IT WITH THE GUIDAN	LLFULLY MAKE ANY FALSE CE OF PR NOTICES 98-10		
Id-Resistant Peckaging Yes No Certification must submitted		Weter Soluble Yes No No. per container Weter Soluble Yes No Peckage wgt	No. per container	2. Type of Containe Metal Plastic Glass Paper Other			
Location of Nat Conter	ts Information Container	4. Size(s) Retail Container	5. Lo	cation of Label Direct	tions		
Manner in Which Labe	is Affixed to Product	Lithograph Paper glued Stenciled	Other				
		Section -	IV .				
Contact Point <i>(Comple</i>	nte items directly below i	or identification of individual to	be contacted, if nec	essary, to process th	is application.)		
ame ROBERT F. BISCHOFF		Title AGENT, TOTA	ENT, TOTALTURF CONSULTING LLC		Telephone No. (Include Area Code) (317) 873-0222		
	any knowlingly false or	Certification this form and all attachments misleading statement may be p			6. Jate Application Received (Stamped)		
signature Robert Fi	Berch	3. Title AGENT, TOTAL	TURF CONSULTING	LLC	# % - 4 # - 6 # - 7		
yped Name	7 1	5. Date			-		

3/10

Section II (Explanation Continued)

SUBJECT TO ENFORCEMENT ACTION AND PENALTIES UNDER SECTIONS 12 AND 14 OF FIFRA.







TURF FERTILIZER WITH RONSTAR® 0.67

50 lb. Bag Treats up to 7,260 Sq. Ft.

See supplemental label for the use of this product on ornamental shrubs, vines, and trees.

FOR SALE TO, USE & STORAGE BY TURF AND LANDSCAPE PERSONNEL ONLY. FOR PARKS, GOLF COURSES, ATHLETIC FIELDS, AND OTHER RECREATIONAL AND ORNAMENTAL TURFGRASS AREAS. A PRE-EMERGENT WEED CONTROL PLUS FERTILIZER FOR PERENNIAL BLUEGRASS, PERENNIAL RYEGRASS, ST. AUGUSTINEGRASS, BERMUDAGRASS, TALL FESCUE AND ZOYSIAGRASS

Controls: crabgrass, annual bluegrass, goosegrass, Florida Pusley, oxalis, stinging nettle, carpetgrass, pigweed and other weeds listed

ACTIVE INGREDIENT:

Oxadiazon: [2-tert-butyl-4-(2,4-dichloro-5-isopropoxyphenyl) x²-1,3,4-oxadiazolin-5-one]. 0.67% INERT INGREDIENTS: 99.33% Total. 100.00%

Contains RONSTAR® Herbicide a Registered Trademark of Bayer Environmental Science.

EPA Reg. No. 52287-3

EPA Est. No. 52287-FL-1

GUARANTEED FERTILIZER ANALYSIS

Derived from Ammonium Sulfate, Ureaform, Muriate of Potash, Manganese as Mn 0.5%, fron as Fe 1.0%, Chlorine < 15%

KEEP OUT OF REACH OF CHILDREN

WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detaile. (If you do not understand the label, find someone to explain it to you in detail.)

STATEMENTS OF PRACTICAL TREATMENT

<u>IF SWALLOWED</u>: Call a physician or poison control center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. <u>IF IN EYES</u>: Flush eyes with plenty of water. Get medical attention if irritation persists. <u>IF ON SKIN</u>; Wash with plenty of soap and water. Get medical attention if irritation persists.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes eye and skin irritation. Avoid contact with skin, eyes or clothing. Thoroughly wash face and hands with soep and water after exposure and before eating or smoking. Wear protective clothing including long-skeved shirt and gloves when handling. Launder exposed clothing before reuse. Do not allow domestic animals to graze treated areas. Avoid contamination of feed and foodstuffs.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertical areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not apply where rucoff is likely to occur. Do not contaminate water when disposing of equipment washwaters. Apply only as specified on this laber. Do not allow people (other than the applicator) or pets on treatment area during application. Do not enter the treatment area until dust has settled.



DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system. Not for use on home lawns. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE. Store product in its original bag, in a cool, dry, locked place out of reach of children. In case of spill, sweep up, and use according to directions or dispose of as follows.

PRODUCT DISPOSAL: If Empty: Do not reuse this container. Place in trash or offer for recycling if available. If Partly Filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

When applied according to directions, this product can be used safely on the following established turfgrass varieties: BERMUDAGRASS, PERENNIAL BLUEGRASS, PERENNIAL RYEGRASS Varieties, ST. AUGUSTINE GRASS Varieties, TALL FESCUE, and ZOYSIAGRASS lawns and ornamental turf. It may be used for Apartment and Condominium complexes, Parks, Golf Courses, Athletic Fields, Cemeteries and all other ornamental and recreational turfgrass areas.

CONTROLS GRASSY WEEDS AND BROADLEAF WEEDS: Annual Bluegrass (Poa Annua), Barnyardgrass, Crabgrass, Fall Panicum, Florida Pusley, Goosegrass, Green Foxtail, Oxalis, Stinging Nettle, Carpetgrass, Pigweed

HOW TO APPLY

For best weed control rake the area free of leaves and thatch and mow if necessary so that granules will reach the soil surface. Measure the area to be treated and uniformly apply the required amount using a fertilizer spreader (see Spreader Settings chart). Do not apply to turf when it is wet. Do not overlap spreader applications on St. Augustinegrass since discoloration "streaking" may temporarily occur from "double rate". After application if rain is not expected within 24 hours sprinkle the treated turf with ¼ to ½ inch of water to get maximum weed control.

HOW MUCH TO USE

FOR CRABGRASS: Apply 50 lbs./7,260 sq. ft. when turf has been in a yearly maintenance program (equal to 2 lbs. A.I. herbicide per acre). Apply 50 lbs. to treat 4,840 sq. ft. of lawn if it has been heavily infested with Crabgrass (equal to 3 lb. A.I. herbicide per acre). Apply in early spring 1 to 2 weeks prior to weed seed germination. This product is a pre-emergence herbicide. It will not control emerged and growing weeds; only germinating weed seeds. Late application will control only those weed seeds that germinate after the application. See "When To Apply" Chart.

FOR GOOSEGRASS AND PROSTRATE SPURGE: These weeds do not usually germinate until late spring or early summer. Apply 50 lbs. to treat 3,630 sq. ft. (4 lbs. A.I./acre) and apply 1 to 2 weeks prior to seed germination. See Use Precautions for St. Augustinegrass.

FOR ANNUAL BLUEGRASS (POA ANNUA): Apply 50 lbs. Per 3,630 to 4,170 sq. ft. of lawn (equal to 3.5 to 4 lbs. A.l. herbicide per acre). Apply in late summer to fall or in early spring 1 to 2 weeks before expected weed seed germination. Will not control annual bluegrass present at time of application, only germinating seeds. Refer to the chart for proper timing in your locale. Annual barnyardgrass and prostrate spurge may also be controlled by this same rate. See Use Precautions for St. Augustinegrass. FOR PIGWEED, FLORIDA PUSLEY, OXALIS, STINGING NETTLE AND CARPETGRASS: Where these weeds are a problem apply 50 lbs. to treat 3,630 to 4,174 sq. ft. of lawn (equal to 3.5 to 4 lbs. A.l. herbicide per acre) in early spring or 1 to 2 weeks before expected weed seed germination. Will not control emerged and growing weeds; only effective against germinating weed seeds. See Use Precautions for St. Augustinegrass.

SPREADER SETTINGS FOR SQUARE FOOT COVERAGE

SPREADER	AMT. OF PRODUCT	7,260 SQ. FT. =2#A,1./ACRE	4,840 SQ.FT. =3#A.I./ACRE	3,630 SQ.FT. ==4#A.I./ACRE
Cyclone	50 lbs.	XX	XX	XX
Scotts #35	50 lbs.	XX	XX	XX

NOTE: spreader rate varies with walking speed. Check spreader over known area.

2



WHEN TO APPLY

APPRORIATE TIME TO APPLY THIS PRE-EMERGENT HERBICIDE											
JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
1,				2,			3	3			
								3A			
4,				2Λ		=		4			

- 1. For CRABGRASS and other Grasses and Broadleaf Weeds germinating in the Spring in Northern turf.
- For GOOSEGRASS and other Late germinating Annual Grasses and broadleaf weeds such as PROSTRATE SPURGE.
- 2a. For GOOSEGRASS in Southern turf.
- 3. For POA ANNUA in Northern turf.
- 3a. For POA ANNUA in Southern turf.
- 4. For Fall-Winter-Spring control of annual weeds in Southern turf.

USE PRECAUTIONS

Do not use on Red Fescue, Bentgrass varieties, Dichondra, or Centipede grass because damage will occur. Do not use on putting greens or "Tees" due to varying cultural practices used. For St. Augustinegrass application rates above the 50 lbs. per 4,840 sq. ft. will temporarily discolor and damage the turf. If a higher rate is necessary (such as 50 lbs./ 3,630 sq. ft. or 50 lbs./ 4,174 sq. ft.) in order to control Goosegrass, Spurge, Poa Annua, Florida Pusley or Carpetgrass, turf damage can be expected to the St. Augustine. Therefore apply only when that damage can be tolerated and when good cultural practices are planned in order to revive the damaged turf. Avoid product contact with flowers and shrubs except as recommended on the ornamental use Supplemental Labeling available for this product. For newly seeded lawns do not apply this product until after the second or third cutting. For best lawn care, place new grass seed in the fall of the year. Spring pre-emergence weed control can then be applied.

RESEEDING TREATED AREAS

Delay new seeding until 4 months after treatment.

NOTICE: Buyer and user assume all risk and liability of use, storage, and/or handling of this product not in accordance with the terms of this label.

NET WEIGHT: XX POUNDS

Harrell's Inc. P. O. Box 807 Lakeland, FL 33802

. . .