



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

May 9, 2022

Sanford R Simon
VP Corporate Compliance
Harrell's LLC.
P.O. Box 807
Lakeland, FL 33802

Subject: Registration Review Label Mitigation for Prodiamine
Product Name: TURF FERTILIZER WITH BARRICADE(R) 0.21
EPA Registration Number: 52287-6
Application Date: July 10, 2019
Decision Number: 553020

Dear Mr. Simon:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Prodiamine Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Page 2 of 2
EPA Reg. No. 52287-6
Decision No. 553020

If you have any questions about this letter, please contact Srijana Shrestha by phone at 202-566-2329, or via email at shrestha.srijana@epa.gov.

Sincerely,

A handwritten signature in blue ink, appearing to be 'Linda Arrington', with a stylized flourish at the end.

Linda Arrington, Branch Chief
Risk Management and Implementation Branch 4
Pesticide Re-Evaluation Division
Office of Pesticide Programs

Enclosure: Stamped Label

Bold, italicized text is for the reader and is not a part of the label.

[Bracketed information is optional] Text separated by / denotes and/or options



PRODIAMINE	GROUP	3	HERBICIDE
------------	-------	----------	-----------

TURF FERTILIZER WITH BARRICADE® 0.21

[Alternate Brand Names: Fertilizer W/ Barricade® 0.21%; Harrell’s PROtect Professional Fertilizer plus Barricade 0.21; Turf & Ornamental Fertilizer W/ Barricade® 0.21%; Landscape Ornamental Fertilizer W/ Barricade® 0.21%; Fertilizer W/ Prodiamine 0.21]

[Batch Number] [SKU Number] [Fertilizer N-P-K]

FOR PRE-EMERGENCE CONTROL OF WEED GRASSES AND BROADLEAF WEEDS IN ESTABLISHED TURF GRASSES, LAWNS, GOLF COURSE FAIRWAYS AND SOD FARMS, (NOT FOR PUTTING GREENS). USE FOR PRE-EMERGENCE WEED CONTROL AROUND LANDSCAPE ORNAMENTALS INCLUDING ESTABLISHED PERENNIALS AND WILDFLOWER PLANTINGS.

ACTIVE INGREDIENT:

Prodiamine:*** 0.21%

OTHER INGREDIENTS:..... 99.79%


Total..... 100.00%

***CAS 29091-21-2

(Note to PM- CAS number may appear on the front or back panel)

EPA Reg. No. 52287-6

EPA Est. No. 52287-FL-1



KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

READ ENTIRE LABEL BEFORE USING

NET WEIGHT: 50 lb (22.7 kg) [or: 1,000 lb, or 2,000 lb or bulk trailer]

Bold, italicized text is for the reader and is not a part of the label.

[Bracketed information is optional] Text separated by / denotes and/or options

FIRST AID

If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice • Have person sip a glass of water if able to swallow • Do not induce vomiting unless told to do so by a poison control center or doctor.
If in eyes:	<ul style="list-style-type: none"> • Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
If on skin:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor of going for treatment.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequent repeated skin contact while handling the material may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long sleeved shirt and long pants, chemical resistant gloves, shoes plus socks and a dust mask. Follow manufacturer’s instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then was thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish.

AQUATIC ORGANISM ADVISORY STATEMENT: Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated area may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment wash water.

Bold, italicized text is for the reader and is not a part of the label.

[Bracketed information is optional] Text separated by / denotes and/or options

NON-TARGET ORGANISM ADVISORY STATEMENT: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. This product is a turf fertilizer and a preemergence herbicide that provides residual control of many grass and broadleaf weeds in established turf grasses, golf courses (excluding putting greens), for sod farms, and lawns, and around landscape ornamentals, established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seed germination and root development.

Do not graze or feed livestock forage cut from areas treated with this product. Do not apply aerially. Do not apply to golf course putting greens. Failure to follow the directions for use and precautions on this label may result in poor weed control, plant injury.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (sod farms, commercial nurseries).

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls; chemical resistant gloves such as butyl rubber > 14 mils or natural rubber > 14 mils or nitrile rubber > 14 mils; shoes plus socks.

NON-AGRUCULTURAL USES REQUIREMENTS

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The "WPS" applies when this product is used to produce agricultural plants on farms, forests or nurseries. Applicators and handlers should wear long-sleeved shirt, long pants, chemical resistant gloves, shoes with socks and protective eyewear. Thoroughly wash face and hands with soap and water after exposure and before eating or smoking. Launder exposed clothing before reuse. Do not allow reentry to treated areas until dusts have settled and the turfgrass or soil is dry.

Bold, italicized text is for the reader and is not a part of the label.

[Bracketed information is optional] Text separated by / denotes and/or options

WEED-RESISTANCE MANAGEMENT

For resistance management, Turf Fertilizer with Barricade® 0.21 contains the active ingredient Prodiamine, a Group 3 Herbicide as determined by the mode of action (MOA) classification system of the Weed Science Society of America. Any weed population may contain or develop plants naturally resistant to Fertilizer with Turf Fertilizer with Barricade® 0.21 Group 3 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed. Continued effectiveness of this product depends on the successful implementation of a weed resistance management program.

To delay herbicide resistance users should take one or more of the following steps:

- Rotate the use of Turf Fertilizer with Barricade® 0.21 and other Group 3 herbicides within a growing season sequence, or among growing seasons with different herbicide groups that control the same weeds in a field.
- Adopt an integrated weed-management program for herbicide use that includes:
 - Scouting and uses historical information related to herbicide use and crop rotation, and
 - Consideration of tillage (or other mechanical control methods), and
 - Cultural (e.g., higher crop seeding rates), and
 - Precision fertilizer application method and timing to favor the crop and not the weeds), and
 - Biological (weed-competitive crops or varieties) and other management practices.
- Scout fields prior to application to identify the weed species present and their growth stage to determine if the intended application will be effective.
- Scout fields after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include:
 - Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds, or
 - A spreading patch of non-controlled plants of a particular weed species or,
 - Surviving plants mixed with controlled individuals of the same species.
- If resistance is suspected, prevent weed seed production in the affected area by using an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- Report any incidence of local non-performance of this product against a particular weed species to your [Harrell's] [company] representative at (800) 282-8007, [and/or at <https://www.harrells.com>].

BEST MANAGEMENT PRACTICES

Proactively develop and implement diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides. A diversified program may include the use of herbicides

Bold, italicized text is for the reader and is not a part of the label.

[Bracketed information is optional] Text separated by / denotes and/or options

with different modes of action along with cultural, mechanical, and/or biological practices. Best management practices include the following:

- To the extent possible, use a diversified approach toward weed management. Whenever possible incorporate multiple weed-control practices such as mechanical cultivation, biological management practices, and crop rotation.
- Plant into weed-free fields and keep fields as weed-free as possible.
- Identify weeds present in the field through scouting and field history and understand their biology. The weed-control program should consider all the weeds present. Apply this herbicide at the correct timing and rate needed to control the most difficult weed in the field.
- Fields with difficult to control weeds should be rotated to crops that allow the use of herbicides with alternative modes of action (MOA) or different management practices.
- Difficult to control weeds may require sequential applications of herbicides with differing MOA's.
- Prevent field-to-field and within-field movement of weed seed or vegetative propagules. Thoroughly clean plant residues from equipment before leaving fields.
- To the extent possible do not allow weed escapes to produce seeds, roots or tubers. Manage weed seeds at harvest and post-harvest to prevent a buildup of the seed seedbank.
- If resistance is suspected, treat weed escapes with a herbicide with a different MOA than those in this product or use non-chemical methods to remove escapes.

DRIFT ADVISORY

WIND: Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

APPLICATION DIRECTIONS

Apply uniformly with suitable, calibrated application equipment. The size and bulk density of granules in this product may vary; therefore, calibrate your granular applicator prior to application. A calibrated applicator must be used. Follow directions in the owner's manual for calibrating your applicator.

APPLICATION RATES

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds. The length of time of residual pre-emergence weed control provided by this product is related to the rate applied. Maximum use rate selection for turf should be based on turf species.

Bold, italicized text is for the reader and is not a part of the label.

[Bracketed information is optional] Text separated by / denotes and/or options

WEEDS CONTROLLED

When used in accordance with this label this product will provide control of the following weeds:		
Barnyardgrass	Henbit	Purslane, Common
Bluegrass, Annual (<i>Poa Annua</i>)	Itchgrass	Pusley, Florida
Carpetweed	Johnsongrass (from seed)	Rescuegrass ⁽³⁾
Chickweed, Common	Junglerice	Shepherd’s Purse ⁽²⁾
Chickweed, Mouseear (from seed)	Knotweed	Signalgrass, Broadleaf
Crabgrass (Large, smooth)	Kochia	Speedwell, Persian
Crowfootgrass	Lambsquarter, Common	Sprangletop
Cupgrass, Woolly	Lovegrass	Spurge, Prostrate
Foxtails, Annual	Panicum (Texas, Fall, Browntop)	Witchgrass
Goosegrass ⁽¹⁾	Pigweed	Woodsorrel, Yellow (from seed)

⁽¹⁾ In many areas a single application of 310 to 715 lbs. per acre will control Goosegrass. However, under heavy Goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 310 to 475 lbs. per acre followed after 60 to 90 days by a second application at doses that would not exceed those given in the Maximum Annual Application Rate section. Do not exceed the maximum rates in the Maximum Annual Application Rate section.

⁽²⁾ Application for Shepherd’s Purse should be made in late summer, fall or winter prior to germination.

⁽³⁾ Suppression only of Rescuegrass.

Sequential application may be made so long as the total amount of product applied does not exceed the maximum annual application rates. All applications must be made prior to germination of the weed seeds.

USE RESTRICTIONS

- Do not apply this product to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*poa annua*) are desirable species
- Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.
- Do not cut (harvest) treated sod before 120 days after application. Do not apply to newly set sod until the following year.

USE PRECAUTIONS

- Application of this product may thin emerged annual bluegrass and newly overseeded grasses.
- Application to turf stressed by drought, low fertility, or pest damage may result in turf injury.
- Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.

Bold, italicized text is for the reader and is not a part of the label.

[Bracketed information is optional] Text separated by / denotes and/or options

MAXIMUM ANNUAL APPLICATION RATE

This product is for use on the turf grass species listed below and on landscape ornamental species as listed in the Tolerant Ornamental Species section of this label. Do not exceed the maximum annual rate as stated in the following table:

TURF GRASS		
MAXIMUM ANNUAL APPLICATION RATE BY TURF GRASS SPECIES ⁽¹⁾		
Turf Species	Lbs. product/A	Lbs. ai/A
Creeping Bentgrass	310 lbs.	0.65 lbs.
Creeping Red Fescue	355 lbs.	0.75 lbs.
Buffalograss	475 lbs.	1.00 lbs.
Kentucky Bluegrass	475 lbs.	1.00 lbs.
Perennial Ryegrass	475 lbs.	1.00 lbs.
Bermudagrass ⁽²⁾	715 lbs.	1.50 lbs.
Bahiagrass	715 lbs.	1.50 lbs.
Centipedegrass	715 lbs.	1.50 lbs.
Seashore Paspalum	715 lbs.	1.50 lbs.
St. Augustinegrass	715 lbs.	1.50 lbs.
Tall Fescue (including turf type)	715 lbs.	1.50 lbs.
Zoysia	715 lbs.	1.50 lbs.
MAXIMUM ANNUAL APPLICATION RATE FOR LANDSCAPE ORNAMENTALS		
Tolerant Ornamental Species	715 lbs.	1.50 lbs.

(1) These are the maximum rates per calendar year by species limitations.

(2) May be used on newly-sprigged or plugged bermudagrass at rates not to exceed 248 lbs. per acre. Newly-sprigged or plugged bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced product rates used in sprigging situations. Do not apply more than 715 lbs. of this product to the same area per calendar year. Use higher rates of product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum application rates specified in the Maximum Annual Application Rate section.

ESTABLISHED TURF

This product is a pre-emergence herbicide that, when properly applied, will control certain grass and broadleaf weeds listed in the Weeds Controlled Table in established turf grass and lawns. The maximum amount of material that may be applied per year is given for each turf grass species in the Maximum Annual Application Rate table. Most effective weed control in turf grasses will be obtained when this product is activated by at least ½ inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map for approximate crabgrass seed germination dates.

Bold, italicized text is for the reader and is not a part of the label.

[Bracketed information is optional] Text separated by / denotes and/or options



- 1: Mid-north & northwestern.....After May 30
- 2: Upper/mid-states & northeast.....After May 10
- 3: Lower mid-states.....After April 20
- 4: Upper southern states.....After March 20
- 5: Warm season states.....January 1 to March 20

WHEN TO APPLY AFTER OVERSEEDING TURF

Delay seeding for 60 days after application or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding and/or reseeding results.

Months Before Overseeding

Product Rate Lbs./A	Lbs. ai/Acre	North*	Transition*	South*
240	0.50	4	4	4
310	0.65	5	4	4
355	0.75	6	5	5
380	0.80	-	6	6
475	1.00	-	7	7
540	1.14	-	-	9
620	1.30	-	-	10
715	1.50	-	-	12

*Transition states are: DE, MD, VA, WV, KY, MO, KS; North states are those north of transition states; South states are those south of transition states. States west of KS, should follow timing for North or South depending on turf type: Cool Season or Warm Season.

LANDSCAPE ORNAMENTS INCLUDING ESTABLISHED PERENNIALS AND WILDFLOWER PLANTINGS

This product may be applied for residual pre-emergence weed control around ornamentals and wildflowers listed on this label.

Bold, italicized text is for the reader and is not a part of the label.

[Bracketed information is optional] Text separated by / denotes and/or options

USE RATES

Apply at 310 to 715 lbs. per acre of landscape plants, in fall and /or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 715 lbs. per acre.

APPLICATION TIMING AND INFORMATION

Apply uniformly to the surface of the soil in ornamental planting beds with a granular applicator. This product is a pre-emergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least ½ inches of rainfall or irrigation, or with shallow (1 to 2 inches) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves. Prior to application of this product, control existing vegetation by hand weeding, cultivation, or the use of an appropriate post-emergence herbicide.

ESTABLISHED LANDSCAPE ORNAMENTALS

This product may be applied to established ornamentals as a broadcast, over the top, or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate this product into the soil.

NEWLY TRANSPLANTED ORNAMENTALS

This product may be applied to newly transplanted ornamentals as a broadcast, over the top, or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply after cuttings from roots have become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union. Irrigate after application as above directed.

TOLERANT ORNAMENTAL SPECIES

This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to this product.

FOR USE AROUND LANDSCAPE ORNAMENTALS, INCLUDING ESTABLISHED PERENNIALS AND WILDFLOWERS, AS LISTED.

Scientific Name	Common name	Scientific Name	Common name
Abies spp.	Fir Species**	Gypsophila repens	Baby’s Breath
Abelia grandiflora	Abelia: Sherwood	Hedera helix	English Ivy
Acer palmatum	Japanese Maple	Helianthemum sp.	Sunrose
A. platanoides	Norway Maple	Hemerocallis sp.	Daylily: Aztec Gold, Stella De Oro, Tender Love
Achillea, sp.	Yarrow: King Edward	Hibiscus	Rose of Sharon**
Actinidia chinensis*	Kiwi*	Hibiscus sp.	Mallow: Disco Belle White
Agapanthus africanus	Lily-of-the-Nile (African Lily)	Hibiscus rosa-sinensis	Chinese Hibiscus
Agapanthus orientalis		Hosta plantaginea	Hosta, Plantain Lily (Fragrant)
Akebia quintata	Five-Leaf or Chocolate Vine	H. sieboldiana	Hosta, “Searsucker”
Allium cemuum	Lady’s Leek, Nodding Onion	Houttuynia cordata var. Variegata	
Anemone hybrid	Japanese Anemone	Hydrangea macrophylla	Bigleaf hydrandea

Bold, italicized text is for the reader and is not a part of the label.

[Bracketed information is optional] Text separated by / denotes and/or options

Aquilegia sp.	Aquilegia: Red and Gold	Ilex comuta	Chinese Holly
Arctostaphylos densiflora	Vine Hill Manzanita	I. crenata	Japanese Holly; Helli
Arctotheca calendula	Cape Weed	I. opacalis	American Holly
Artemisia sp.	Wormwood: Silver Mound, Castle	I. pernyi	Holly
Aster sp.	Aster: Bonny Blue, Purple Dome	I. vomitoria	Yaupon Holly; Schillings
Aster X Frikartii		Inula ensifolia	
Athrium Filiz-femina	Lady Fern: Fern Lady	Iris ensata	Sword-Leaved iris; Jodlesong
Aucuba japonica	Japanese Aucuba	Iris siberica	Siberian Iris; Cabernet Iris species**
Begonia sp.	Fibrous Begonia: Hardy Grandis	Iris spp.	Iris species**
Berberis gladwynensis	Barberry	Jasminium nudiflorum	Winter Jasmine
B. julianae	Wintergreen Barberry	Juniperus chinensis	Chinese Juniper; Nick's Compact, Parsonii
B. mentorensis	Mentor Barberry	J. conferta	Shore Juniper; Blue Pacific
B. thunbergii	Japanese Barberry	J. davurica	Parsoni
B. verruculosa	Warty Barberry	J. horizontalis	Creeping Jupiter
Bergenia cordifolia		Juglans sp.	Walnut*
Boltonia asteroides	Snowbank	Justicia brandegeana	Shrimp Plant
Bougainvillea spp.	Bougainvillea	Lagerstromia indica	Crape Myrtle
Buddleia davidii	Butterfly-Bush (Dwarf Blue); Royal Red	Lagerstromia indica X Fauriei	Crape Myrtle; Tuscarora
Buxus microphyllia	Japanese Boxwood	Lantana montevidensis	Weeping Lantana
Callistermon viminalis	Weeping Bottlebrush	Lavender sp.	Lavender; Munstead
Calluna vulgaris	Scotch Heather	Leontopodium alpinum	Edelweiss
Campanula carpatica	Tussock Bellflower (white clips)	Ligstrum sinense	Chinese Privet; Variegata
Campis X Tagliabuana	Trumpet Creeper, Trumpet Flower, Madame Galen	L. amurense	Amur Privet
Carpobrotus edulis	Hottentot Fig (Ice Plant)	L. japonicum	Japanese Privet
Cassia artemisoides	Feathery Cassia	L. lucidum	Glossy Privet (Waxleaf)
Ceanothus rigidus	Wild Lilac	Lillum sp.	Lily; Jazz
Ceratostigma Plumbagonoides		Liriope muscari	Big Blue Lily; Turf; Silver Mound; Evergreen Giant
Chamaecyparis pisifera	False Cypress	Liriope Muscari var. Variegata	Liriope, Variegated
Chrysanthemum nipponicum		Liriope spicata	Liriope, Creeping
Cleyera japonica	Cleyera	Lobelia Cardinalis	Cardinal Flower, Indian Pink
Citrus spp.*	Citrus species*	Lonicera japonica	Japanese Honeysuckle
Coreopsis sp.	Coreopsis (Calliopsis): Early Sunrise, Moonbeam	L. tatarica	Tatarian Honeysuckle
Cornus florida	Flowering Dogwood	Loropetalum Chinese	Burgundy
C. stolonifera	American Dogwood	Lythrum sp.	Loosestrife; Modern Pink
Cortaderia seloana	Pampas Grass	Magnolia spp.	Magnolia species**
Cotonneaster apiculatus	Cranberry Cotoneaster	Maleophora luteola	Ice Plant
C. buxifolius	Cotoneaster	Malus sp.	Crabapple*
C. dammeri	Bearberry Cotoneaster	Miscanthus sinensis	Yaku Jima, Silberfeder
C. microphyllus	Rockspray Cotoneaster	Nandina domestica	Heavenly Bamboo
Cretaegus spp.	Hawthorne	Narcissus spp.	Narcissus species**
Crocsmia sp.	Lucifer	Nerium oleander	Oleander
Cupressus sempervirens	Italian Cypress	Olea europea	Olive*
Delosperma alba	White Trailing Ice Plant	Ophiopogon japonicas	Mondo Grass
Delosperma sp.	Cooperi Pink	Osmanthus heterphyllus	Osmanthus (False Holly); Gulf tide
Delphinium sp.	Larkspur: Blue Elf	Osteospermum fruticosum	Trailing African Daisy
Dianthus deltoids	Dianthus: Maiden Pinks "Zing"	Oxydendrum arboretum	Sourwood
D. gratianopolitanus	Cheddar Pink	Paeonia suffruticosa	Tree Peony
Dodonea Viscosa	Hop Bush	Pennisetum setaceumi	Fountain Grass (Dwarf)

Bold, italicized text is for the reader and is not a part of the label.

[Bracketed information is optional] Text separated by / denotes and/or options

Scientific Name	Common Name	Scientific Name	Common Name
Echinacea purpurea	Coneflower, Purple; Mangus	Perovskia atriplicifolia	
Elaeagnus pungens	Silverberry	Persea Americana	Avocado*
Euonymus fortune	Wintercreeper	Photinia fraseri	Frasier's Photinia ((Redtip)
E. japonica	Japanese Spindle Tree (Evergreen Euonymus)	Physotegia Viginiana	Dragonhead, False; Vivid Spruce species**(Colorado Blue, Norway, etc.)
E. kiautschovica	Spreading Euonymus	Pieris japonica	Lily-of-the-Valley Bush
Fatsia japonica	Japanese Aralia	Pinus brutia	Calabrian Pine
Forsythia intermedia	Border Forsythia	P. canariensis	Canary island Pine
F. suspense	Weeping Forsythia	P. elliotii	Slash Pine
F. viridissima	Greenstem Forsythia	P. halepensis	Aleppo Pine
Gaillardia sp.	Gaillardia, Blanket Flower: "Goblin"	P. nigra	Austrian Pine
Gardenia jasminoides	Gardenia, Cape-Jasmine, Radicans Gaur asp.	P. palustris	Longleaf Pine
Gentiana daturica	Gentian	P. radiata	Monterey Pine
Geranium cinereum	Cranesbill	P. thunbergiana	Japanese Black Pine
Gladiolus spp.	Gladiolus species**	P. strobus	Eastern White Pine
P. sylvestris	Scotch Pine	Syzygium paniculatum	Australian Brushcherry
P. taeda	Loblolly Pine	Taxus cuspidate	Japanese Yew
P. virginiana	Virginia Pine	T. media	Yew
Pistachio sp.	Pistachio*	Teucrium sp.	Germander
Pittosporum rhombifolium	Queensland Pittosporum	Thalictrum dipterocarpum	Meadow Rue
P. tobira	Japanese Pittosporum	Thuja occidentalis	American Arborvitae
Posocarpus Macrophyllus	Japanese Yew	Trachelospermum asiaticum	Star Jasmine
Prunus fauocerasus	English Laurel	Tsuga Canadensis	Canada Hemlock
Prunus sp.	Almond, Apricot, Nectarine, Peach, Plum and Prune*	Tulipa spp.	Tulip species
Pseudotsuga menziesii	Douglas Fir	Veronica sp.	Veronica, Speedwell; Sunny Border
Pyracantha Coccinea	Firethorn (Scarlet)	Viburnum japonicum	Japanese Viburnum
P. fortuneana	Firethorn	V. odoratisimum	Sweet Viburnum
P. koidzumii	Firethorn	V. plicatum	Japanese Snowball
Pyrus Sp.	Bradford Pear sp.	V. rigidum	Canary Island Viburnum
Quercus Shumardii	Oak, Shumard's Red	V. suspensum	Arrowwood Viburnum
Quercus spp.	Oak species	V. tinus	Laurustinus
Raphiolepis indica	Indian Hawthorne	V. trilobium	Cranberry Bush
Raphiolepis umbellate	Yedda Hawthorn	V. wrightii	Leatherleaf Viburnum
Rhododendron (including Azalea)	'Coro Bells' 'Delaware, Valley White', 'Flame Creeper', 'formosa', 'Girard Crimson', George L. Tabor', Hino-crimson', 'PJM', 'Roseum Elegans', 'Wakeiebisu', 'White Gumbo'	Vinca major	Greater Periwinkle, Vinca
Rosa banksiae	Lady Bank's Rose	Vinca minor	Common Periwinkle, Vinca
Rosmarinus officinalis	Rosemary*	Vitis Sp.	Grape*
Rudbeckia sp.	Black-Eyed susan; Goldstrum	Weigela florida	Old Fashioned Weigela
Santolina, virens	Santolina	Yucca aloifolia	Spanish Bayonet, Yucca
Saxifraga sp.	Axifrage; Purple Dome	Y. filaentosa	Adam's Needle, Yucca
Scabiosa sp.	Pincushion Flower		
Sedum album	Stonecrop		
Sedum cauticola	Stonecrop; Lidakense		
Sedum dasphyllum	Stonecrop		
Sedum spurium	Sonecrop; Dragon's Blood		
Spiraea Bumalda	Spirea; Anthony Waterer		

*Do not use on food producing trees, vines or plants.

**Do not use on container grown plants.

Bold, italicized text is for the reader and is not a part of the label.

[Bracketed information is optional] Text separated by / denotes and/or options

STORAGE AND DISPOSAL

PESTICIDE STORAGE

Do not store above 120°F. Store in a cool, dry, and locked storage area inaccessible to children and domestic animals. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING

For 50, 1000, or 2000-pound bags and totes:

Non-refillable container. Do not reuse or refill this container, offer for recycling if available. Completely empty container into application equipment. Dispose of empty container in a sanitary land fill or by other procedures allowed by state and local authorities.

For Killebrews and other bulk transport vehicles:

Refillable container: Container handling instructions are not required. Cleaning before refilling is the responsibility of the refiller.

(Note: End Directions for Use)

[Turf Fertilizer with Barricade® 0.21]

Guaranteed [Fertilizer] Analysis [N-P-K]

Total Nitrogen (N).....XX%

 XX % Ammoniacal Nitrogen

 XX % Urea Nitrogen*

Available Phosphate(P₂O₅).....XX%

Soluble Potash (K₂O).....XX%

Derived from: e.g.: Polymer-Coated Urea, Urea, Ammonium Phosphate, Muriate of Potash.

*X% Slowly Available/Slow Release/Controlled release nitrogen from polymer-coated urea.

[Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco/metals.org>]

F-352 (Florida Fertilizer License Number)

[Fertilizer] [Density] [Bulk Density]

[Any other state specific fertilizer required language]

(Note to PM: Appropriate fertilizer specific information will be inserted on Final Printed Label to replace "X".)

Bold, italicized text is for the reader and is not a part of the label.

[Bracketed information is optional] Text separated by / denotes and/or options

CONDITIONS OF SALE AND WARRANTY

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitation of Liability before using this product. If the terms are not acceptable, return the unopened container at once. By using this product, user and buyer accept the following Conditions, Disclaimer of Warranties, and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Harrell's, Inc. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: There are no warranties express or implied, of merchantability or of fitness for a particular purpose or otherwise, which extend beyond the statements made on this label. No agent of Harrell's, LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Harrell's, LLC disclaims any liability whatsoever for special, incidental, or consequential damages resulting from the use or handling of this product.

LIMITATION OF WARRANTIES: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price, or at Harrell's, LLC option, the replacement of the product.

Manufactured By/Manufactured For/Sold By/ [and Guaranteed By]

Harrell's, LLC
P. O. Box 807
Lakeland, FL 33802

[***CAS 29091-21-2]

(Note to PM: CAS No. may appear on the front or back panel)

[Growing a Better World® is a registered trademark of Harrell's, LLC]

[BARRICADE® IS A REGISTERED TRADEMARK OF SYNGENTA, INC]