07/24/2009



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

JUL 2 4 2009

ş

Mr. Robert F. Bischoff Product Registration TotalTurf Consulting LLC Agent for Harrell's LLC 26 A.A. Cooper Road Megs, GA 31765

SUBJECT: Application for Pesticide Notification (PRN 98-10) Request Alternate Brand Name "Fertilizer W/Barricade 0.45%" EPA Reg. No. 52287-7 Application Dated June 17, 2009

Dear Registrant:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 06/17/09 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

 \sim

Linda Arrington Notifications & Minor Formulations Team Leader Registration Division (7505P) Office of Pesticide Programs

		mpleting form.	Form A	boroved.	OMB No. 207	0-0060, Approval expires 2-28-95
\$EPA	Environme	United States Ntal Protection A /eshington, DC 20460			Registratio Amendme Other	OPP Identifier Number
		Application fo	or Pesticide - Se	ction I		
I. Company/Product Nu 52287-7	mber		2. EPA Product M Joanne I. Miller	-		3. Proposed Classification
. Company/Product (Na Turf Fertilizer with E			РМ# 23		_	
Ame and Address of Harrell's LLC P. O. Box 807 Lakeland, FL 3380 Check in			-	t is simil. N/A	lar or identical	with FIFRA Section 3(c)(3) in composition and labeling OTIFICATION JUL 2 4 2009
		S	ection - II			
Notification - Exp xplanation: Use edd Notification of Alternate CFR 152.46, and no oth 18 U.S.C. Sec. 1001 to b	response to Agency lo plain below. ditional page(s) if nece Brand Name per PR No per changes have been n willfully make any false	ossary. (For section I and otice 98-10 (IIA). This notif made to the labeling or the	Agency I "Me Too "Me Too Other - E Section II.) fication is consistent with confidential statement of ar understand that if this i	etter date " Applicat xplain bel the provise of formula on hotification	tion. ow. sions of PR Notic of this product. I is not consisten	e 98-10 and EPA regulations at 40 I understand that it is a violation of It with the terms of PR Notice 98-10
· · · · · · · · · · · · · · · · · · ·			action - III ter Soluble Packaging		· · · · · · · · · · · · · · · · · · ·	
child-Resistant Packagir Yes V No Certification mus	ng Unit Packaging Yes √ No	Wa ✓ No.per If "	ter Soluble Packaging	r	2. Type of Con	
Child-Resistant Packagir Yes Vo Certification mus Certification mus De submitted	ng Unit Packaging Yes Vo If "Yes" Unit Packaging ents Information Container	No. per wgt. container Pac 4. Size(s) Retail Cor	ter Soluble Packaging Yes No Yes" No. pe skage wgt contain Intainer 50 lbs	r ner	2. Type of Con	ntainer letal lastic lass aper ther (Specify) birections
Hild-Resistant Packagir Yes √ No Certification mus Se submitted Label	ng Unit Packaging Yes Vo If "Yes" Unit Packaging ents Information Container	No. per wgt. container 4. Size(s) Retail Cor t Lithograph Paper glued Stenciled	ter Soluble Packaging Yes No Yes" No. pe kage wgt contain ntainer 50 lbs	r ner 5. Loc	2. Type of Con M V Pl V Pi G G Pi O O Stion of Label C	ateiner letal lastic lass aper ther (Specify) birections
hild-Resistant Packagir Yes √ No Certification mus to submitted Location of Net Conte √ Label	ng Unit Packaging Yes √ No If "Yes" Unit Packaging ents Information Container el is Affixed to Produc	No. per wgt. container 4. Size(s) Retail Cor t Lithograph Paper glued Stenciled	ter Soluble Packaging Yes No Yes" No. pe contain ntainer 50 lbs Other Ction - IV	r ner 5. Loc -/	2. Type of Con ✓ Pl ✓ Pd ✓ Co Stion of Label C ☐ On Affixed L	ntainer letal lastic lass aper ther (Specify) Directions _abel
Child-Resistant Packagir Yes No Certification mus Certification mus Certification for the Conte Label [ng Unit Packaging Yes √ No If "Yes" Unit Packaging ents Information Container el is Affixed to Produc	No. per wgt. container If " wgt. container Pac 4. Size(s) Retail Cor 4. Size(s) Retail Cor 20 4. Size(s) Retail Cor 21 22 23 24 24 25 25 25 25 26 26 26 26 26 26 26 26 26 26 26 26 26	ter Soluble Packaging Yes No Yes" No. pe contain ntainer 50 lbs Other Ction - IV	ner 5. Loc her d, if nece	2. Type of Con ✓ Pl ✓ Pd ✓ Co etion of Label C On Affixed L ssary, to process Tel	ntainer letal lastic lass aper ther (Specify) Directions _abel
hild-Resistant Packagir Yes √ No Certification must e submitted Location of Net Conte ✓ Label [Ing Unit Packaging Yes No If "Yes" Unit Packaging ents Information Container el is Affixed to Produc elete items directly belo tatements I have made at any knowlingtly false	No. per wgt. container If " wgt. container Pac 4. Size(s) Retail Cor 4. Size(s) Retail Cor 20 4. Size(s) Retail Cor 21 22 23 24 24 25 25 25 25 26 26 26 26 26 26 26 26 26 26 26 26 26	ter Soluble Packaging Yes No Yes" No. pe skage wgt contain ntainer 50 lbs Ction - IV ection - IV dividual to be contacte	r ner 5. Loc 5. Loc d, if neces	2. Type of Con Pi G G Pi O ation of Label C On Affixed L Ssary, to process Tel (31 (31) (3) (3) (3) (3) (3) (3) (3) (3	Iteiner Iteiner Iteial Iassic Iass Iaper ther (Specify) Directions Iabel Inter Specify) Solutions Iabel Inter Specify Inter Specify
✓ No Certification must be submitted b. Location of Net Conte ✓ Label ✓ Label ✓ Label I. Contact Point (Complame Robert F. Bischoff I certify that the s Lacknowledge tha both under applica . Signature	Ing Unit Packaging Yes No If "Yes" Unit Packaging ents Information Container el is Affixed to Produc elete items directly belo tatements I have made at any knowlingtly false	No. per No. per wgt. container 4. Size(s) Retail Cor 4. Size(s) Retail Cor 4. Size(s) Retail Cor Paper glued Stenciled Sec Sec Sec Sec Sec Sec Sec Sec	ter Soluble Packaging Yes No Yes" No. pe contain intainer 50 lbs Oth Continuer 50 lbs Oth Continuer 50 lbs Oth Container 50 lbs Container 50 lbs Containe	d, if neces	2. Type of Con Pl G G G C Pl O C etion of Label C O On Affixed L Ssary, to process Tel (31	Ateiner letal lastic lass aper ther (Specify) Directions .abel ess this application.) ephone No. (Include Area Code) 17) 873-0222 6. Dete Application Received

EPA Form 8570-1 (Rev. 3-94) Previous editions are obsolete.

.

TotalTurf Consulting LLC 26 A. A. Cooper Rd. Meigs, GA 31765

Direct: 843-384-2807 Fax: 229-683-3426 E-Mail: rbcooper@juno.com

June 17, 2009

产

Document Processing Desk (NOTIF) Office of Pesticide Programs (7505P) U. S. Environmental Protection Agency Room S-4900, One Potomac Yard 2777 South Crystal Drive Arlington, VA 22202-4501

Attn: Joanne I. Miller

Subject: Turf Fertilizer with Barricade 0.45 EPA Reg. No. 52287-7 Notification of Alternate Brand Name

Dear Ms. Miller:

Enclosed labeling shows the addition of an alternate brand name (Fertilizer W/Barricade 0.45%) for the subject registered product. This notification is being submitted on behalf of Harrell's LLC, P. O. Box 807, Lakeland, FL 33802 by TotalTurf Consulting LLC, an authorized agent for Harrell's LLC.

The following documents are included in the subject submission:

Cover Letter Application for Pesticide (EPA Form No. 8570-1) One Copy of Labeling Showing the Alternate Brand Name [Coded as 20090617.Final Label] One Copy of Current Labeling [Coded as 20090303.Final Label (PRN2007-4Corr.)]

If there are any questions or additional information is needed concerning this submission, do not hesitate to contact me at (317) 873-0222 or Raymond B. Cooper at (843) 384-2807.

Sincerely,

Porbert F. Buchoff

Robert F. Bischoff Agent TotalTurf Consulting LLC 26 A. A. Cooper Road Meigs, GA 31765



惷

20090617.Final Label

.

4/10 NOTIFICATION JUL 2 4 2009

. . .

FERTILIZER W/BARRICADE® 0.45%

FOR SELECTIVE PRE-EMERGENCE CONTROL OF WEED GRASSES AND BROADLEAF WEEDS IN ALL ESTABLISHED TURF GRASSES, LAWNS, GOLF COURSE FAIRWAYS AND SOD FARMS, (NOT FOR PUTTING GREENS). USE FOR PRE-EMERGENCE WEED CONTROL AROUND LANDSCAPE ORNAMENTALS INCLUDING ESTABLISH PERENNIALS AND WILDFLOWER PLANTINGS.

	Total100.00%
INERT INGREDIENTS:	
Prodiamine***:	
ACTIVE INGREDIENT:	

***CAS [2909-21-2]

Barricade® is a registered trademark of Syngenta.Inc.

EPA Reg. No. 52278-7

EPA Est. No. 52287-FL-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detaile. (If you do not understand the label, find someone to explain it to you in detail.)

READ ENTIRE LABEL BEFORE USING

FIRST AID

<u>IF SWALLOWED:</u> Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, and then continue rinsing. Call a poison control center or doctor for treatment advice. IF ON SKIN: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice.

Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment. For information, call 1-800-282-8007.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

<u>CAUTION:</u> Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequent repeated skin contact while handling the material may cause allergic reactions in some individuals.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated area risy t = hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long sleeved shirt and long pants; chemical resistant gloves; shoes plus 3.xku and a dust mask. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

1

USER SAFETY RECOMMENDATIONS

<u>Users should</u>: Wash hands before eating, drinking, chewing gym, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. <u>STORAGE</u>: Store in original container away from other fertilizer, feed or foodstuffs and separated from other pesticides. <u>PESTICIDE DISPOSAL</u>: If Empty: Do not reuse this container. Place in trash or offer for recycling if available. If Partly Filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused

<u>reactly three</u>. Call your local solid waste agency of 1-000-CELEAROT for disposal histocerologies. Rever place unused product down any indoor or outdoor drain. <u>IF PRODUCT IS SPILLED</u>: Sweep up and use as per label instruction or dispose of as indicated above.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container, offer for recycling if available.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. This product is a turf fertilizer and a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in established turf grasses, golf courses (excluding putting greens), for sod farms, and lawns, and around landscape ornamentals, established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when this product is activated by at least ½ inch of rainfall, irrigation or shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application. Do not graze or feed livestock forage cut from areas treated with this product. Do not apply to golf course putting greens. Failure to follow the directions for use and precautions on this label may result in poor weed control, crop injury, or illegal residues.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (sod farms, commercial nurseries). Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls; chemical resistant gloves such as butyl rubber > 14 mils or natural rubber > 14 mils or natural rubber > 14 mils.

NON-AGRUCULTURAL USES REQUIREMENTS

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The "WPS" applies when this product is used to produce agricultural plants on farms, forests or nurseries. Applicators and handlers should wear long-sleeved shirt, long pants, chemical resistant gloves, shoes with socks and protective eyewear. Thoroughly wash face and hands with soap and water after exposure and before eating or smoking. Launder exposed clothing before reuse. Do not allow reentry to treated areas until dusts have settled and the turfgrass or soil is dry.

APPLICATION DIRECTIONS

Apply uniformly with suitable, calibrated application equipment. Do not apply otherwise.

ESTABLISHED TURF

This product is a selective pre-emergence herbicide that, when properly applied, will control certain grass and broadleaf weeds listed below in established turf grass and lawns. The maximum amount of material that may be applied per year is given for each turf grass species in the <u>Maximum Annual Rates</u> section of this label. Most effective weed control in turf grasses will be obtained when this product is activated by at least ½ inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map for approximate crabgrass seed germination dates.



Approximate Date After May 30 After May 10 After April 20 After March 20 January 1 to March 20

2

.

USE PRECAUTIONS

The following precautions apply to the use of this product in turf grasses and lawns: (1) Application of this product may thin emerged annual bluegrass and newly overseeded grasses. (2) Do not apply to overseeded turf within 60 days after seeding or until after the second mowing whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil. (3) Do not cut (harvest) treated sod before 120 days after application. Do not apply to newly set sod until the following year. (4) Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. (5) Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control. (6) Do not apply this product to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.

RATES OF APPLICATION

This product may be applied as a single application or in sequential application to control weeds germinating throughout the year. All applications must be made prior to germination of the weeds listed below. This product will not control established weeds. Maximum use rate section should be based on turf species. The length of time of residual preemergence weed control provided by this product is related to the rate applied.

MAXIMUM ANNUAL RATES

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

	i yearry rate as given in the ton	
MAXIMUM APPLICATION RATE/	CALENDAR YEAR OF PRO	DUCT BY TURF GRASS SPECIES (1)
Turf Species	Lbs product per acre	Lbs. a.i. per acre
Creeping Bentgrass	144 lbs.	0.65 lbs.
Creeping Red Fescue	167 lbs.	0.75 lbs.
Buffalograss	222 lbs.	1.00 lbs.
Kentucky Bluegrass	"	n
Perennial Ryegrass	п	н
Bermudagrass ⁽²⁾	333 lbs.	1.50 lbs.
Bahiagrass	"	n
Centipedegrass	"	n
Seashore Paspalum	н	п
St. Augustinegrass	"	11
Tall Fescue (including turf type)	"	"
Zovsia	"	"

⁽¹⁾ These are the maximum rates per calendar year by species limitations. ⁽²⁾ May be used on newly-sprigged or plugged Bermudagrass at rates not to exceed 116 lbs. per acre. Newly-sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations.

Do not apply more than 333 lbs. of this product to the same area per calendar year. Use higher rates of product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum application rates specified in the <u>Maximum Annual Rates</u> section.

WEEDS CONTROLLED

	TEEDO CONTROEDOD	
When used in accordance with this la	bel this product will provide control o	f the following weeds:
Barnyardgrass	Henbit	Purslane, Common
Bluegrass, Annual (Poa Annua)	Itchgrass	Pusley, Florida
Carpetweed	Johnsongrass (from seed)	Rescuegrass ⁽³⁾
Chickweed, Common	Junglerice	Shepherd's Purse ⁽²⁾
Chickweed, Mouseear (from seed)	Knotweed	Signalgrass, Broadleaf
Crabgrass (Large, smooth)	Kochia	Speedwell, Persian
Crowfootgrass	Lambsquarter, Common	Sprangletop
Cupgrass, Woolly	Lovegrass	Spurge, Prostrate
Foxtails, Annual	Panicum (Texas, Fall, Browntop)	Witchgrass
Goosegrass ⁽¹⁾	Pigweed	Woodsorrel, Yellow (from seed)
0)- · · · · · · · · · · · · · · · · · · ·	· · · · · · · · · · · · · · · · · · ·	

⁽¹⁾In many areas a single application of 144 to 333 lbs. per acre will control Goosegrass. However, under heavy Goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by ⁽¹⁾, ⁽¹⁾ making an initial application of 144 to 222 lbs per acre followed after 60 to 90 days by a second application at doses that would not exceed those given in the <u>Maximum Annual Rates</u> section. Do not exceed the maximum rate for turf grass species listed in the <u>Maximum Annual Rates</u> section. ⁽²⁾Application for Shepherd's Purse', ⁽¹⁾ should be made in late summer, fall or winter prior to germination. ⁽³⁾ Suppression only of Rescuegrass.

Sequential application may be made so long as the total amount of product applied does not exceed the maximum endured application rates recommended for each turf species. All applications must be made prior to germination of the weight seeds.

WHEN TO APPLY AFTER OVERSEEDING TURF

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding and or reseeding results.

Product Rate Lbs./Acre	A.I./Acre	North	Transition	South
111	0.50	4	4	4
144	0.66	5	4	4
167	0.74	6	5	5
178	0.80	-	6	6
222	2.00	-	7	7
253	1.14	-	-	9
289	1.30	-	-	10
333	1.50	-	-	12

APPLICATION EQUIPMENT AND SPREADER SETTINGS

The size and bulk density of granules in this product may vary; therefore, calibrate your granular applicator prior to application of this product. Follow directions in the owner's manual for calibrating your spreader. The settings indicated in the following table should only be used as a guide in calibrating your spreader.

	SPREADER SETTINGS				
To apply the recommended amounts of a.i. per acre which are as follows:	0.65 lbs.	0.75 lbs.	1.0 lbs.	1.50 lbs.	
50 LBS. TREATS =	14,851 SQ. FT.	13,248 SQ.FT.	9,811 SQ.FT.	6,540 SQ. FT	
Scott's Evergreen 18'	# XX	# XX	# XX	# XX	
Scotts Speedy Broadcast	# XX	# XX	# XX	# XX	
Red Devil Broadcast	# XX	# XX	# XX	# XX	
Republic EZ Drop 18"	# XX	# XX	# XX	# XX	
Republic EZ Broadcast	# XX	# XX	# XX	# XX	
Republic EZ Tractor Broadcast	# XX	# XX	# XX	# XX	

LANDSCAPE ORNAMENTALS INCLUDING ESTABLISHED WILD FLOWER PLANTINGS

This product may be applied for residual pre-emergence weed control in ornamentals listed on this label.

USE RATES

Apply at 144 to 333 lbs. per acre (3.33 lbs. to 7.64 lbs. per 1,000 sq. ft.) of landscape plants, in fall and /or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 333 lbs. per acre.

APPLICATION TIMING AND INFORMATION

Apply uniformly to the surface of the soil in ornamental planting beds with a granular applicator. Calibrate your applicator to assure appropriate dosage as described above under <u>Use Rates</u>, prior to application. Calibrate your applicator as described in the owner's manual. The bulk density of this product may vary; therefore, to assure that the product is applied at the rates described, you must use a calibrated applicator.

This product is a pre-emergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least ½ inches of rainfall or irrigation, or with shallow (1 to 2 inches) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves. Prior to application of this product, control existing vegetation by hand weeding, cultivation, or the use of an appropriate post-emergence herbicide.

ESTABLISHED LANSCAPE ORNAMENTALS

This product may be applied to established ornamentals as a broadcast, over the top, or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate this product in the soil.

NEWLY TRANSPLANTED ORNAMENTALS

This product may be applied to newly transplanted ornamentals as a broadcast, over the top, or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union. Irrigate after application as above directed.

,

TOLERANT ORNAMENTAL SPECIES

This product will not harm most trees, shrubs, vines and flowers. The species listed here are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to this product.

FOR USE AROUND LANDSCAPE ORNAMENTALS, INCLUDING ESTABLISHED PERENNIALS AND WILDFLOWERS, AS LISTED.

Scientific Name Abies spp.

Abella grandiflora Acer palmatum A. platanoides Achillea, sp. Actinidia chinensis^a Agapanthus africanus Agapanthus orientalis Akebia quintata

Allium cernuum Anemone hybrida Aquilegia sp. Arctostaphylos densiflora Arctotheca calendula Artemesia sp.

Aster sp. Aster X Frikartii Athrium Filiz-femina Aucuba japonica Begonia sp.

Berberis gladwynensis B. julianae B. mentorensis B. thunbergil

B. verruculosa Bergenia cordifolia Boltonia asteroids Bougainvillea spp. Buddleia davidii Buxus microphylla Callistemon viminalis Calliana vulgaris Campia X. Tagliabuana

Capobrotus edulis Cassia artemisoides Ceanothus rigidus Ceratostigma plumbagonoides Chamaccyparis pisifera Chrysanthemum nipponicum Cleyera japonica Citrus spp.*

Coreopsis sp.

Cornus florida C. stolonifera Cortaderia selloana Contoneaster apiculatus

C. buxifolius

C. dammeri C. microphyllus Cretaegus spp. Crocosmia sp. Cupressus sempervirens Delosperma alba Fir Species** (Balsam, Fraser, Noble, Etc.) Abelia: Sherwood Japanese Maple Norway Maple Yarrow: King Edward Kiwi* Lily-of-the-Nile (African Lily)

Common Name

Five-Leaf or Chocolate Vine

Lady's Leek, Nodding Onion Japanese Anemone Aquilegia: Red and Gold Vine Hill Manzanita Cape Weed Wormwood: Silver Mound, Castle

Aster: Bonny Blue, Purple Dome

Lady Fern: Fern Lady Japanese Aucuba Fibrous Begonia: Hardy Grandis Barberry Wintergreen Barberry Mentor Barberry Japanese Barberry

Warty Barberry

Snowbank Bougainvillea Butterfty-Bush (Dwarf Blue); Royal Red Japanese Boxwood Weeping Bottlebrush Scotch Heather Tussock Bellflower (White Clips) Trumpet Creeper, Trumpet Flower, Madame Galen Hottentot Fig (Ice Plant) Feathery Cassin Wild Liac

Faise Cypress

Cleyera Citrus species*

Coreopsis (Calliopsis): Early Sunrise, Moonbeam Flowering Dogwood American Dogwood Pampas Grass Cranberry Cotoneaster

Cotoneaster

Bearberry Cotoneaster Rockspray Cotoneaster Hawthorne Lucifer Italian Cypress White Trailing Ice Plant

Scientific Name Delosperma sp.

Delphinium sp. Dianthus deltoids D. gratianopolitanus Dodonea viscosa Echinacea purpurea Elacagnus pungens Euonymus fortunei E. japonica

E. kiautschovica Fatsia japonica Forsythia intermedia F. suspense F. viridissima Gaillardia sp.

Gardenia jasminoides Gaura sp. Gentiana daburica Geranium cinereum Gladiolus spp.**

Gypsophilia repens Hedera helix Hellanthemum sp. Remerocallis sp.

Hibiscus** Hibiscus sp. Hibiscus rosa-sinensis Hosta plantaginea H. Sieboldiana Houttuynia cordata var. variegate Hydrangea macrophylla Llex corauta L creanta L opaca

I. pernyi I. vornitoria Inula ensifolia Iris ensata

Iris siberica Iris spp.** Jasminium udiflorum Juniperus chinensis

J. conferta

J. davurca J. horizontalis Juglans sp.* Justicia brandegeana Lagerstromia Indica Lagerstromia indica X Fauriei Lantana montevidensis Lavender sp. Leontopodium alpinum Ligustrum sinense L. amurense L. japonicum Common Name Cooperi Pink

Larkspor: Blue Elf Dianthus: Maiden Pinks "Zing" Cheddar, Pink Hop Bush Coneflower, Purple: Magnus Silverberry Wintercreeper Japanese Spindle Tree (Evergreen Euooymus) Sprending Euonymus Japanese Aralia Border Forsythia Greenstem Forsythia Gaillardia, Blanket Flower: "Goblin"

Gentian Cranesbill Gladiolus species**

Baby's Breath English Ivy Suarose Daylily: Aztec Gold, Stella De, Oro, Tender Love Rose of Sbaron** Mallow: Disco Belle White Chinese Hibbiscus Hosta, Plantain Lily (Fragrant) Hosta, "Searsucker"

Bigleaf Hydrangea Chinese Holly Japanese Holly; Helleri American Holly

Holly Yaupon Holly; Schillings

Sword-Leaved Iris; Jodlesong

Siberian Iris; Cabernet Iris species** Winter Jasmine Chinese Juniper; Nick's Compact, Parsonnil Shore Juniper; Blue Pacific

Parsoni Creeping Juniper Walnut* Shrimp Plant

Crape Myrtle; Crape Myrtle; Tuscarora Weeping Lantana Lavender; Munstead Edelweiss Chinese Privet; Variegata Amur Privet Japanese Privet

9/10

Scientific Name L. jucidum

Lillum sp. Liriope muscari Liriope Muscari var. variegate Liriope spicata Lobelia Cardinalis Lonicera japonica L. tatarica Loropetalum chinense Lythrum sp. Magnolia spp.** Maleophora luteola Malus sp.* Miscanthus sinensis Nandian domestica Narcissus spp.** Nerium oleander Olea europea* Ophiopogon japonicus Osmanthus beterphyllus Osteospermum fruticosum Oxydendrum arboretum Paeonia suffruticosa Pennisetum setaceumi Perovskia atriulicifolia Persea americana Photinia fraseri Physotegia Virginiana Picea spp.** Pier is japonica Pinus brutia P. canariensis P. elliottii P. halepensis

Pinus brautia P. canariensis P. elliotti P. halepensis P. palustrus P. palustrus P. radiate P. tradiate P. strobes P. strobes P. stylvestris P. trobes P. virginiana Pistachio sp.* Pittosporum rhombifolium P. tobira Podocarpus macrophyllus Prunus fauroserasus Prunus fauroserasus

Pseudotsuga menziesii** Pyracantha coccinea P. fortuneana P. hotdzumii Pyrus sp. Quercus Shumardii Quercus Spp. Raphiolepis indica Raphiolepis umbellate Rhododendron (including Azalea)

Rosa banksiae Rosmarinus officinalis* Rudbeckia sp. Santolina, virens Saxifraga sp.

Scabiosa sp. Sedum cauticola Sedum tayphyliom Sedum spuriam Spiraea Bumalda Syzygium paniculatum Taxus cuspidate T. media Teucrium sp. Thalictrum dipterocarpom Thuja occidentalis Trachelospermum asiatum Tsuga canadensis Tulipa spp. Common Name Glossy Privet (Waxleaf)

Liriope, Variegated

Lily; Jazz Big Blue Lily; Turf Silver Mound; Evergreen Giant

Liriope, Creeping Cardinal Flower, Indian Pink Jananese Honeysuckel Tatarian Honeysuckel Burgundy Looseleaf; Modern Pink Magnolia species** Ice Plant Crabapple* Yaku Jima, Silberfeder Heavenly Bamboo Narcissus species** Oleander Olive* Mondo Grass Osmanthus (False Holly): Gulf Tide Trailing African Daisy Sourwood Tree Peony Fountain Grass (Dwarf) Avocado* Frasier's Photinia (Redtip) Dragonhead, False; Vivid Spruce species**(Colorado Blue, Norway,etc.) Lily-of-the-Valley Bush Calabrian Pine Canary Island Pine Splash Pine Aleppo Pine Australian Pine Longleaf Pine Monterev Pine Japanese Black Pine Eastern White Pine Scotch Pine LobloHy Pine Virginia Pine Pistachio* Queensland Pittosporum Japanese Pittosporum Japanese Yew English Laurel Almond, Apricot, Nectarine, Peach, Plum and Prune* Douglas Fir** Firethorn (Scarlet) Firethorn Firethorn Bradford Pear sp. Oak, Shumard's Red Oak, species Indian Hawthorn Yedda Hawthorn 'Coral Bells' Delaware Valley White' Flame Creeper 'Formosa' Girard Crimson 'George L. Tabor' 'Hino-crimson' 'PJM' 'Roseum Elegans' 'Wakeiebisu' 'White Gumbo' Lady's Bank's Rose Rosemary* Black-Eyed Susan; Goldstrum Santolina Saxifrage; Purple Dome Pincushion Flower Stonecrop Stonecrop; Lidakense Stonecrop Stonecrop; Dragon's Blood Spirea; Anthony Waterer Australian Brushcherry Japanese Yew Yew Germander Meadow Rue American Arborvitae Star Jasmine Canada Hemlock Tulip species

Scientific Name Veronica so.

Vibrurnum japonicum V. odoratissimum V. plicatum

V. rigidum V. suspensum V. tinus V. trilobium V. trilobium V. Wrightii Vinca major Vinca minor Vitis sp.* Weigela florida Yucca aloifolia Y. filamentosa Common Name Veronica, Speedwell; Sunny Border Japanese Viburnum Sweet Viburnum Japanese Snowball

Canary Island Viburnum Arrow wood Viburnum Laurustinus Cranberry Bush Leatherteaf Viburnum Greater Periwinkle, Vinca Common Periwinkle, Vinca Grape* Old Fashioned Weigela Spanish Bayonet, Yucca Adam's Needle, Yucca ٩,

а н

* Do not use on food producing trees, vines, or plants

** Not for use on container grown plants

CONDITIONS OF SALE AND WARRANTY

Read the entire <u>Directions for Use</u>, <u>Conditions</u>, <u>Disclaimer of Warranties</u> and <u>Limitation of Liability</u> before using this product. If the terms are not acceptable, return the unopened container at once. By using this product, user and buyer accept the following <u>Conditions</u>, <u>Disclaimer of Warranties</u>, and <u>Limitations of Liability</u>.

<u>CONDITIONS</u>: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Harrell's, Inc. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: There are no warranties express or implied, of merchantability or of fitness for a particular purpose or otherwise, which extend beyond the statements made on this label. No agent of Harrell's, LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Harrell's, LLC disclaims any liability whatsoever for special, incidental, or consequential damages resulting from the use or handling of this product.

LIMITATION OF WARRANTIES: The exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price, or at Harrell's,LLC option, the replacement of the product.

NET WEIGHT: 50 POUNDS

Harrell's, LLC P. O. Box 807 Lakeland, FL 33802

Batch No.