JUN 25 2003

Mr. Robert F. Bischoff Agent for HARRELL'S INC. Total Turf Consulting LLC P.O. Box 7928 Hilton Head, SC 29938

Dear Mr. Bischoff:

This is in response to your request dated May 30, 2003 to change the primary brand name for **EPA Registration 52287-9** from <u>Wilbro Fertilizer With .75% Ronstar</u> to:

TURF FERTILIZER WITH RONSTAR® 0.75

Your request for the name change is accepted and the change will be made to the Agency's records. All future correspondence on this product should reference the new name.

Sincerely yours,


Sherada D. Hobgood Notifications Review Coo

Notifications Review Coordinator Registration Division (7505C)

	يمو	æ	erse before comple	tina form.			Form Apr	proved	. OM	IB No.	2070-00	60. Approvel expires 2-28-95		
≎EPA	tion Agency			~		- 1end:	ation ment	OPP Identifier Number						
			1	Applicati	ion for F	Pestici	de - Seci	tion	<u> </u>					
1. Company/Product Num 52287-9	2. EPA Product Manager JOANNE 1. MILLER						3. Proposed Classification None Restricted							
4. Company/Product (Nan WLBRO FERTILIZE	ne) R	M	TH .75% RONS	TAR	PM# 23									
5. Name and Address of A	App	lice	ant <i>(Include ZIP Cod</i>	de)								th FIFRA Section 3(c)(3)		
HARRELL'S, INC. P. O. BOX 807 LAKELAND, FL 33	(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. NOTIFICATION													
Check if t	his	is (a new address		Product Name N/A JUN 2 5 2003									
		_			Sect	tion - i	(
Amendment - Expl	Final printed labels in repsonse to Agency letter dated "Me Too" Application. Other - Explain below.													
Explanation: Use addit			/ if											
NOTIFICATION OF PRIM NOTIFICATION IS CONS REGULATIONS AT 40 CF STATEMENT OF FORMU STATEMENT TO EPA. I AND 2001-6 AND THE RE	ISTI FR 1 JLA FUF	EN 156. OF RTH	T WTH THE PROVI 5.10 AND 40 CFR 152 T THIS PRODUCT. I HER UNDERSTAND	'ISIONS/GUIE 12.46, AND NO I UNDERSTA) THAT IF TH	DANCE OF F O OTHER C AND THAT IT IIS NOTIFICA D 40 CFR 15	PR NOTIC HANGES TIS A VIO ATION IS 52.46. THI	ES 98-10 AN HAVE BEEN LATION OF NOT CONSI IS PRODUCT	ID 200 I MADI 18 U.S STENT)1-6 A1 E TO 1 S.C. SE T WITH	ND THE THE LA EC.100 H THE	E REQUIF ABELING 11 TO WIL GUIDAN	REMENTS OF EPA OR THE CONFIDENTIAL LIFULLY MAKE ANY FALSE CE OF PR NOTICES 98-10		
			<u> </u>		Sect	ion - II								
1. Material This Product V Child-Resistant Packaging	—-т				15/2005	Calabla De			T. T		Containe			
Child-Hesistant Packaging		ان.	nit Packaging Yes		Water Soluble Packaging				2. 1) r				
No			No	No							Plastic Glass	;		
Certification must			"Yes" nit Packaging wgt.	No. per container	If "Yes" Package						Paper	(Specify)		
					etail Contain			5 Lor	cation	of Lat	oel Directi	inna		
Label									[
6. Manner in Which Label	xed to Product	Lithograph Othe Stanciled												
		_			Secti	on - I\	/					. s		
1. Contact Point Comple	te i	ten	ns directly below fo	r identificati	on of individ	dual to be	contacted, i	if nece	essery	r, to pr	oc ess thi	is application.)		
Name ROBERT F. BISCHOFF					1 .					•	one No. (Include Area Code) 173-0222			
i certify that the sta i acknowledge that both under applicable	any	kn	nowlinglly false or m		d all attachn							6. Cate Application Received (Stamped)		
2. Signature Retex F. Bischoff					3. Title AGENT, TOTALTURF CONSULTING LLC									
4. Typed Name ROBERT F. BISCHOFF	5. Date MAY 30, 2003													

Section II (Explanation Continued)

SUBJECT TO ENFORCEMENT ACTION AND PENALTIES UNDER SECTIONS 12 AND 14 OF FIFRA. NON-NOTIFICATION LABEL CHANGES CONSISTENT WITH PR NOTICE 98-10 (IV-D) INVOLVE NEW REGISTRANT NAME, ADDRESS, REGISTRATION NUMBER, AND ESTABLISHMENT NUMBER AS A RESULT OF THE TRANSFER OF REGISTRATION FROM WILBRO TO HARRELL'S.





MOTIFICATION
JUN 2 5 2003

TURF FERTILIZER WITH RONSTAR® 0.75

50 Lb bag treats up to 8,150 Sq. Ft.

See supplemental label for the use of this product on ornamental shrubs, vines, and trees

FOR SALE TO, USE AND STORAGE BY COMMERCIAL NURSERY, TURF AND LANDSCAPE PERSONNEL ONLY. FOR COMMERCIAL USE ON PARKS, GOLF COURSES, ATHLETIC FIELDS, AND OTHER RECREATIONAL AND ORNAMENTAL TURFGRASS AREAS. A PREEMERGENT WEED CONTROL PLUS FERTILIZER FOR PERENNIAL BLUEGRASS, PERENNIAL RYEGRASS, ST. AUGUSTINEGRASS, BERMUDAGRASS, TALL FESCUE & ZOYSIAGRASS. Controls: Crabgrass, Annual Bluegrass, Goosegrass, Florida Pusley, Oxalis, Stinging Nettle, Carpetgrass, Pigweed and Other Weeds Listed

ACTIVE INGREDIENT:

Oxadiazon: 2-tert-butyl-4-(2,4-dichloro-5-isopropoxyphenyl)-z-1,3,4 oxadiazolin-5-one. 0.75% INERT INGREDIENTS: 99.25% TOTAL 100.00%

Contains RONSTAR® herbicide, a registered trademark of Bayer Environmental Science

EPA Reg. No. 52287-9

EPA Est. No. 52287-FL-1

GUARANTEED FERTILIZER ANALYSIS

Total nitrogen (N).....variable%

Nitrate nitrogen.....X%

Ammoniacal nitrogen.....X%

Water soluble nitrogen and/or urea nitrogen.....X%

Water insoluble nitrogen...X%

Available phosphoric acid (P2O5).variable%

Soluble potash (K2O).....variable%

Derived from ammonium sulfate, ureaform, muriate of potash; manganese as Mn .5%; iron as Fe 1.0%, chlorine <15%

KEEP OUT OF REACH OF CHILDREN

WARNING—AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detaile. (If you do not understand the label, find someone to explain it to you in detail.)

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 tall glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes eye and skin irritation. Avoid contact with skin, eyes or clothing. Do not allow domestic animars to graze treated areas. Avoid contamination of feed and foodstuffs.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear coveralls over short-sleeved shirt and short pants; waterproof gloves; chemical resistant footwear plus socks; protective eyewear, chemical-resistant headgear for overhead exposure; chemical resistant apron when cleaning equipment. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.



USER SAFETY RECOMMENDATIONS

Users should: Wash hands before eating, drinking, chewing gym, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean highwater mark. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters. Apply only as specified on this label.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS
Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls; waterproof gloves, chemical resistant footwear plus socks, protective eyewear, and chemical resistant headgear for overhead exposure.

NON-AGRUCULTURAL USES REQUIREMENTS

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants only on farms, forest, nurseries, sod farms or geeenhouses. Thoroughly wash face and hands with soap and water after exposure and before eating or smoking. Wear protective clothing including long-sleeved shirt and impervious gloves when handling. Launder exposed clothing before reuse.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store product in its original bag, in a cool, dry, tocked place out of reach of children. In case of spill, sweep-up, and use according to directions or dispose of as follows:

PRODUCT DISPOSAL: If Empty: Do not reuse this container. Place in trash or offer for recycling if available. If Partly Filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

FOR LAWNS AND ORNAMENTAL TURF

NOT FOR USE ON HOME LAWNS. FOR SALE TO, USE AND STORAGE BY COMMERCIAL NURSERY, TURF AND LANDSCAPE PERSONNEL ONLY

When applied according to directions, this product can be used safely on the following established turfgrass species: BERMUDAGRASS, PERENNIAL BLUEGRASS, PERENNIAL RYEGRASS VARIETIES, ST. AUGUSTINEGRASS VARIETIES, TALL FESCUE, AND ZOYSIAGRASS lawns and ornamental turf. It may be used for apartment and condominium complexes, parks and golf courses, athletic fields, cemeteries and all other ornamental and recreational turfgrass areas.

CONTROLS GRASSY WEEDS AND BROADLEAF WEEDS: Annual Bluegrass (Poa annua), Barnyardgrass, Crábgrass, Fall Panicum, Florida Pusley, Goosegrass, Green Foxtail, Oxalis, Stinging Nettle, Carpetgrass, Pigweed.

HOW TO APPLY

For best weed control rake the area free of leaves and thatch and mow if necessary so that granules will reach the goll surface. Measure the area to be treated and uniformly apply the required amount using a fertilizer spreader (see Spreader Setrings chart). Do not apply to turf when it is wet. Do not overlap spreader applications on St. Augustinegrass since discoloration "streaking" may temporarily occur from "double rate". After applications, if rain is not expected within 24 hours, sprinkle the treated turf with 1/2 to 1/2. inch of water to get maximum weed control.



HOW MUCH TO USE

CRABGRASS: Apply 50 lbs./8,150 sq. ft. when turf has been in a yearly maintenance program (equal to 2 lbs. Al herbicide per acre). Apply 50 lbs. to treat 5,500 sq. ft. of lawn if it has been heavily infested with crabgrass; equal to 3 lbs. Al herbicide per acre. Apply in early spring 1 to 2 weeks prior to weed seed germination. This product is a pre-emergence herbicide. It will not control emerged and growing weeds; only germinating weed seeds. Late applications will control only those weed seeds that germinate after the application. See "When to Apply" chart.

CONSEGRASS & PROSTRATE SPURGE: These weeds do not usually germinate until late spring or early summer. Apply 50 lbs. to treat 4,100 sq. ft (4 lbs. Al per acre) and apply 1 to 2 weeks prior to seed germination. See Chart for When to Apply. Observe maximum rate for St. Augustinegrass.

ANNUAL BLUEGRASS (POA ANNUA): Apply 50 lbs. per 4,100 to 4,600 sq. ft. of lawn; equal to 3.5 to 4 lbs. A.l.herbicide per acre. Apply in late summer to fall or in early spring 1 to 2 weeks before expected weed seed germination. Will not control annual bluegrass present at time of application; therefore only germinating seeds. Refer to chart for proper timing in your locale. Annual barnyardgrass and prostrate spurge may also be controlled by this same rate. Observe maximum rate for St. Augustinegrass. PIGWEED, FLORIDA PUSLEY, OXALIS, STINGING NETTLE, AND CARPETGRASS: Where these weeds are a problem, apply 50 lbs. to treat 4,100 to 4,600 sq. ft. of lawn (equal to 3.5 to 4 lbs. A.l.herbicide per acre), in early spring or 1 to 2 weeks before expected weed seed germination. Will not control emerged and growing weeds; only effective against germinating weed seeds. Observe maximum rate for St. Augustinegrass (50 lbs. per 5,500 sq. ft.).

SPREADER SETTINGS FOR SOUARE FOOT COVERAGE

SPREADER	AMT. OF PRODUCT	8,150 SQ. FT. =2#A.I./ACRE	5,500 SQ.FT. =3#A.I./ACRE	4,100 SQ.FT. =4#A.L/ACRE
Cyclone B-1	50 lbs.	4 1/2	5 1/2	5 3/4
Scotts #35	50 lbs.	5	6	7

NOTE: Spreader rate varies with walking speed. Check spreader over known area.

WHEN TO APPLY APPROPRIATE TIMES TO APPLY THIS PRE-EMERGENT HERBICIDE											
JAN	FEB	MAR 1	APR	MAY	JUNE 2	JULY	AUG	SEP	OCT 3	NOV	DEC
	4			2a.——				3a			
1.	For CRABGRASS and other grasses and broadleaf weeds germinating in the Spring in Northern turf.										
2.	For GOOSEGRASS and other late germinating annual grasses and broadleaf weeds such as PROSTRATE SPURGE.										
2a.	GOOSEGRASS in Southern turf (PRECAUTIONS IN USING).										
3.	For POA ANNUA in Northern turf.										
3a.	For POA ANNUA in Southern turf.										
4.	For fall,	winter, spr	ing control	of ANNUA	AL WEEDS	in Southern	n turf.				

PRECAUTIONS IN USING

Do not use on desirable red fescue, bentgrass varieties, dichondra, or centipedegrass since damage will occur. Do not use on putting greens or tees due to varying cultural practices used. Use only the rate of 50 lbs. to tread 5,500 sq. ft on St. Augustinegrass since temporary discoloration will occur from heavier rates. Avoid contact with flowers and shrubs except as recommended on the ornamental use supplemental labeling available for this product. For newly seeded lawns do no apply this product until after the second or third cutting. For best lawn care, place new grass seed in the fall of the year. Spring pre-emergence weed control can then be applied.

RESEEDING TREATED AREAS

Delay new seeding until 4 months after treatment.

NOTICE: Buyer and user assume all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

NET WEIGHT: XX POUNDS

Harrell's, Inc. P. O. Box 807 Lakeland, FL 33802