

52287-12

6-25-2003

4/8

JUN 25 2003

Mr. Robert F. Bischoff
Agent for HARRELL'S INC.
Total Turf Consulting LLC
P.O. Box 7928
Hilton Head, SC 29938

Dear Mr. Bischoff:

This is in response to your request dated May 30, 2003 to change the primary brand name for **EPA Registration 52287-12** from Wilbro Fertilizer With Starteem #3 to:

FERTILIZER WITH STARTEEM® # 3

Your request for the name change is **accepted** and the change will be made to the Agency's records. All future correspondence on this product should reference the new name.

Sincerely yours,



Sherada D. Hobgood
Notifications Review Coordinator
Registration Division (7505C)



United States
Environmental Protection Agency
 Washington, DC 20460

| | |
|-------------------------------------|--------------|
| <input type="checkbox"/> | Registration |
| <input type="checkbox"/> | Amendment |
| <input checked="" type="checkbox"/> | Other |

OPP Identifier Number

Application for Pesticide - Section I

| | | |
|--|---|--|
| 1. Company/Product Number 52287-12 | 2. EPA Product Manager JOANNE I. MILLER | 3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted |
| 4. Company/Product (Name) WLBRO FERTILIZER WITH STARTEEM #3 | PM# 23 | |
| 5. Name and Address of Applicant (Include ZIP Code) HARRELL'S, INC. P. O. BOX 807 LAKELAND, FL 33802 <input type="checkbox"/> Check if this is a new address | 6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. <u>N/A</u> NOTIFICATION Product Name <u>N/A</u> JUN 25 2003 | |

Section - II

| | |
|--|--|
| <input type="checkbox"/> Amendment - Explain below. | <input type="checkbox"/> Final printed labels in response to Agency letter dated _____ |
| <input type="checkbox"/> Resubmission in response to Agency letter dated _____ | <input type="checkbox"/> "Me Too" Application. |
| <input checked="" type="checkbox"/> Notification - Explain below. | <input type="checkbox"/> Other - Explain below. |

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

NOTIFICATION OF PRIMARY BRAND NAME CHANGE PER PR NOTICE 98-10 AND LABEL CHANGE RELATIVE TO PR NOTICE 2001-6. THIS NOTIFICATION IS CONSISTENT WITH THE PROVISIONS/GUIDANCE OF PR NOTICES 98-10 AND 2001-6 AND THE REQUIREMENTS OF EPA REGULATIONS AT 40 CFR 156.10 AND 40 CFR 152.46, AND NO OTHER CHANGES HAVE BEEN MADE TO THE LABELING OR THE CONFIDENTIAL STATEMENT OF FORMULA OF THIS PRODUCT. I UNDERSTAND THAT IT IS A VIOLATION OF 18 U.S.C. SEC. 1001 TO WILLFULLY MAKE ANY FALSE STATEMENT TO EPA. I FURTHER UNDERSTAND THAT IF THIS NOTIFICATION IS NOT CONSISTENT WITH THE GUIDANCE OF PR NOTICES 98-10 AND 2001-6 AND THE REQUIREMENTS OF 40 CFR 156.10 AND 40 CFR 152.46. THIS PRODUCT MAY BE IN VIOLATION OF FIFRA AND I MAY BE

Section - III


| | | | | | |
|---|---|--|-------------------|---|----------------------------------|
| 1. Material This Product Will Be Packaged In: | | | | 2. Type of Container | |
| Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No | Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No | Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No | | <input type="checkbox"/> Metal | <input type="checkbox"/> Plastic |
| Certification must be submitted | | If "Yes" Unit Packaging wgt. | No. per container | If "Yes" Package wgt. | No. per container |
| | | | | <input type="checkbox"/> Glass | <input type="checkbox"/> Paper |
| | | | | <input type="checkbox"/> Other (Specify) _____ | |
| 3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container | | 4. Size(s) Retail Container | | 5. Location of Label Directions <input type="checkbox"/> | |
| 6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled | | | | <input type="checkbox"/> Other _____ | |

Section - IV

| | | |
|---|---|---|
| 1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.) | | |
| Name ROBERT F. BISCHOFF | Title AGENT, TOTALTURF CONSULTING LLC | Telephone No. (Include Area Code) (317) 873-0222 |
| Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment, or both under applicable law. | | 6. Date Application Received (Stamped) |
| 2. Signature <i>Robert F. Bischoff</i> | 3. Title AGENT, TOTALTURF CONSULTING LLC | |
| 4. Typed Name ROBERT F. BISCHOFF | 5. Date MAY 30, 2003 | |

Section II (Explanation Continued)

SUBJECT TO ENFORCEMENT ACTION AND PENALTIES UNDER SECTIONS 12 AND 14 OF FIFRA. NON-NOTIFICATION LABEL CHANGES CONSISTENT WITH PR NOTICE 98-10 (IV-D) INVOLVE NEW REGISTRANT NAME, ADDRESS, REGISTRATION NUMBER, AND ESTABLISHMENT NUMBER AS A RESULT OF THE TRANSFER OF REGISTRATION FROM WILBRO TO HARRELL'S.



FERTILIZER WITH STARTEEM® # 3

A NEW! PRE-EMERGENCE HERBICIDE COMBINATION FOR GOLF COURSES, PARKS, ATHLETIC FIELDS, CEMETERIES, UNIVERSITY, FEDERAL, MUNICIPAL, CORPORATE LAWNS and SOD FARMS

NOTIFICATION
JUN 25 2003

50 lb. Bag Treats up to 10,870 Sq. Ft.

FOR SALE TO, USE AND STORAGE BY: GOLF COURSES, SOD FARMS, COMMERCIAL NURSERIES, TURF MANAGEMENT FIRMS, MUNICIPAL, FEDERAL, AND CORPORATE PARKS, LAWNSCAPES, CEMETERIES, AND GROUNDS MAINTENANCE PERSONNEL.

NOT FOR USE ON HOME LAWNS.

STARTEEM® is a pre-emergent weed control product plus a quality fertilizer for use on varieties of Perennial Bluegrass, Bentgrass, Bermudagrass, Buffalograss, Perennial Ryegrass, St. Augustinegrass, Seashore Paspalum, Tall Fescue and Zoysiagrass.

Also for use around Ornamental Shrubs, Vines, Trees, & Ground Covers

ACTIVE INGREDIENTS:

| | |
|--------------------------|---------------|
| Oxadiazon | 1.00% |
| Benefin | 0.25% |
| Trifluralin | 0.25% |
| OTHER INGREDIENTS: | 98.50% |
| | Total 100.00% |

EPA Reg. No. 52287-12

EPA Est. No. 52287-FL-1

GUARANTEED FERTILIZER ANALYSIS

| | |
|---|-------------|
| Total Nitrogen (N)..... | Variable X% |
| Ammoniacal Nitrogen..... | X% |
| Water Soluble Nitrogen and/or Urea Nitrogen..... | X% |
| Water Insoluble Nitrogen..... | X% |
| Available Phosphoric Acid (P ₂ O ₅)..... | Variable X% |
| Soluble Potash (K ₂ O)..... | Variable X% |
| Derived from Ammonium Sulfate, Ureaform, Muriate of Potash; Manganese as Mn X%; Iron as Fe X%; Chlorine <X% | |

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Remove person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

IF SWALLOWED: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

FERTILIZER WITH STARTEEM® # 3

A NEW! PRE-EMERGENCE HERBICIDE COMBINATION FOR GOLF COURSES, PARKS, ATHLETIC FIELDS, CEMETERIES, UNIVERSITY, FEDERAL, MUNICIPAL, CORPORATE LAWNS and SOD FARMS

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear long-sleeved shirt and long pants, socks and shoes. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Do not allow domestic animals to graze treated areas. Avoid contamination of feed and foodstuffs.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear protective eyewear; long-sleeved shirt and long pants; waterproof gloves; chemical resistant footwear plus socks. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system. Apply only as specified on this label.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, and nurseries, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. For early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear: coveralls over long-sleeved shirt and long pants, socks, shoes, and waterproof gloves.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants only on farms, forests, nurseries or greenhouses. Observe and wear (PPE) protective eyewear, long-sleeved shirt and long pants, impervious gloves and chemical resistant footwear plus socks, when handling. Launder exposed clothing before reuse. Thoroughly wash face and hands with soap and water after exposure and before eating or smoking.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.
STORAGE: Store product in its original bag, in a cool dry locked place out of reach of children. In case of spill, sweep up and use according to directions or dispose of as follows. **PRODUCT DISPOSAL:** If Empty: Do not reuse this container. Place in trash or offer for recycling if available. If Partly Filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Harrell's now offers the STARTEEM-SPLIT - a new weed control program that finally gets the maximum pre-emergence weed control results, that you should expect with 3 active ingredients.

Controls Crabgrass, Goosegrass, Poa Annuua, Annual Sedge and many weeds as listed; On Golf Courses, Commercial Lawns, and other Athletic and/or Recreational, Ornamental Turf. If you are serious about getting rid of annual grassy and certain broadleaf weeds before they go to seed again, use STARTEEM # 1 and then STARTEEM # 2 OR for hard-to-kill weeds follow # 1 with STARTEEM # 3.

FERTILIZER WITH STARTEEM® # 3

A NEW! PRE-EMERGENCE HERBICIDE COMBINATION FOR GOLF COURSES, PARKS, ATHLETIC FIELDS, CEMETERIES, UNIVERSITY, FEDERAL, MUNICIPAL, CORPORATE LAWNS and SOD FARMS

HOW MUCH TO USE AND WHEN TO APPLY THEM!

STARTEEM # 1 is to be used as the first application; STARTEEM # 2 (a different formula) is to be used as the second application of a seasonal (Split-application program) both products to be used at a rate of 200 lbs./acre (equal to 4.6 lbs. /1000 sq. ft.) STARTEEM may be applied to "established" perennial Bluegrass, Bentgrass (see restrictions), Bermudagrass, Buffalograss, Perennial Ryegrass, St. Augustinegrass, Seashore Paspalum, Tall Fescue and Zoysiagrass turf, at condominium complexes, corporate and public parks, golf courses, sod farms, athletic fields, playgrounds, and other ornamental and recreational turfgrass areas. Not for use on home lawns. The turf should be free of leaves and thatch and in condition to allow granules to reach the soil surface. Know the square footage of the area to be treated and uniformly apply using a granule spreader. Do not apply to turf when it is wet. When using, do not overlap spreader applications, especially on St. Augustinegrass since discoloration "streaking" may temporarily occur from the "doubled rate". Observe the maximum rate for St. Augustinegrass and Bentgrass of one application of each STARTEEM # 1 AND STARTEEM # 2 or # 3 per 1,000 sq. ft. per year. See below restrictions when using on Bentgrass. After applications, if rain is not expected within 24 hours, sprinkler irrigate the treated turf with up to a ¼ inch of water to get maximum weed control and quicker response from the fertilizer nutrients. STARTEEM #1, #2, and #3 must be applied prior to weed seed germination (see map of U.S.). They will not control growing weeds. See complete list of weeds that STARTEEM controls under Section for GOLF COURSES.

| Pre-emergence Weed Control with the "STARTEEM® SPLIT" | | | |
|---|--|---|--|
| Weeds Controlled | FIRST APPLICATION STARTEEM #1 | SECOND APPLICATION STARTEEM #2 or #3 | Directions |
| Crabgrass (Digitaria, spp.) Goosegrass (Bleusine indica) Field Sandbur (Canchusincertus) Annual Sedge (Cyperus compressus) Annual Bluegrass (Poa annua spp.) (See Note below) | *Apply 200 lbs. product per acre equal to 4.6# / 1,000 sq. ft. | Apply 60 days later at 200 lbs. per acre equal to 4.6# / 1000 sq. ft. | *Apply #1 40 to 60 days prior to weed seed germination in the late winter or early spring. Apply #2 or #3, 60 days later. Application made after May, may not be fully effective. Apply in late summer to early fall prior to weed seed germination. |
| Virginia Buttonweed (Diodia virginiana) Pigweed (Amaranthus spp.) Florida Pusley (Richards scabra) Oxalis (Oxalis spp.) Stinging Nettle (Urtica dolca) Carpetgrass (Axonagus officinis) | Apply 200 lbs product per acre equal to 4.6# / 1,000 sq. ft. | Apply 60 days later at 200 lbs per acre equal to 4/6# / 1,000 sq. ft. | Apply in early spring prior to weed seed germination. |
| Fall Application for Spring Weed Control in Cool Season Turf | | | |
| Crabgrass, Goosegrass and other annual weeds as listed | Apply #1 at 200 lbs per acre equal to 4.6# / 1,000 sq. ft. | Apply #3 (60 days later) at 200 lbs. per acre equal to 4.6# / 1,000 sq. ft. | For use in Bluegrass, Bentgrass, Perennial Ryegrass and Tall Fescue. Make application in the fall after turf has become dormant. For late fall applications, seeding into treated areas should be delayed until 4 months after greenup. |
| Annual Bluegrass Control in Bermudagrass Fairways Overseeded with Perennial Ryegrass only | | | |
| Annual Bluegrass (Poa annua spp.) | 200 lbs. / A | 4.6# lbs. / 1,000 sq. ft. | Apply in late summer to early fall prior to Poa Annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use when Poa trivialis is present in the overseed mixture. Apply the full 200 lb. rate per acre in one application. |
| <p>NOTE: In many areas annual bluegrass exists in at least two biotypes or subspecies of Poa annua. The true <u>annual</u> biotype, "Poa annua spp. annua" which germinates from seed every year <u>is controlled</u> while the biotype, "Poa annua spp. raptans" which behave like a <u>perennial</u> and survives through the summer as an established plant, <u>cannot be controlled</u>. Consult the Extension service or University weed specialist in your area for more information concerning your locale.</p> | | | |

| Annual Weed Control during establishment from Sprigs of Bermudagrass, Zoysiagrass and Seashore Paspalum | | | |
|--|--|---|--|
| Crabgrass and Goosegrass, Annual Sedge and other annual weeds. | Apply 40-60 days before germination at 200 lbs. / per acre | Apply 200 lbs. of #3 / per acre; 60 days later. | |
| For Bermudagrass make application immediately prior to or immediately after sprigging. For Seashore Paspalum make application 10 to 14 days after sprigging. Applications made close to the time of sprigging may cause objectionable Seashore Paspalum injury. Disturbance of the soil surface after application may disrupt the herbicide barrier and reduce effectiveness. For best results sprinkler irrigate the "sprigged" area with up to ¼ inch water after application. | | | |

PRECAUTIONS IN USING: For St. Augustine grass and Bentgrass turf. DO NOT exceed the rate of one application of each STARTEEM #1 and STARTEEM #2 or #3 per 1,000 sq. ft. / year. DO NOT apply to Bentgrass turf mowed less than 3/8 inches. This product may cause discoloration of Bentgrass, Bermudagrass and St. Augustinegrass but this discoloration is temporary and normally outgrown within 2 to 3 weeks.

20030530.Final Label

FERTILIZER WITH STARTEEM® # 3

A NEW! PRE-EMERGENCE HERBICIDE COMBINATION FOR GOLF COURSES, PARKS, ATHLETIC FIELDS, CEMETERIES, UNIVERSITY, FEDERAL, MUNICIPAL, CORPORATE LAWNS and SOD FARMS

Avoid contact with flowers and shrubs except as recommended elsewhere for ornamental plants on this label. DO NOT apply to newly seeded areas. Seeding into treated areas should be delayed until 4 months after treatment except as noted above. DO NOT apply to Dichondra or Centipedegrass. DO NOT use on golf course putting greens or tees because of the varying cultural practices used.

FOR SPRING APPLICATION ON GOLF COURSES UNDER TURF MANAGEMENT

Use the STARTEEM-SPLIT application!

Weed control and better turf management can be programmed by using the "STARTEEM-SPLIT" weed control program. Starting 40 to 60 days before expected weed seed germination, apply STARTEEM #1 at a rate of 200 lbs. per acre and follow within 60 days with an application of STARTEEM #2 at the 200 lb. per acre rate. This program supplies 2.5 lbs. of Oxadiazon per acre, per year and 2.5 lbs. of the combination of the Benefin and Trifluralin per acre, per year.

If "Hard-To-Kill" weeds (as listed below) with the asterisk (*) are known to be prevalent in the turfgrass area, use STARTEEM® #3 as the second application after STARTEEM® #1 applied at 200 lbs. per acre (equal to the 4.6 lbs per 1,000 sq ft)

Controls Annual Broadleaf Weeds and Annual Grasses

| | | | |
|------------------|-------------------------|------------------------|------------------------|
| Annual Bluegrass | *Evening Primrose | **Jungle Ricegrass | Sow Thistle |
| Annual Sedge | Fireweed | Lambsquarter | Speedwell |
| *Barnyardgrass | Field Sandbur | **Liverwort | Spotted Catsear |
| Bittercress | Foxtail; Green, Bristly | Oxalis | Spurge; Garden, Petty' |
| Carpetweed | Fall Panicum | **Panama Paspalum | **Prostrate |
| Carpetgrass | Florida Pusley | Pennsylvania Smartweed | Stinging Nettle |
| Common Groundsel | **Galinsoga | Redroot Pigweed | *Swinecress |
| Common Purslane | Golden Ragwort | Sensitive Plant | Virginia Buttonweed |
| Crabgrass | *Goosegrass | Shepardspurse | Yellow Woodsorrel |

Use the STARTEEM® #1 and STARTEEM® #2 SPLIT application for control of these weeds listed above not bearing the asterisk (). For hard-To-Kill weeds use STARTEEM #3 as second application.

NOTE: For newly seeded lawns do not apply this product until after the second or third cutting. For best turf management, place new grass seed in the fall of the year. Spring pre-emergence weed control can then be applied.

SPREADER SETTINGS

| Spreader Settings for | STARTEEM® #1 | STARTEEM® #2 | STARTEEM® #3 |
|-----------------------|--------------|--------------|--------------|
| Cyclone B3-Pro | #x.x | #x.x | #x.x |
| For Scotts Pro R-8A | #x.x | #x.x | #x.x |

All STARTEEM products #1; #2; and #3 apply at 200 lbs / Acre = to 4.6# / 1,000 sq. ft.

Plan on an approximately 3 mph spreader speed to obtain best coverage. Know the exact square footage of your area before application. Incorrect application can result in turf damage or poor weed control.

FOR SPRING APPLICATION TO UNIVERSITY, FEDERAL, CORPORATE, MUNICIPAL LAWNS, CEMETERIES, PARKS, ATHLETIC FIELDS, SOD FARMS, CONDOMINIUMS, UNDER MANAGEMENT. FOR PRE-EMERGENCE WEED CONTROL, use a STARTEEM®-SPLIT PROGRAM!

STARTEEM® may be applied to "established" varieties of perennial Bluegrass, Bentgrass (see restrictions), Bermudagrass, Buffalograss, Perennial Ryegrass, St. Augustinegrass, Seashore Paspalum, Tall Fescue, and Zoysiagrass. Weed control and better turf management can be programmed by using the "STARTEEM-SPLIT" weed control program. Starting 40 to 60 days before expected weed seed germination, apply STARTEEM® #1 at a rate of 4.6 lbs per 1,000 sq. ft. and follow with a second application using STARTEEM® #2 at the same rate; within 60 days. This program supplies 2.5 lbs of Oxadiazon per acre, per year and 2.5 lbs of the combination of the Benefin and Trifluralin per acre, per year.

If "Hard-To-Kill" weeds, as listed below with the asterisk (*) are known to be prevalent in the area, use STARTEEM® #3 as the second application after STARTEEM® #1 applied at 200 lbs per acre (equal to the 4.6 lbs per 1,000 sq. ft.)

WHEN TO APPLY AFTER OVERSEEDING TURF

Do not apply to over-seeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Delay overseeding or reseeding 60 to 90 days.

FERTILIZER WITH STARTEEM® # 3

A NEW! PRE-EMERGENCE HERBICIDE COMBINATION FOR GOLF COURSES, PARKS, ATHLETIC FIELDS, CEMETERIES, UNIVERSITY, FEDERAL, MUNICIPAL, CORPORATE LAWNS and SOD FARMS


Controls Annual Broadleaf Weeds and Annual Grasses

Annual Bluegrass
 Annual sedge
 *Barnyardgrass
 Bittercress
 Carpetweed
 Carpetgrass
 Common Groundsel
 Common Purslane
 Crabgrass
 *Evening Primrose
 Fireweed
 Field Sandbur

Fall Panicum
 Florida Pusley
 *Galinsoga
 Golden Ragwort
 *Goosegrass
 **Jungle Ricegrass
 Lambsquarter
 **Liverwort
 Oxalis
 **Panama Paspalum
 Pennsylvania Smartweed

Redroot Pigweed
 Sensitive plant
 Shepardspurse
 Sow Thistle
 Speedwell
 Spotted Catsear
 Spurge, Garden, Petty,
 **Prostrate
 Stinging Nettle
 *Swinecress
 Virginia Buttonweed
 Yellow Woodsorrel

Approximate Crabgrass Seed Germination Dates In the U.S.A.



- Approximate Date
- After May 30
 - After May 10
 - After April 20
 - After March 20
 - January 1 to March 20

NOTICE: Buyer and user assume all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

Net Weight 50 Pounds

Harrell's, Inc.
 P. O. Box 807
 Lakeland, FL 33802