

52287-15

10/14/2008

1/4



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OCT 14 2008

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Mr. Robert F. Bischoff
Agent, TotalTurf Consulting LLC
Harrell's Inc.
P.O. Box 807
Lakeland, FL 33802

Dear Mr. Bischoff:

Subject: Harrell's Granular Herbicide 75
Label Amendment
EPA Registration No. 52287-15

The labeling referred to above, submitted in accordance with registration under the Federal Insecticide, Fungicide, and Rodenticide act, as amended is acceptable.

Submit 1 (one) copy of your final printed labeling before you release the product for shipment. A stamped copy of the label is enclosed for your records.

Sincerely,

A handwritten signature in black ink that reads "Joanne I. Miller".

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505P)

ACCEPTED

2/14

OCT 14 2008

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

(Base Label):

(logo) Harrell's, LLC

57287-15

HARRELL'S GRANULAR HERBICIDE 75

A selective preemergence herbicide for control of certain annual grasses and broadleaf weeds in:

- Container Grown Ornamentals
- Landscape Ornamentals
- Nursery Stock

Active Ingredients:

Oxyfluorfen.....	2.0%
Trifluralin.....	3.0%
Inert Ingredients.....	95.0%
Total.....	100.0%

Contains 2.5 pounds active ingredient per 50 pound bag

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Keep Out of Reach of Children

Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

Personal Protective Equipment (PPE)

Mixers, loaders, applicators and other handlers must wear:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves
- Chemical-resistant apron for mixers and loaders

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs or in a manner that meets with requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d) (4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advise. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advise.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advise.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advise.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

Environmental Hazards

This product is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates including shrimp and oyster and is toxic to wildlife. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash water or rinsate. Use with care when applying in areas frequented by wildlife or adjacent to any body of water or wetland area. Do not apply when weather conditions favor drift or erosion from target areas. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. See Directions for Use for additional restrictions.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

EPA Reg. No. 52287-15

EPA Est. No. 52287-SC-01

Harrell's, LLC • Lakeland, FL 33802, U.S.A.

Herbicide

Net Contents 50 Pounds

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Non-Agricultural Use Requirements

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter the treated area until dusts have settled. If soil incorporation is required after the application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If the incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container only. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General Information

Harrell's Granular Herbicide 75 herbicide is a preemergence herbicide for control of certain annual grasses and broadleaf weeds in container grown ornamentals, landscape ornamentals and nursery stock. Apply Harrell's Granular Herbicide 75 before or after planting but prior to germination of target weeds, or immediately after cultivation. Length of weed control will vary with weed population, potting media or soil conditions, temperature, watering regime, and other factors. Following application, user should monitor and observe level of weed control over time to determine when additional applications may be needed. Repeat applications should not be made sooner than 60 days after a previous application of Harrell's Granular Herbicide 75.

Use Precautions

Harrell's Granular Herbicide 75 does not control established weeds. Existing weeds should be controlled by cultivation or with postemergence herbicides. Weed residues, prunings and trash should be removed or thoroughly mixed into soil prior to treatment. Soil should be in good condition and free of clods at the time of application. Do not apply when wind conditions favor drift of Harrell's Granular Herbicide 75 granules from the target area. Optimum weed control will be obtained when followed by ½ to 1 inch overhead irrigation or rainfall within 3 to 4 days after surface application.

Weeds controlled by Harrell's Granular Herbicide 75 when applied at 100 lb/acre (2.3 lb/1000 sq ft)

Common Name	Scientific Name
AMARANTH, SPINY	<i>Amaranthus spinosus</i>
BARLEY, HARE	<i>Hordeum leporinum</i>
BARNYARDGRASS (WATERGRASS)	<i>Echinochloa crus-galli</i>
BEDSTRAW, CATCHWEED	<i>Galium aparine</i>
BITTERCRESS, LESSER	<i>Cardamine oligosperma</i>
BLUEGRASS, ANNUAL	<i>Poa annua</i>
BROME	<i>Bromus spp.</i>
BUCKWHEAT, WILD	<i>Polygonum convolvulus</i>
BURCLOVER	<i>Medicago hispida</i>
CANARYGRASS (ANNUAL)	<i>Phalaris canariensis</i>
CARPETWEED	<i>Mollugo verticillata</i>
CHICKWEED, COMMON	<i>Stellaria media</i>
CLOVER, RED	<i>Trifolium pratense</i>
CLOVER, WHITE	<i>Trifolium repens</i>
COCKLEBUR, COMMON	<i>Xanthium pensylvanicum</i>
CRABGRASS	<i>Digitaria spp.</i>
FIDDLENECK, COAST	<i>Amsinckia intermedia</i>
FILAREE, BROADLEAF	<i>Erodium botrys</i>
FILAREE, REDSTEM	<i>Erodium cicutarium</i>
FILAREE, WHITESTEM	<i>Erodium moschatum</i>
FIREWEED (FROM SEED)	<i>Epilobium angustifolium</i>
FLIXWEED	<i>Descurainia sophia</i>
FOXTAIL, GIANT	<i>Setaria faberi</i>
FOXTAIL, GREEN	<i>Setaria viridis</i>
FOXTAIL, YELLOW	<i>Setaria lutescens</i>
GOOSEFOOT, NETTLELEAF	<i>Chenopodium murale</i>
GOOSEGRASS	<i>Eleusine indica</i>

GROUNDCHERRY, CUTLEAF	<i>Physalis angulata</i>
GROUNDCHERRY, WRIGHT	<i>Physalis wrightii</i>
GROUNDSEL, COMMON	<i>Senecio vulgaris</i>
HENBIT	<i>Lamium amplexicaule</i>
JIMSONWEED	<i>Datura stramonium</i>
JUNGLERICE	<i>Echinochloa colonum</i>
JOHNSONGRASS, SEEDLING	<i>Sorghum halepense</i>
KNOTWEED, PROSTRATE	<i>Polygonum aviculare</i>
KOCHIA	<i>Kochia scoparia</i>
LADYSTHUMB (SMARTWEED)	<i>Polygonum persicaria</i>
LAMBSQUARTERS, COMMON	<i>Chenopodium album</i>
LETTUCE, PRICKLY (CHINA LETTUCE)	<i>Lactuca serriola</i>
LOVEGRASS	<i>Eragrostis spp.</i>
MALLOW, LITTLE (MALVA)	<i>Malva parviflora</i>
MAYWEED (DOG FENNEL)	<i>Anthemis cotula</i>
MINERSLETTUCE	<i>Montia perfoliata</i>
MORNINGGLORY SPECIES, ANNUAL	<i>Ipomoea species</i>
MORNINGGLORY, IVYLEAF	<i>Ipomoea hederacea</i>
MORNINGGLORY, TALL	<i>Ipomoea purpurea</i>
MUSTARD, BLUE (PURPLE MUSTARD)	<i>Chorispora tenella</i>
MUSTARD, TUMBLE (JIM HILL MUSTARD)	<i>Sisymbrium altissimum</i>
MUSTARD, WILD	<i>Brassica kaber</i>
NETTLE, BURNING	<i>Urtica urens</i>
NETTLE, STINGING	<i>Urtica dioica</i>
NIGHTSHADE, AMERICAN BLACK	<i>Solanum americanum</i>
NIGHTSHADE, BLACK	<i>Solanum nigrum</i>
NIGHTSHADE, HAIRY	<i>Solanum sarrachoides</i>
OATS, WILD	<i>Avena fatua</i>
ORACH, RED	<i>Atriplex rosea</i>
PANICUM, FALL	<i>Panicum dichotomiflorum</i>
PANICUM, TEXAS	<i>Panicum texanum</i>
PEPPERWEED, YELLOWFLOWER	<i>Lepidium perfoliatum</i>
PIGWEEED, PROSTRATE	<i>Amaranthus blitoides</i>
PIGWEEED, REDROOT	<i>Amaranthus retroflexus</i>
PIMPERNEL, SCARLET	<i>Anagallis arvensis</i>
PURSLANE, COMMON	<i>Portulaca oleracea</i>
PUSLEY, FLORIDA	<i>Richardia scabra</i>
REDMAIDS	<i>Calandrinia caulescens</i>
ROCKET, LONDON	<i>Sisymbrium irio</i>
SANDBUR, FIELD	<i>Cenchrus incertus</i>
SANDSPURRY, RED	<i>Spergularia rubra</i>
SHEPHERDSPURSE	<i>Capsella bursa-pastoris</i>
SIDA, PRICKLY (TEAWEED)	<i>Sida spinosa</i>
SIGNALGRASS	<i>Brachiaria spp.</i>
SMARTWEED, PENNSYLVANIA	<i>Polygonum pennsylvanicum</i>
SORREL, RED (FROM SEED)	<i>Rumex acetosella</i>
SOWTHISTLE, ANNUAL	<i>Sonchus oleraceus</i>
SPURGE, PROSTRATE	<i>Euphorbia supina</i>
SPURGE, SPOTTED	<i>Euphorbia maculata</i>
SPURRY, CORN	<i>Spergula arvensis</i>
STINKGRASS	<i>Eragrostis cilianensis</i>
TANSYMUSTARD	<i>Descurainia pinnata</i>

THISTLE, BULL	<i>Cirsium vulgare</i>
THISTLE, RUSSIAN	<i>Salsola kali</i>
VELVETLEAF	<i>Abutilon theophrasti</i>
WITCHGRASS	<i>Panicum capillare</i>
WOODSORREL, COMMON YELLOW	<i>Oxalis stricta</i>

Application Directions

Apply Harrell's Granular Herbicide 75 using a drop or rotary-type spreader designed to apply granular herbicides or insecticides. Calibrate application equipment prior to use according to manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or plant injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first.

Application Techniques for Applying Harrell's Granular Herbicide 75

When using a drop-type spreader, a splash board mounted under the hopper will provide more even granule distribution.

A chain fastened to the side of the spreader and allowed to drag on the soil surface can be used to mark the edge of the treated swath and help prevent skips or overlaps.

For treating smaller areas or rows of nursery stock or ornamental beds, a hand held or push-type rotary applicator such as whirlybird or cyclone unit is recommended. For hand held units, walk and turn the crank at a constant rate of speed.

Uses

Container Grown Ornamentals, Landscape Ornamentals and Nursery Stock

Harrell's Granular Herbicide 75 is recommended as a preemergence treatment for control of certain annual grasses and broadleaf weeds in container grown ornamentals, landscape ornamentals, and nursery stock. Apply Harrell's Granular Herbicide 75 before or after planting but prior to germination of target weeds, or immediately after cultivation.

Refer to the "General Information" section of this label for general use precautions and information on application, application rates and weeds controlled.

Special Use Restrictions and Precautions

- Do not apply more than 2 lbs of oxyfluorfen active ingredient per acre for each application on all container grown ornamentals or more than 6 lbs of oxyfluorfen active ingredient per acre per season.
- To avoid possible injury, do not apply Harrell's Granular Herbicide 75 to:
 - nursery forest or Christmas tree seedling beds, cutting beds, or transplant beds
 - unrooted liners or cuttings that have been planted in pots for the first time
 - pots less than four inches wide
- Do not apply Harrell's Granular Herbicide 75 to newly transplanted ornamentals and nursery stock until soil or potting media has been settled by packing and irrigation or rainfall and no cracks are present or injury may occur.
- Do not apply Harrell's Granular Herbicide 75 to wet or dew covered foliage. Should this happen, irrigate immediately to completely remove herbicide particles from leaf surfaces.

- Do not apply Harrell's Granular Herbicide 75 to whorl-leaved plants as their leaves may channel the herbicide granules to the leaf base and may result in foliage injury.
- Harrell's Granular Herbicide 75 should not be applied two (2) weeks prior to or two (2) weeks following leaf bud break or during periods of flush growth. Newly forming leaves are particularly sensitive to this product and spotting or leaf drop may result from such applications.
- Do not apply Harrell's Granular Herbicide 75 within two (2) weeks prior to shipping.
- Do not apply Harrell's Granular Herbicide 75 in greenhouse or other enclosed structures. Do not apply within two (2) weeks prior to enclosure in greenhouse or other structures.
- Repeat applications should not be made sooner than 60 days after a previous application of Harrell's Granular Herbicide 75.

Users who wish to use Harrell's Granular Herbicide 75 on plant species not recommended on this label may determine suitability for such uses by making trial application of Harrell's Granular Herbicide 75 at a recommended rate to a small number of plants. Prior to using Harrell's Granular Herbicide 75 on a larger number of plants, the treated plants should be observed for signs of herbicidal injury during 30 to 60 days of normal growing conditions to determine if the treatment is non-injurious to the target plant species. **The user assumes responsibility for any plant damage or other liability resulting from the use of Harrell's Granular Herbicide 75 on plant species not recommended on this label.**

Harrell's Granular Herbicide 75 May Be Used On The Following Established Plant Species When Container Grown or Field Grown:

Trees

Scientific Name	Common Name
<i>Abies balsamea</i>	Balsam fir
<i>Abies concolor</i>	White fir
<i>Abutilon hybridum</i>	Albus-flowering maple
	Luteus-flowering maple
	Roseus-flowering maple
	Tangerine-flowering maple
	Vesuvius red-flowering maple
<i>Acer ginnala</i>	flame maple
<i>Acer platanoides</i>	Norway maple
<i>Acer rubrum</i>	Red maple
	Red sunset maple
<i>Acer saccharinum</i>	Silver maple
<i>Acer saccharum</i>	Sugar maple
<i>Areacastrum romanzoffianum</i>	Queen palm
<i>Betula nigra</i>	River birch
<i>Betula papyrifera</i>	Paper birch
<i>Betula pendula</i>	European white birch
<i>Brachychiton populneus</i>	Bottle tree
<i>Bucida buceras</i>	Black olive
<i>Castanea mollissima</i>	Chinese chestnut
<i>Ceratonia siliqua</i>	Carob
<i>Cercis canadensis</i>	Redbud
<i>Chamaecyparis obtusa</i>	Fillicoides-fernspray cypress
	Gracilis-slender Hinoki cypress
<i>Chamaecyparis pisifera</i>	Swara false cypress
	Squarrosa-moss cypress
<i>Chamaedorea cataractarum</i>	Palm

<i>Chamaedorea costaricana</i>	Palm
<i>Chamaedorea elegans</i>	Parlor palm
<i>Cornus florida</i>	Cloud nine dogwood
	Flowering dogwood
<i>Cornus kousa</i>	Dogwood, kousa
<i>Crataegus viridis</i>	Green hawthorn
<i>Cupaniopsis anacardioides</i>	Carrot wood
<i>Cupressus glabra</i>	Arizona cypress
<i>Elaeagnus angustifolia</i>	Russian olive
<i>Eucalyptus camaldulensis</i>	Redgum eucalyptus
<i>Eucalyptus cinerea</i>	Mealy eucalyptus
	Silver dollar eucalyptus
<i>Eucalyptus microtheca</i>	Coolibah tree
<i>Eucalyptus sideroxylon</i>	Red ironbark eucalyptus
<i>Ficus benjamina</i>	Ficus
	Mini ficus
<i>Fraxinus americana</i>	White ash
<i>Fraxinus udhei</i>	Shamel ash
<i>Ginkgo biloba</i>	Ginkgo-maidenhair tree
<i>Gleditsia triacanthos</i>	Honey locust
	Shademaster honey locust
<i>Heteromeles arbutiflora</i>	Toyon
<i>Illicium floridanum</i>	Florida anise-tree
<i>Juniperus virginiana</i>	Eastern red cedar
<i>Larix kaempferi</i>	Japanese larch
<i>Liquidambar styraciflua</i>	American sweet gum
<i>Liriodendron tulipifera</i>	Tuliptree
<i>Magnolia spp.</i>	Magnolia
<i>Malus spp.</i>	Crabapple
<i>Morus alba</i>	White mulberry
<i>Musa aluminata</i>	Banana
<i>Nyssa sylvatica</i>	Blackgum
<i>Oxydendrum arboreum</i>	Sourwood
<i>Picea abies</i>	Norway spruce
	Pendula-weeping Norway spruce
	Repens-spreading Norway spruce
<i>Picea glauca conica</i>	Dwarf Alberta spruce
<i>Picea glauca</i>	White spruce
<i>Picea pungens</i>	Dwarf globe blue spruce
	Glauca-Colorado blue spruce
	Hoopsii-Hoop's blue spruce
	Koster-Koster blue spruce
<i>Pinus aristata</i>	Bristlecone pine
<i>Pinus canariensis</i>	Canary Island pine
<i>Pinus contorta</i>	Shore pine, beach pine
<i>Pinus eldarica</i>	Eldarica pine
<i>Pinus leucodermis</i>	Bosnian pine
<i>Pinus mugo</i>	Pumilio-shrubby Swiss mountain pine
<i>Pinus nigra</i>	Austrian black pine
<i>Pinus radiata</i>	Monterey pine
<i>Pinus resinosa</i>	Red pine
<i>Pinus taeda</i>	Loblolly pine
<i>Pinus strobus</i>	White pine
<i>Pinus sylvestris</i>	Columnar Scotch pine
	Scotch pine
<i>Pinus thunbergiana</i>	Japanese black pine

<i>Platanus acerifolia</i>	London planetree
<i>Platanus occidentalis</i>	American sycamore
<i>Platanus racemosa</i>	Califorina sycamore
<i>Podocarpus</i> spp.	Podocarpus
<i>Populus deltoides</i>	Cottonwood
<i>Prosopis chilensis</i>	Chilean mesquite
<i>Prunus yedoensis</i>	Yoshino flowering cherry
<i>Pseudotsuga menziesii</i>	Douglasfir
<i>Quercus coccinea</i>	Scarlet oak
<i>Quercus ilicifolia</i>	Bear oak
<i>Quercus palustris</i>	Pin oak
<i>Quercus phellos</i>	Willow oak
<i>Quercus rubra</i>	Red oak
<i>Quercus virginiana</i>	Live oak
<i>Robinia pseudoacacia</i>	Black locust
<i>Salix</i> spp.	Willow
<i>Sequoiadendron giganteum</i>	Giant sequoia
<i>Swietenia mahogani</i>	Mahogany
<i>Tabebuia caraiba</i>	Yellow tab
<i>Taxodium distichum</i>	Baldcypress
<i>Tilia</i> spp.	Linden
<i>Tsuga canadensis</i>	Eastern hemlock
<i>Ulmus parvifolia</i>	Chinese elm
<i>Washingtonia robusta</i>	Mexican fan palm

Ornamental Shrubs

Common Name	Scientific Name
<i>Abelia grandiflora</i>	Edward Goucher abelia
	Glossy abelia
<i>Acacia abyssinica</i>	Abyssinica acacia
<i>Acacia redolens</i>	Prostrate acacia
<i>Acacia stenophylla</i>	Shoestring acacia
<i>Acalypha wilkesiana</i>	Copper leaf
<i>Acer ginnala</i>	Amur maple
<i>Acer palmatum</i>	Coral bark Japanese maple
	Dwarf Japanese maple
<i>Agave americana</i>	Century plant
<i>Astilbe chinensis</i>	False spiraea
<i>Athyrium nipponicum</i>	Japanese painted fern
<i>Baccharis pilularis</i>	Coyotebush
<i>Berberis gladwynensis</i>	William Penn barberry
<i>Berberis mentorensis</i>	Mentor barberry
<i>Berberis thunbergii</i>	Atropurea-redleaf Japanese barberry
	Aurea-golden Japanese barberry
	Crimson pygmy barberry
	Rose glow barberry
<i>Bougainvillea</i> spp.	Barbara Karst
	California gold
	Pink pixie
	Scarlet O'Hara
	Temple fire
	Texas dawn
<i>Buxus microphylla japonica</i>	Japanese boxwood
<i>Buxus microphylla Koreana</i>	Korean boxwood
<i>Buxus sempervirens</i>	Common boxwood

<i>Callistemon citrinus</i>	Lemon bottlebrush
<i>Callistemon viminalis</i>	Weeping bottlebrush
<i>Calluna vulgaris</i>	Spring torch scotch heather
<i>Camellia sasanqua</i>	Sasanqua camellia
<i>Camellia japonica</i>	Japanese camellia
<i>Cassia artemisioides</i>	Feathery cassia
<i>Ceanothus</i> spp.	Wild lilac
<i>Cephalotaxus drupacae</i>	Plum yew
<i>Cerastium tomentosum</i>	Snow-in-summer
<i>Chamaecyparis obtusa</i> spp.	Kosteri cypress
	Nana-dwarf Hinoki cypress
	Torulosa cypress
<i>Chamaecyparis pisifera</i>	Filifera-thread cypress
<i>Chrysalidocarpus lutescens</i>	Areca palm
<i>Clethra alnifolia</i>	Summersweet
<i>Cleyera japonica</i>	Japanese cleyera
<i>Cornus alba</i>	Sibirica-Siberian dogwood
<i>Cornus stolonifera</i>	Baileyi red osier dogwood
	Flaviramea-yellowtwig dogwood
<i>Cotinus coggygria</i>	Royal purple smoke tree
<i>Cotinus dammeri</i>	Coral beauty smoke tree
	Eichholz smoke tree
<i>Cotoneaster adpressus</i>	Praecox-early cotoneaster
<i>Cotoneaster apiculatus</i>	Cranberry cotoneaster
<i>Cotoneaster congestus</i>	Pyrenees cotoneaster
<i>Cotoneaster dammeri</i>	Bearberry cotoneaster
<i>Cotoneaster himalayan</i>	Himalayan cotoneaster
<i>Cotoneaster horizontalis</i>	Rock cotoneaster
<i>Cotoneaster zabelii</i>	Zabel cotoneaster
<i>Cycas revoluta</i>	Sago palm
<i>Cytisus praecox</i>	Hollandia-warminster broom
<i>Cytisus scoparius</i>	Lena-Scotch broom
<i>Daphne odora</i>	Fragrant daphne
<i>Deutzia</i> spp.	Deutzia
<i>Dodonea viscosa</i>	Hopseed bush
<i>Elaeagnus pungens</i>	Fruitland silver berry
<i>Erica cinerea</i>	Purple bell heather
<i>Erica vagans</i>	Cornish heather
<i>Euonymus fortunei</i>	Canadale gold euonymus
	Emerald'n gold euonymus
	Sunspot euonymus
	Wintercreeper euonymus
<i>Euonymus japonica</i>	Silver king-euonymus
	Variegated evergreen euonymus
<i>Euonymus kiautschovica</i>	Spreading euonymus
<i>Feijoa sellowiana</i>	Pineapple guava
<i>Forsythia</i> spp.	Forsythia
<i>Gardenia jasminoides</i>	August beauty gardenia
	Gardenia
	Radican gardenia
<i>Gaultheria shallon</i>	Salal/lemon leaf
<i>Gelsemium sempervirens</i>	Carolina jessamine
<i>Genista pilosa</i>	Woadwaxen
<i>Hibiscus rosa-sinensis</i>	Ross Estey-hibiscus
<i>Hibiscus syriacus</i>	Rose of Sharon-heart
	Rose of Sharon-red bird

<i>Ilex</i> spp.	Rose of Sharon-woodbridge
<i>Illicium annisatum</i>	Holly
<i>Itea ilicifolia</i>	Mystery gardenia
<i>Ixora collinea</i>	Henry Garnet holly leaf sweetspire
<i>Juniperus</i> spp.	Ixora
<i>Kalmia latifolia</i>	Juniper
<i>Lagerstroemia indica</i>	Mountain laurel
<i>Lantana</i> spp.	Crape myrtle
<i>Leucothoe axillaris</i>	Lantana
<i>Leucothoe fontainesiana</i>	Coast leucothoe
<i>Ligustrum</i> spp.	Drooping leucothoe
<i>Livistona chinensis</i>	Privet
<i>Lonicera periclymenum</i>	Chinese fountain palm
	Flowering woodbine
<i>Lonicera sempervirens</i>	Serotina woodbine
<i>Mahonia bealei</i>	Trumpet honeysuckle
<i>Mahonia repens</i>	Leather leaf mahonia
<i>Myrica cerifera</i>	Creeping mahonia
<i>Nandina domestica</i>	Wax myrtle
	Compact-dwarf heavenly bamboo
	Harbour dwarf-heavenly bamboo
	Heavenly bamboo
	Nana compacta-heavenly bamboo
	Nana purpurea-heavenly bamboo
	Woods dwarf heavenly bamboo
<i>Nerium oleander</i>	Hardy red oleander
	Oleander
	Ruby lace oleander
<i>Osmanthus fortunei</i>	Fortunes osmanthus
<i>Philadelphus</i> spp.	Mockorange
<i>Phoenix roeloenii</i>	Pigmy date palm
<i>Photinia fraseri</i>	Fraser's photinia
<i>Pieris japonica</i>	Japanese andromeda
	Mountain fire lily-of-the-valley
	Snowdrift lily-of-the-valley
	Templebells lily-of-the-valley
	Valley rose lily-of-the-valley
	Valley valentine lily-of-the-valley
<i>Pieris japonica x forestii</i>	Forest flame lily-of-the-valley
<i>Pinus mugo</i>	Mugo-mugho pine
<i>Pittosporum tobira</i>	Green pittosporum
	Japanese pittosporum
	Wheeler's dwarf pittosporum
<i>Plumbago ariculata</i>	Blue cape plumbago
<i>Plumbago capensis</i>	Plumbago
<i>Podocarpus macrophyllus</i>	Yewpine
<i>Polystichum polyblepharum</i>	Tassel fern
<i>Potentilla</i> spp.	Cinquefoil
<i>Prunus caroliniana</i>	Carolina laurel cherry
<i>Prunus glandulosa</i>	Dwarf pink flowering almond
<i>Pyracantha</i> spp.	Pyracantha
<i>Raphiolepis indica</i>	Charisma-Monruce raphiolepis
	Enchantress-Moness raphiolepis
	India hawthorn
	Springtime-Monme raphiolepis
<i>Raphiolepis ovata</i>	Roundleaf raphiolepis

<i>Rhododendron</i> spp.	Azalea/rhododendron
<i>Rhus lancea</i>	Africa sumac
<i>Rosa rugosa</i>	Ramanas rose
<i>Rosmarinus officinalis</i>	Rosemary
<i>Skimmia japonica</i>	Japanese skimmia
<i>Skimmia revesiana</i>	Reeve's skimmia
<i>Spiraea bumalda</i>	Anthony Waterer spiraea
<i>Spiraea japonica</i>	Dolchia spiraea
	Japanese alpine spiraea
	Shiobana spiraea
<i>Spiraea vanhouttii</i>	Bridal wreath
<i>Syringa rothomagensis</i>	Chinese lilac
<i>Syringa vulgaris</i>	Common lilac
<i>Taxus cuspidata</i>	Upright Japanese yew
<i>Taxus media</i>	Anglojap yew
<i>Tecomaria capensis</i>	Cape honeysuckle
<i>Ternstroemia gymnanthera</i>	
<i>Thuja occidentalis</i>	American arborvitae
	Emerald arborvitae
	Globosa-globe arborvitae
	Little giant-dwarf arborvitae
	Nigra-dark American arborvitae
	Pyramidalis-pyramid arborvitae
	Rheingold arborvitae
	Techny arborvitae
<i>Thuja orientalis</i>	Aureus nana-dwarf golden arborvitae
	Minima glauca-dwarf arborvitae
<i>Veitchia merrilli</i>	Christmas palm
<i>Viburnum</i> spp.	Viburnum
<i>Weigela</i> spp.	Weigela
<i>Xylosma congestum</i>	Xylosma
<i>Yucca filamentosa</i>	Yucca

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, to the extent consistent with applicable law, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. To the extent consistent with applicable law, all such risks shall be assumed by buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer and Inherent Risks of Use above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.