

52287-16

9/28/2004

1/2



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
52287-16

Date of Issuance:
SEP 28 2004

Term of Issuance:
Conditional

Name of Pesticide Product:
Weedfree 63 Herbicide

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Harrell's Inc.
P.O. Box 807
Lakeland, FL 33802

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is registered in accordance with FIFRA sec. 3(c) (7) (A) provided that you:

1. Add the phrase "EPA Registration No.52287-16" to the label before you release the product for shipment.

COMMENTS CONTINUED ON PAGE 2 OF THIS NOTICE OF REGISTRATION

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product under the enclosed stamped copy of the label constitutes acceptance of these conditions.

Enclosure

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Signature of Approving Official:

151

Date:

SEP 28 2004

412

2. Submit the following data required for the registration of this pesticide product within 1 year of the date of this Notice of Registration:

<u>EPA Guideline Data Number</u>	<u>Guideline Descriptor</u>
830.6317	Storage Stability Study
830.6320	Corrosion Characteristics

3. Revise the "Ingredient Statements" by adding the CAS Number of the active ingredient, oxyfluorfen.
4. Following the Signal Word, CAUTION, add "Contains Petroleum Distillates".
5. Rewrite the "Personal Protective Equipment (PPE)" statement that reads "Chemical-resistant gloves" to read: "Chemical-resistant gloves, such as Barrier Laminate, Butyl Rubber, Nitrile Rubber, Viton, Barrier Laminate, (Category F, and G).
6. Revise the "First Aid" statements by adding the following: "NOTE TO PHYSICIAN: May pose an aspiration pneumonia hazard. Contains petroleum distillate." as last statement in the block.
7. The claim "grassy weeds" must be revised to read "weed grasses" on page 4 and 6 of the labeling.
8. Under "Conifer Transplants and Container Stock (includes 2-0 Seedling and Christmas Tree Plantings" delete the statement that reads: "Many container-grown conifers".
9. On page 8, tell the using-public when and where to use the range of dosages stated under "Dosage" on page 8.
10. On the front panel, either delete the descriptor "selective" or define it in terms of the labeling the follows.
11. Submit one (1) copy of the final printed labeling before you release this product for shipment.
12. Submit and/or cite all data required for the registration of this product when the Agency requires all registrants of similar products to submit data; and submit acceptable responses required for reregistration of this product under FIFRA, section 4.

A Stamped copy of the label is enclosed for your records.

Attachment

Base Label:

(Logo) Harrell's, Inc.

WEEDFREE 63 HERBICIDE

A selective preemergence herbicide for control of certain annual grasses and broadleaf weeds in:

- Container Grown Ornamentals
- Landscape Ornamentals
- Nursery Stock
- Conifer Seedbeds

**ACCEPTED
with COMMENTS
In EPA Letter Dated:
SEP 28 2004**

Active Ingredient:	
Oxyfluorfen.....	2%
Inert Ingredients.....	98%
Total.....	100%

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

52289-16

Contains 1 pound active ingredient per 50 pound bag.

Keep Out of Reach of Children

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing.

Wash thoroughly with soap and water after handling.

Personal Protective Equipment (PPE):

Mixers, loaders, applicators and other handlers must wear:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves
- Chemical-resistant apron for mixers and loaders

Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems and enclosed cabs in a manner that meets with requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d) (4-6), the handler PPI requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

Environmental Hazards

This product is highly toxic to aquatic invertebrates, aquatic plants, wildlife, and fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Use with care when applying in areas frequented by wildlife or adjacent to any body of water or wetland area. Do not apply when weather conditions favor drift or erosion from target areas. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. See Directions for Use for additional restrictions. Do not contaminate water when disposing of equipment wash water.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in original container away from other fertilizer, feed, or foodstuffs and separated from other pesticides.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label booklet. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

EPA Reg. No. 52287-

EPA Est. No. 52287-FL-01

Harrell's, Inc. • Lakeland, FL 33802, U.S.A.

Herbicide

Net Contents 50 Pounds

Directions for use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements:

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Non-Agricultural Use Requirements:

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Do not enter or allow others to enter until dusts have settled.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in original container away from other fertilizer, feed, or foodstuffs and separated from other pesticides.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL USE INFORMATION

Unless otherwise directed by registered supplemental labels, follow the Directions for Use in each crop group section.

CULTURAL CONSIDERATIONS

In order for Weedfree 63 to provide maximum preemergence activity:

Prior to application, the bed, soil, or growing media surface should be smooth and free of crop and weed trash (decaying leaves, clippings, dead weeds, etc.). Leaves and trash may be removed by blowing the area to be treated or by thoroughly mixing the trash into the soil through cultivation prior to herbicide application. After application, at least one-quarter inch (1/4 inch) of irrigation or rainfall should occur within 3 or 4 days after application. The best results from Weedfree 63 are from applications to established beds, soil, or growing media surfaces that are left undisturbed during the time period for which weed control is desired. Cultural practices that result in redistribution or disturbance of the soil surface after treatment will decrease the herbicidal effectiveness of Weedfree 63.

6/12

APPLICATION DIRECTIONS

Apply Weedfree 63 using a drop or rotary-type spreader designed to apply granular herbicides or insecticides. Calibrate application equipment prior to use according to manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or plant injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first.

Application Techniques for Applying Weedfree 63

When using a drop-type spreader, a splash board mounted under the hopper will provide more even granule distribution.

A chain fastened to the side of the spreader and allowed to drag on the soil surface can be used to mark the edge of the treated swath and help prevent skips or overlaps.

For treating smaller areas or rows of nursery stock or ornamental beds, a hand held or push-type rotary applicator such as whirlybird or cyclone unit is recommended. For hand held units, walk and turn the crank at a constant rate of speed.

CONIFER SEEDBEDS, TRANSPLANTS, CONTAINER STOCK and SELECTED FIELD GROWN DECIDUOUS TREES

Weedfree 63 is effective as a preemergence herbicide for the control of certain annual grassy and broadleaf weeds in conifer seedbeds, transplant and container stock, and in selected field-grown deciduous trees. Preemergence weed control is most effective when granules are applied to clean, weed-free soil surfaces. Treated soil surfaces should not be disturbed as the herbicidal effectiveness of Weedfree 63 may be decreased. Seedling weeds are controlled during emergence as they come in contact with the soil-applied herbicide.

IMPORTANT: When applied as directed, the conifer and selected deciduous species listed on this label have shown tolerance to Weedfree 63. It is impossible, however, to evaluate this product on all varieties, biotypes and cultivars of listed species on this label or under all possible growing conditions. The user should exercise reasonable judgment and caution with this product. Until familiar with results under user growing conditions, limit application of this product to a few plants in a small treated area to determine plant tolerance and extent of injury, if such occurs, prior to initiating large-scale applications.

WEEDS CONTROLLED

When Weedfree 63 is applied preemergence at recommended dosages and weed stages, the following grasses and broadleaf weeds are controlled.

† BARNYARDGRASS	MUSTARD, BLUE
BEDSTRAW, CATCHWEED	MUSTARD, TUMBLE
BITTERCRESS, LESSER	MUSTARD, WILD
† BLUEGRASS, ANNUAL	NETTLE, BURNING
BUCKWHEAT, WILD	NIGHTSHADE, BLACK
BURCLOVER	NIGHTSHADE, HAIRY
CARPETWEED	OATS, WILD
† CLOVER, RED	ORACH, RED
† CLOVER, WHITE	PEPPERWEED
	YELLOWFLOWER
COCKLEBUR, COMMON	PIGWEEED, PROSTRATE
† CRABGRASS, LARGE	PIGWEEED, REDROOT
† FIDDLENECK, COAST	PIMPERNEL, SCARLET
FILAREE, BROADLEAF	PURSLANE, COMMON

1/12

FILAREE, REDSTEM	REDMAIDS
FIREWEED (FROM SEED)	ROCKET, LONDON
FLIXWEED	SANDSPURRY, RED
† FOXTAIL, GIANT	† SHEPHERDSPURSE
† GOOSEGRASS	SIDA, PRICKLY
GROUNDCHERRY, CUTLEAF	SMARTWEED, PENNSYLVANIA
GROUNDCHERRY, WRIGHT	SORREL, RED (FROM SEED)
GROUNDSEL, COMMON	SOWTHISTLE, ANNUAL
HENBIT	SPEEDWELL, BIRDSEYE
JIMSONWEED	SPURGE, PROSTRATE
KNOTWEED, PROSTRATE	SPURGE, SPOTTED
LADYSTHUMB	SPURRY, CORN
LAMBSQUARTERS, COMMON	TANSYMUSTARD
LETTUCE, PRICKLY	THISTLE, BULL
MALLOW, LITTLE	THISTLE, RUSSIAN
MAYWEED	VELVETLEAF
MINERSLETTUCE	WITCHGRASS
† MORNINGGLORY, IVYLEAF	WOODSORREL, YELLOW
† MORNINGGLORY, TALL	

† Highest rate and/or multiple applications may be required for acceptable control.

Weedfree 63 is most effective when applied preemergence to annual grasses.

CONIFER SEEDBEDS

To assist in the establishment of conifer seedbeds, Weedfree 63 can be applied as a preemergence application following seeding. During periods of cool, cloudy weather, make certain that seedlings have hardened off prior to application.

Conifers are tolerant to preemergence applications of Weedfree 63. Weedfree 63 will provide residual preemergence control of many broadleaf weeds and annual grass species.

CONIFER SPECIES

Weedfree 63 may be applied to conifer seedbeds of species:

DOUGLAS FIR	<i>Pseudotsuga menziesii</i>
FIR	
FRASER	<i>Abies fraseri</i>
GRAND	<i>Abies grandis</i>
NOBLE	<i>Abies procera</i>
HEMLOCK	
EASTERN HEMLOCK	<i>Tsuga canadensis</i>

PINE

AUSTRIAN
EASTERN WHITE
HIMALAYAN
JACK
LOBLOLLY
LODGEPOLE
LONGLEAF
MONTEREY
MUGHO
PONDEROSA
SCOTCH
SHORTLEAF
SLASH
VIRGINIA

Pinus nigra
Pinus strobus
Pinus wallichiana
Pinus banksiana
Pinus taeda
Pinus contorta
Pinus palustris
Pinus radiata
Pinus mugo
Pinus ponderosa
Pinus sylvestris
Pinus echinata
Pinus elliottii
Pinus virginiana

SPRUCE

BLUE
DWARF ALBERTA
NORWAY
SITKA

Picea pungens
Picea glauca Conica
Picea abies
Picea sitchensis

PREEMERGENCE DOSAGE

Apply 12.5 to 50 lb. (0.25 lb. to 1.0 lb. active) of Weedfree 63 per broadcast acre as a preemergence application prior to conifer emergence. Where grassy weeds are present, a rate of 25 to 50 lb. (0.5 to 1.0 lb. active) of Weedfree 63 per broadcast acre is recommended. In known areas of high weed competition, 50 lb. (1.0 lb. active) of Weedfree 63 per broadcast acre are recommended.

TIMING AND METHOD OF APPLICATION

Broadcast Weedfree 63 to beds and irrigate prior to weed emergence with 1/2 to 3/4 inch of sprinkler irrigation.

CONIFER TRANSPLANTS AND CONTAINER STOCK (INCLUDES 2-0 SEEDLING AND CHRISTMAS TREE PLANTINGS)

Many container-grown conifers and conifer transplants are tolerant to preemergence and postemergence applications of Weedfree 63.

The following conifer species in addition to species listed under the CONIFER SEEDBED section have been shown to be tolerant to Weedfree 63.

ARBORVITAE

Thuja occidentalis
Thuja orientalis

RED CEDAR

Juniperus virginiana

JUNIPER

Juniperus chinensis
Juniperus horizontalis
Juniperus procumbens
Juniperus sabina
Juniperus scopulorum

WESTERN HEMLOCK

Tsuga heterophylla

YEW

Taxus species

Users who wish to use Weedfree 63 on plant species not recommended on this label may determine suitability for such uses by making trial application of Weedfree 63 at a recommended rate to a small number of plants. Prior to using Weedfree 63 on a larger number of plants, the treated plants should be

observed for signs of herbicidal injury during 30 to 60 days of normal growing conditions to determine if the treatment is non-injurious to the target plant species. **The user assumes responsibility for any plant damage or other liability resulting from the use of Weedfree 63 on plant species not recommended on this label.**

DOSAGE

For preemergence weed control apply 50 to 100 lb. (1.0 lb. to 2.0 lb active) of Weedfree 63 per broadcast acre.

TIMING AND METHOD OF APPLICATION

For optimum weed control, preemergence applications should be made immediately after transplanting seedlings or to weed-free container stock. Two applications may be necessary, in fall-transplanted conifer fields, for season-long weed control.

CONIFER - SPECIFIC USE RESTRICTIONS

In addition to the following, also observe GENERAL USE RESTRICTIONS listed at the end of this label.

- Do not apply more than 100 lb. (2.0 lb active) of this product per broadcast acre per year.
- **NOT FOR CONIFER RELEASE IN FOREST MANAGEMENT PROGRAMS OR FOR FOREST REGENERATION APPLICATIONS.**
- Do not apply Weedfree 63 to wet or dew covered foliage. Should this happen, irrigate immediately to completely remove herbicide particles from leaf surfaces.
- Do not apply Weedfree 63 to whorl-leaved plants as their leaves may channel the herbicide granules to the leaf base and may result in foliage injury.
- Do not apply Weedfree 63 in an enclosed greenhouse structure as injury to plant foliage may result.
- Do not store or transport treated container stock in an enclosed structure until completion of 4 irrigations (minimum 21 days) as injury to non-labeled plants may occur.
- Always apply Weedfree 63 only to healthy conifer stock. Do not apply Weedfree 63 to conifers that are under stress from excessive fertilizer or soil salts, disease, nematodes, frost, drought, flooding, previously applied pesticides, soil insects, or winter injury, as severe injury may result.
- Do not graze or feed livestock forage cut from areas treated with Weedfree 63.

SELECTED FIELD-GROWN DECIDUOUS TREES

Many field-grown deciduous trees are tolerant to applications of Weedfree 63 directed to the soil and base of the plant. Weedfree 63 will provide preemergence control of many broadleaf weeds and grasses.

DECIDUOUS TREE SPECIES	
Almond	<i>Prunus spp.</i>
Apple	<i>Malus X domestica</i>
Apricot	<i>Prunus spp.</i>
Ash, Green	<i>Fraxinus pennsylvanica</i>
Ash, White	<i>Fraxinus americana</i>
Birch, River	<i>Betula nigra</i>
Cherry	<i>Prunus spp.</i>
Chestnut	<i>Castanea spp.</i>
Crabapple	<i>Malus spp.</i>
Dogwood	<i>Cornus florida</i>
Eucalyptus	<i>Eucalyptus viminalis</i> <i>Eucalyptus pulverulenta</i> <i>Eucalyptus camaldulensis</i>
Filbert	<i>Corylus spp.</i>
Lilac	<i>Syringa vulgaris</i>
Locust, Black	<i>Robinia pseudoacacia</i>

10/12

† Maple, Black	<i>Acer nigrum</i>
† Maple, Red	<i>Acer rubrum</i>
† Maple, Sugar	<i>Acer saccharum</i>
Myrtle, Crepe	<i>Lagerstroemia indica</i>
Nectarine	<i>Prunus spp.</i>
Nut, Hickory	<i>Carya sp.</i>
Nut, Macadamia	<i>Macadamia ternifolia</i>
Oak, Chestnut	<i>Quercus prinus.</i>
Oak, Pin	<i>Quercus palustris</i>
Oak, Red	<i>Quercus. rubra</i>
Oak, Water	<i>Quercus nigra</i>
Oak, Willow	<i>Quercus phellos</i>
Olive, Russian	<i>Elaeagnus angustifolia</i>
Poplar	<i>Populus spp.</i>
Poplar, Tulip	<i>Liriodendron tulipifera</i>
Peach	<i>Prunus persica</i>
Pear	<i>Pyrus spp.</i>
Pecan	<i>Carya spp.</i>
Pistachio	<i>Pistacia vera</i>
Plum	<i>Prunus spp.</i>
Prune	<i>Prunus spp.</i>
Redbud	<i>Cercis canadensis</i>
Sweetgum	<i>Liquidamber styraciflua</i>
Sycamore	<i>Platanus occidentalis</i>
Walnut, Black	<i>Juglans nigra</i>

† Do not apply to maple trees used for production of maple sap or maple syrup.

DOSAGE

Apply 25 to 100 lb. (0.5 lb. to 2.0 lb active) of Weedfree 63 per acre to the soil area surrounding deciduous plants for preemergence weed control. This product may be applied as a single or split application.

DO NOT apply more than 100 lb. of product per acre per season.

For spot treatments, refer to the following table for dosage recommendations.

Pounds Active/Acre	Pounds Weedfree 63 /Acre	Pounds of Weedfree 63 to treat 400 Sq. Ft	Pounds of Weedfree 63 to treat 100 Sq. Ft
2	100	1.0	0.25

TIMING

Weedfree 63 can be applied after transplanting or to established deciduous trees. For optimum weed control, applications should be made prior to weed germination.

For maximum safety to deciduous species mentioned on this label, post-directed applications of Weedfree 63 should be made to the soil prior to bud swell in the spring or after trees have initiated dormancy in the fall.

Weedfree 63 may be phytotoxic to the foliage of non-target plants. Avoid making applications of this product under conditions that favor drift to non-target areas.

11/12

Note: Applications made after bud swell may result in injury to deciduous trees and are not recommended. If a non-dormant application is required due to weed competition, do not apply during periods of new foliage growth. Applications should be made after foliage has fully expanded and hardened off. Direct material toward the soil at the base of the trees. Splashing soil can carry Weedfree 63, which may injure the leaves of some deciduous trees.

METHOD OF APPLICATION

Weedfree 63 should be directed to the soil. Avoid direct contact with foliage, flowers or green bark. Provide uniform distribution and coverage to the soil surface. Application equipment should be calibrated carefully before each use.

FIELD-GROWN DECIDUOUS TREES SPECIFIC USE RESTRICTIONS

- DO NOT apply more than 100 lb. (2.0 lb active) of this product per broadcast acre per year.
- The use directions described here for field-grown deciduous trees do not apply for bearing treefruit, nut and vine crops.
- DO NOT apply Weedfree 63 to trees that have been weakened or are under stress from excessive fertilizer or soil salts, disease, nematodes, frost, wind injury, drought, flooding, previously applied pesticides, insects, or winter injury as severe injury may result.
- DO NOT graze or feed livestock forage cut from areas treated with Weedfree 63.

USE RESTRICTIONS THAT APPLY TO ALL REGISTERED APPLICATIONS ARE LISTED BELOW:

GENERAL USE RESTRICTIONS:

- Read and observe all label directions before using.
 - Do not contaminate irrigation water or water used for domestic purposes.
 - Do not use any plants treated with Weedfree 63 for feed or forage.
 - Do not feed or allow animals to graze on any areas treated with Weedfree 63.
 - Weedfree 63 should be applied only by ground application equipment.
 - Do not apply when weather conditions favor drift. Avoid drift to all non-target areas.
 - Use Weedfree 63 only for recommended purposes and at recommended rates.
 - Do not treat ditch banks or waterways with Weedfree 63.
- **ROTATION CROP RESTRICTIONS**
Do not plant any crop other than the tolerant plant species listed on this label within 10 months following a Weedfree 63 treatment.
 - Do not direct seed any crops, other than those on this label, within 60 days following a Weedfree 63 treatment.
 - Do not transplant seedling crops, other than Weedfree 63-labeled crops, within 30 days following a Weedfree 63 treatment

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

12/12

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer and Inherent Risks of Use above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.