



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

52287-27

Date of Issuance:

1/24/23

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

FERTILIZER W/ Ronstar 0.75 + Barricade
0.25

Name and Address of Registrant (include ZIP Code):

Harrell's, LLC
P.O. Box 807
Lakeland, FL 33802

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.


On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. The data requirements for storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) are not satisfied. You have 18 months from the date of registration to provide these data.
3. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 52287-27."

Signature of Approving Official:


Shaja B. Joyner, Product Manager 20
Fungicide-Herbicide Branch
Registration Division 7505P

Date:

1/24/23

4. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 06/01/2021
- Alternate CSF #1 dated 06/01/2021
- Alternate CSF #2 dated 06/01/2021

If you have any questions, please contact Ernest Kraka by phone at (202)-566-2811, or via email at kraka.ernest@epa.gov

Enclosure

Note to PM:

Bold, italicized text is for the reader and is not a part of the label.

EPA Reg. No. 52287-XX

[Bracketed information is optional] Text separated by / denotes and/or options

Restricted Use Pesticide due to potential human health risks of concern (cancer) from chronic dietary or dermal exposure when used in accordance with label directions. For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.



OXADIAZON	GROUP	14	HERBICIDE
PRODIAMINE	GROUP	3	HERBICIDE

FERTILIZER W/ Ronstar 0.75 + Barricade 0.25

[Alternate Brand Names: Fertilizer With Ronstar 0.75 + Barricade 0.25; Fertilizer W/ Oxadiazon 0.75 + Prodiamine 0.25; Harrell's PROtect Professional Fertilizer W/ Ronstar 0.75 + Barricade 0.25; Turf Fertilizer W/ Ronstar 0.75 + Barricade 0.25; Fertilizer W/ Ronstar 0.75 and Barricade 0.25; Fertilizer W/ Ronstar 0.75 & Barricade 0.25; Fertilizer W/ Oxadiazon 0.75 and Prodiamine 0.25; Fertilizer W/ Oxadiazon 0.75 & Prodiamine 0.25]

[Batch Number] [Lot Number] [SKU Number] [Fertilizer N-P-K] [Bulk] [Density] [xx lbs. per cu. ft]

For use in established turf in non-agricultural areas (e.g., golf courses, parks, athletic fields) and sod farms only.

Not for use on residential turfgrass

SELECTIVE PREEMERGENT HERBICIDE for the control of annual grasses and broadleaf weeds in turfgrass by a professional applicator.

ACTIVE INGREDIENT:

Oxadiazon:0.75%

Prodiamine:0.25%

OTHER INGREDIENTS:99.00%

TOTAL100.00%


Net Weight: 50 lb. (22.7 kg), 1,000 lb. (453.4 kg), 2,000 lb. (907.2 kg), or bulk trailers

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

READ THE ENTIRE LABEL BEFORE USING



EPA Reg. No. 52287-XX

EPA Est. No. 52287-FL-1

FIRST AID

If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	

For **MEDICAL** Emergencies Call 24 Hours a Day: 1-800-304-0824

For **TRANSPORTATION** Emergencies Call 24 Hours a Day 1-800-424-9300

For **PRODUCT USE** information Call 1-800-282-8007

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Harmful if inhaled. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Loaders, applicators and other handlers must wear coveralls worn over short-sleeved shirt and short pants, socks, chemical-resistant footwear, and waterproof gloves. Follow manufacturer’s instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For Terrestrial uses: **Do not** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. **Do not** contaminate water when disposing of

Note to PM:

Bold, italicized text is for the reader and is not a part of the label.

EPA Reg. No. 52287-XX

[Bracketed information is optional] Text separated by / denotes and/or options

equipment washwater or rinsate. Runoff may be hazardous to aquatic organisms in water adjacent to treated areas. This pesticide is toxic to oysters and shrimp.

GROUNDWATER ADVISORY: This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

SURFACE WATER ADVISORY:

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water.

This product is classified as having high potential for reaching surface water via runoff for several months or more after application.

A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of oxadiazon from runoff water and sediment.

NON-TARGET ORGANISM ADVISORY: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize drift.

DIRECTIONS FOR USE

RESTRICTED USE PESTICIDE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. **DO NOT** use on home lawns.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (sod farm, commercial nurseries). **Do not** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water, is coveralls, shoes plus socks and waterproof gloves.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The “WPS” applies when this product is used to produce agricultural plants on farms, forests or nurseries. Applicators and handlers should wear long-sleeved shirt, long pants, waterproof gloves, shoes with socks and protective eyewear. Thoroughly wash face and hands with soap and water after exposure and before eating or smoking. Launder exposed clothing before reuse. **Do not** allow reentry to treated areas until dusts have settled and the turfgrass or soil is dry.

WEED-RESISTANCE MANAGEMENT

For resistance management, Fertilizer with Ronstar 0.75 + Barricade 0.25 contains the active ingredients Oxadiazon, a Group 14 Herbicide, and Prodiamine, a Group 3 Herbicide, as determined by the mode of action (MOA) classification system of the Weed Science Society of America. Any weed population may contain or develop plants naturally resistant to Group 14 and Group 3 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies must be followed. Continued effectiveness of this product depends on the successful implementation of a weed resistance management program.

To delay herbicide resistance users must take one or more of the following steps:

- Rotate the use of this product and other Group 14 and Group 3 herbicides within a growing season sequence, or among growing seasons with different herbicide groups that control the same weeds in a field.
- Adopt an integrated weed-management program for herbicide use that includes:
 - Scouting and uses historical information related to herbicide use and crop rotation, and
 - Consideration of tillage (or other mechanical control methods), and
 - Cultural (e.g., higher crop seeding rates), and
 - Precision fertilizer application method and timing to favor the crop and not the weeds), and
 - Biological (weed-competitive crops or varieties) and other management practices.
- Scout fields prior to application to identify the weed species present and their growth stage to determine if the intended application will be effective.
- Scout fields after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include:
 - Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds, or
 - A spreading patch of non-controlled plants of a particular weed species or,
 - Surviving plants mixed with controlled individuals of the same species.
- If resistance is suspected, prevent weed seed production in the affected area by using an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use and switch to another management strategy or herbicide with a different mode of action, if available.

- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- Report any incidence of local non-performance of this product against a particular weed species to your [Harrell's] [company] representative at (800) 282-8007, [and/or at <https://www.harrells.com>].

BEST MANAGEMENT PRACTICES

Proactively develop and implement diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides. A diversified program may include the use of herbicides with different modes of action along with cultural, mechanical, and/or biological practices. Best management practices include the following:

- To the extent possible, use a diversified approach toward weed management. Whenever possible incorporate multiple weed-control practices such as mechanical cultivation, biological management practices, and crop rotation.
- Plant into weed-free fields and keep fields as weed-free as possible.
- Identify weeds present in the field through scouting and field history and understand their biology. The weed-control program should consider all the weeds present. Apply this herbicide at the correct timing and rate needed to control the most difficult weed in the field.
- Fields with difficult to control weeds should be rotated to crops that allow the use of herbicides with alternative modes of action (MOA) or different management practices.
- Difficult to control weeds may require sequential applications of herbicides with differing MOA's.
- Prevent field-to-field and within-field movement of weed seed or vegetative propagules. Thoroughly clean plant residues from equipment before leaving fields.
- To the extent possible **do not** allow weed escapes to produce seeds, roots or tubers. Manage weed seeds at harvest and post-harvest to prevent a buildup of the seed seedbank.
- If resistance is suspected, treat weed escapes using a herbicide with a different MOA than those in this product or use non-chemical methods to remove escapes.

DRIFT ADVISORY

WIND: Drift potential generally increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.**

PRODUCT INFORMATION

Fertilizer W/Ronstar 0.75 + Barricade 0.25 is a preemergent herbicide for control of many annual grasses and broadleaf weeds in turf. Fertilizer W/Ronstar 0.75 + Barricade 0.25 controls weeds by killing the weed seedlings as they come in contact with the herbicide during germination. Apply Fertilizer W/Ronstar 0.75 + Barricade 0.25 using a properly calibrated spreader for granular applications. Uniform application is essential for satisfactory control of listed weeds. Calibrate the spreader prior to use following the manufacturer's directions and check frequently during the application to be certain that the equipment is working properly. Water-in the product as soon as practical after application for best results. Failure to follow the directions for use and precautions on this label may result in poor weed control or crop injury.

After application, user must thoroughly irrigate the site as soon as possible on the same day of application.

TURF

For use in established Kentucky Bluegrass, Creeping Bentgrass, Bermudagrass, Buffalograss, Perennial Ryegrass, St. Augustinegrass, Seashore Paspalum, Tall Fescue, and Zoysiagrass turf on golf courses, parks, athletic fields and sod farms only. Rake turf area free of leaves, mowing if necessary, so granules reach soil surface. Measure area to be treated and uniformly apply required amount, using a fertilizer spreader. Cross-treating with one-half required amount in each direction will give best results. If rain is not expected shortly after application, irrigating turf thoroughly will increase effectiveness.

USE RESTRICTIONS:

- For use in established turf non-agricultural areas (e.g., golf courses, parks, athletic fields) and sod farms only.
- Not for use in lawns, or residential turf.
- Not for use by homeowners/consumers.
- **Do not** use on turfgrass on Residential Properties.
- **Do not** graze or feed livestock forage cut from areas treated with this product.
- **Do not** apply aerially.
- **Do not** cut (harvest) treated sod before 120 days after application. **Do not** apply to newly set sod until the following year.
- **DO NOT** APPLY this product to golf course fairway turf dichondra, colonial bentgrass, velvet bentgrass, annual bluegrass (*Poa annua*), or centipedegrass are desirable species.
- **Do not** apply more than 400 lbs. of product (Oxadiazon: 3 lbs. a.i.; Prodiamine: 1.0 lb. a.i.) per acre per application.
- For St. Augustinegrass, **do not** exceed the per/application rate of 400 pounds per acre (equivalent to 3 lbs. a.i. of Oxadiazon and 1 lb. a.i. of Prodiamine).
- The maximum annual application rate allowed is 600 pounds of product per acre per year (equivalent to 4.5 lbs. a.i. of Oxadiazon and 1.5 lbs. a.i. of Prodiamine).
- **Do not** exceed the rates indicated in the Maximum Annual Rates Table.
- **Do not** make more than 2 applications per year.
- For Creeping Bentgrass turf, **do not** exceed the rate of 264 pounds per acre (equivalent to 2 lbs. a.i. of Oxadiazon and 0.66 lb. a.i. of Prodiamine).
- **RETREATMENT INTERVAL:** Subsequent applications can be made 60 days after the previous treatment.
- **Do not** apply this product through any type of irrigation system.
- **Do not** apply to Creeping Bentgrass turf mowed at less than 1/2 inch.
- **Do not** apply to newly seeded areas.
- **Do not** overseed fairways before 120 days after application of this product.
- **Do not** use where *Poa trivialis* is present in overseed mix.
- **Do not** use on newly sprigged or plugged bermudagrass.

For use on golf courses, parks, athletic fields and sod farms:

- A maximum single application rate of 400 pounds of product (3.0 lbs. a.i. of oxadiazon and 1.0 lb. a.i. of prodiamine)/A, and a maximum annual application rate of 600 pounds of product (4.5 lbs. a.i. of oxadiazon and 1.5 lbs. a.i. of prodiamine)/A.

- For applications on non-agricultural turf (e.g., golf courses, parks, athletic fields) that follow a liquid application, Fertilizer W/Ronstar 0.75 + Barricade 0.25 must be applied 60 days or later after the liquid application. After applying this product, no liquid applications of oxadiazon are allowed within the same calendar year on the same area of turf.
- For applications to non-agricultural turf (e.g., golf courses, parks, athletic fields), this product may not be applied prior to any liquid applications of oxadiazon within the same calendar year on the same area of turf.
- On golf course turf only: In areas of heavy weed infestation only, the maximum single application rate is 533 pounds of product (4.0 lbs. a.i. of oxadiazon and 1.08 lbs. a.i. of prodiamine) per acre, and the maximum annual application rate is 600 pounds of product (4.5 lbs. a.i. of oxadiazon and 1.5 lbs. a.i. of prodiamine) per acre. No liquid applications may be applied on areas of turf treated at this maximum application rate in the same year.
- On golf course turf only: total applications of oxadiazon of all physical forms are limited to 30% of the total managed turf area of a golf course per year. Applications may be made to any turf surface type, including fairways, tees, greens, roughs, non-playable turf surfaces, or other managed turf areas. This 30% limit may be applied to the same or different turf areas from year to year.
- On golf course turf and sod farms: a minimum 10-foot-wide untreated vegetative buffer must be maintained between the treated area and any surface water features.
- The minimum retreatment interval (RTI) between applications is 60 days
- Total applications of oxadiazon of all physical forms must not exceed 6.0 lbs. a.i./A/year if this product is used on the same site with liquid formulations.

(Note to PM: Optional State Nitrogen and Phosphorus Fertilizer Use Restrictions to be inserted as required. Example statements can be found under the Optional State Fertilizer Statements heading.)

USE PRECAUTIONS:

- Fertilizer W/Ronstar 0.75 + Barricade 0.25 may cause discoloration of Creeping Bentgrass, Bermudagrass and St. Augustine grass. This discoloration is temporary and normally outgrown within 2 to 3 weeks.
- Avoid contact with flowers and shrubs except as stated elsewhere on the label.
- For late, fall applications, seeding into treated areas MUST be delayed until 4 months after green up.

APPLICATION RATE:

Fertilizer W/Ronstar 0.75 + Barricade 0.25 may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. The length of time of residual pre-emergence weed control provided by this product is related to the rate applied. This product will not control established weeds.

In many areas a single application will provide control of listed weeds. However, under heavy weed pressure and/or an extended growing season, most effective weed control may be obtained

Note to PM:

Bold, italicized text is for the reader and is not a part of the label.

EPA Reg. No. 52287-XX

[Bracketed information is optional] Text separated by / denotes and/or options

by making an initial application followed a second application 60 days after the first application at rates that **do not** exceed those given in the Maximum Annual Rates Section.

Apply 264 to 400 lbs. of Fertilizer W/Ronstar 0.75 + Barricade 0.25 (equivalent to 2 – 3 lbs. a.i. of Oxadiazon and 0.66 – 1 lb. a.i. of Prodiamine)/per application per acre.

WEEDS CONTROLLED

When used in accordance with this label Fertilizer W/Ronstar 0.75 + Barricade 0.25 will provide control of the following weeds:		
Barnyardgrass	Henbit	Pusley, Florida
Bluegrass, Annual (<i>Poa Annua spp.</i>)*(1)	Itchgrass	Rescuegrass
Carpetgrass	Johnsongrass (from seed)	Shepherd’s Purse
Carpetweed	Junglerice	Sedge, Annual
Chickweed, Common	Knotweed	Signalgrass, Broadleaf
Chickweed, Mouseear (from seed)	Kochia	Speedwell, Persian
Crabgrass (Large, smooth)	Lambsquarter, Common	Sprangletop
Crowfootgrass	Lovegrass	Spurge, Prostrate
Cupgrass, Woolly	Oxalis	Stinging Nettle
Field Sandbur	Panicum (Texas, Fall, Browntop)	Virginia Buttonweed
Foxtails, Annual	Pigweed	Witchgrass
Goosegrass	Purslane, Common	Woodsorrel, Yellow (from seed)

¹Use not permitted in California

***NOTE:** In many areas, annual bluegrass exists as at least two biotypes or subspecies of *Poa annua*. The true annual biotype is *Poa annua spp. annua* which germinates every year from seed, while the biotype that behaves like a perennial is *Poa annua spp. reptans*. This perennial biotype can survive through the summer and established plants will not be controlled by this product. Consult the extension service or University weed specialists in your area for more information.

MAXIMUM ANNUAL RATES

Fertilizer W/Ronstar 0.75 + Barricade 0.25 is for use on the turf grass species listed below. The maximum annual rate for Prodiamine is lower than the maximum rate for Oxadiazon. **Do not** exceed the maximum annual rate for Prodiamine as given in the table below.

MAXIMUM ANNUAL RATE BY TURF GRASS SPECIES (1)		
Turf Species	Lbs. of product/A	Lbs. a.i. of Prodiamine/A
Creeping Bentgrass	264 lbs.	0.66 lb.
Buffalograss	400 lbs.	1.00 lb.
Kentucky Bluegrass	400 lbs.	1.00 lb.
Perennial Ryegrass	400 lbs.	1.00 lb.
Bermudagrass	600 lbs.	1.50 lbs.
Seashore Paspalum	600 lbs.	1.50 lbs.
St. Augustinegrass	600 lbs.	1.50 lbs.
Tall Fescue (including turf type)	600 lbs.	1.50 lbs.
Zoysia	600 lbs.	1.50 lbs.

(1) These are the maximum rates per calendar year by species limitations.

Note to PM:

Bold, italicized text is for the reader and is not a part of the label.

EPA Reg. No. 52287-XX

[Bracketed information is optional] Text separated by / denotes and/or options

(Note: Spreader Section is optional)

[SPREADER SETTINGS] / [COVERAGE INFORMATION]

[For 200 lb/A Rate: 50 lb will cover up to 10,870 sq. ft.]

		Application Rates		
		[4.6 lb/1,000 sq. ft. = 200 lb/A Covers: 10,870 sq. ft.]	[6.9 lb/1,000 sq. ft. = 300 lb/A Covers: 7,246 sq. ft.]	[9.2 lb/1,000 sq. ft. = 400 lb/A Covers: 5,435 sq. ft.]
[Spreader Name]	[Amount of Product]	[Spreader Settings]		
[Spyker 288 Super]	[XX]	[XX]	[XX]	[XX]
[Lesco Rotary (Model 021820)]				
[Scott's ProTurf R8A-1]				
[Lesco 36" Drop Spreader]				
[Lely HR Three Point Hitch Single Disc Broadcaster]				
[Vicon Super Flow PS 203]				

("XX" will be replaced with appropriate value on final label if table is used.)

STORAGE AND DISPOSAL

PESTICIDE STORAGE

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING

For 50 lb., 1000 lb., 2,000 lb. bags and totes:

Non-refillable container. **Do not** reuse or refill this container. Completely empty container into application equipment. Then dispose of empty container in a sanitary land fill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

For Killebrew and other bulk transport vehicles:

Refillable container: Container handling instructions are not required. Cleaning before refilling is the responsibility of the refiller.

(End Directions for Use)

(Fertilizer Information)

Note to PM:

Bold, italicized text is for the reader and is not a part of the label.

EPA Reg. No. 52287-XX

[Bracketed information is optional] Text separated by / denotes and/or options

[Fertilizer W/ Ronstar 0.75 + Barricade 0.25]	
Guaranteed [Fertilizer] Analysis	[N-P-K]
Total Nitrogen (N).....	XX%
XX % Ammoniacal Nitrogen	
XX % Urea Nitrogen*	
Available Phosphate(P ₂ O ₅).....	XX%
Soluble Potash (K ₂ O).....	XX%
Derived from: Polymer-Coated Urea, Urea, Ammonium Phosphate, and Muriate of Potash.	
*X% Slowly Available/Slow Release/Controlled release nitrogen from polymer-coated urea.	
[Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.html]	
F-352 (<i>Florida Fertilizer License Number</i>)	
[Fertilizer] [Density] [Bulk Density]	
[Any other state specific fertilizer required language]	

Note to PM : *Appropriate fertilizer information will be inserted on Final Printed Label to replace "X".*

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Harrell’s, LLC. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, HARRELL’S, LLC MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Harrell’s, LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, HARRELL’S, LLC DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

Note to PM:

Bold, italicized text is for the reader and is not a part of the label.

EPA Reg. No. 52287-XX

[Bracketed information is optional] Text separated by / denotes and/or options

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT HARRELL'S, LLC'S ELECTION, THE REPLACEMENT OF PRODUCT.

NET WEIGHT: 50 lbs., 1,000 lbs., 2,000 lbs.; or BULK TRAILER

Manufactured By/Manufactured For/Sold By/ and Guaranteed By:

Harrell's, LLC

PO Box 807

Lakeland, FL 33802

RONSTAR® is a registered trademark of the Bayer

BARRICADE® is a registered trademark of a Syngenta Company

Growing a Better World® is a registered trademark of Harrell's, LLC

Optional State Fertilizer Restriction Statements

(Note to PM: Fertilizer composition of products distributed in states with nitrogen and phosphorus fertilizer application restrictions will be formulated to contain percentages of nutrients that meet state fertilizer restrictions when applied according to the application rates for the active ingredient as specified in the Directions for Use.)

Appropriate statement to be inserted in the Application Restrictions Section of Label as necessary for State specific fertilizer restriction language regarding application of nitrogen and phosphorus. Some example statements are, but not limited to:

[“In MD, NH, NJ, VA: Application rates of soluble nitrogen are not to exceed 0.7 lb. soluble N/1000 sq.ft/application.”]

[And/Or]: [In NH, NJ, VA: Application of a product containing phosphorus to turf is limited to establishing turf in the first growing season.”]

[And/Or]:“Florida specific requirements for fertilizer use. For applications to Golf Course Fairways: **DO NOT** exceed fertilizer rates specified in the document titled SL 191, “Recommendations for N,P,K and Mg for Golf Course and Athletic Field Fertilization based on Mehlich III Extractant.”

[And/Or: Other state specific fertilizer language that may be required.]

[For Fertilizer Mixtures Containing Nitrogen]: [It is a violation of law in [Maryland][New Jersey][New Hampshire][list state(s)] to apply this product to turf at a rate that results in greater than 0.7 lb per 1,000 sq.ft. of readily available nitrogen, or more than 0.9 lb per 1,000 sq.ft. of total nitrogen, with a least 20% of which shall consist of slow release nitrogen, per application. Additional restrictions apply to total nitrogen applied to turf per season.]

Note to PM:

Bold, italicized text is for the reader and is not a part of the label.

EPA Reg. No. 52287-XX

[Bracketed information is optional] Text separated by / denotes and/or options

[Products sold in Florida: The following statement shall be placed in the Directions for Use section (see that section above in label): [Apply only to actively growing turf. **DO NOT** apply near water, storm drains, or drainage ditches. **DO NOT** apply if heavy rain is expected. Apply this product only to your turf, and sweep any product that lands in the driveway, sidewalk, or street, back onto your turf. Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices. Check with your county or city government to determine if there are local regulations for fertilizer use.]

[Virginia applications: [Apply no more than 0.7 lb per 1,000 sq. ft. of readily available nitrogen during any given 30-day period. Apply no more than 0.9 lb (cool season grasses) or 1.0 lb (warm season grasses) per 1,000 sq. ft. of total nitrogen, with at least 20% of which shall consist of slow release nitrogen, during any given 30-day period.]

[For Fertilizer Mixtures Containing Phosphorus]:

[For starter turf in MD]: [In Maryland, a soil test is not required for a new turf. Apply either 2 lbs. of phosphorus per 1,000 sq. ft. initially or apply 1 lb. of phosphorus per 1,000 sq. ft. twice during the starter year. On established turf, this product must be used in accordance with a soil test that was conducted by a laboratory identified under § 8-803.7 of the Agriculture Article, Annotated Code of Maryland, performed no more than 3 years before the application.] [Other state and local restrictions may apply]

[In Maryland] [For levels of phosphorus above 5%]: [This fertilizer contains phosphorus and may not be used on turf in the state of Maryland except when (1) providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory identified under §8-803.7 of the Agriculture Article, Annotated Code of Maryland, performed no more than 3 years before the application; (2) establishing vegetation for the first time, including after land disturbance, provided the application is conducted in accordance with the specified application rates established by the State; or (3) reestablishing or repairing a turf area. Not for use on established turf grass in Maryland.]

[For all %'s of phosphorus-containing fertilizers in VA]:

[Virginia Applications]: [This turf fertilizer contains phosphorous and is only for nonagricultural use on (i) turf during its first growing season, (ii) on turf areas being repaired or renovated, and (iii) on turf where a soil test performed within the last 3 years indicates a phosphorous deficiency. This fertilizer is not for the routine maintenance of turf.]

[Virginia & Maryland]: [When applied as directed, [we]/[Harrell's] guarantees the [controlled release]/[slow release]/[polymer-coated]/[Enhanced Efficiency] nitrogen fertilizer in this product will release at a rate not to exceed 0.7 lb. N/1,000 sq. ft./month.]

[Maryland Applications] [**DO NOT** apply near water storm drains or drainage ditches. **DO NOT** apply if heavy rain is expected. Apply this product only to your turf, and sweep any product that lands on the driveway, sidewalk, or street back onto your [turf].]

OPTIONAL MARKETING CLAIMS AND LOGO'S

(Note: The following logos and/or claims may or may not be used on the FPL)

Company Specific Information

Note to PM:

Bold, italicized text is for the reader and is not a part of the label.

[Bracketed information is optional] Text separated by / denotes and/or options

EPA Reg. No. 52287-XX

(Harrell's Corporate Logo)



Growing a Better World® is a registered trademark of Harrell's, LLC

Harrell's PROFertilizer® is a registered trademark of Harrell's, LLC

(Harrell's "H" Logo)



Contact us/ For more information

[call] 800.282.8007

[or] [visit us at] www.harrells.com

[or] use the QR code to visit Harrell's website

[Insert] QR Code Image (***Note to PM: If QR code is used, the link will be to the company website home page***)

Harrell's label identifier format: [EPA Reg No.Product Name.label revision date (yyyymmdd)]

FERTILIZER CLAIMS

Contains/Formulated with/ Harrell's PROFertilizer®

Fertilizer with Ronstar 0.75 + Barricade 0.25 is part of the Harrell's PROFertilizer® portfolio/line/family of products

Contains POLYON® Slow release/controlled release/slowly available nitrogen/phosphate/potassium/fertilizer

POLYON® Turf Fertilizer

POLYON® Polymer Coated Fertilizer

POLYON® [ST]/[short term] [Slow]/[Controlled]- release fertilizer

POLYON® [ST] – Distributed exclusively by Harrell's, LLC

POLYON® [ST]/[short term] [Slow]/[Controlled]- release fertilizer

POLYON® [ST] – Distributed exclusively by Harrell's

[Powered by] POLYON® [Slow]/[Controlled]-Release Fertilizer

POLYON® is a registered trademark of Harrell's Coatings Solutions, LLC

POLYON® feeds/releases for up to X weeks/months at X° F

Note to PM:

Bold, italicized text is for the reader and is not a part of the label.

EPA Reg. No. 52287-XX

[Bracketed information is optional] Text separated by / denotes and/or options

(note to PM: release time and temperature is based on the coating thickness of the polymer coated fertilizer used in the blend and inserted for the appropriate product – this information is being required by fertilizer control officials in some states)