

BEST AVAILABLE COPY

Microbiocide 15

ACCEPTED
with COMMENTS
in EPA Letter Dated:

JUL 31 1985

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

52377-2

ACTIVE INGREDIENT:

Poly(ethylene glycol) dimethylsiloxane
[Siloxane] 15.0 %

INERT INGREDIENT 85.0 %

This product contains 1.29 lb. of active ingredient per gallon and weighs
8.58 lb. per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing vapors. Avoid contact with skin, eyes, or clothing.

FIRST AID: If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes, streams, or ponds. Permits may be required for discharges containing this pesticide into lakes, streams, ponds, or public water. For guidance, contact the regional office of the Environmental Protection Agency.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Microbiocide 15 is used to control bacteria and fungi in recirculating commercial and industrial water systems. In its use, systems must be cleaned to remove slimes, and other deposits. An initial slug addition of fluid ounces of Microbiocide 15

of water to provide a concentration of 32 to 64 Microbiocide 15 based on the volume of water in the system, is recommended. Repeat initial dosage

Subsequent slug additions of 1.0 to 2.0 Microbiocide 15 per 1000 gallons

80 parts per million of Microbiocide 15 employed every 2 to 5 days, or as needed. The frequency of application depends upon the relative amount of biofilm and the severity of the problem. Slug additions should be made in the pump of water.

Microbiocide 15 is used to clean air-washing systems that maintain effective mist output. In its use, systems should be cleaned to remove deposits. An initial slug dose of 14.8 to 29.6 Microbiocide 15 per 1000

gallons of water should be employed. Repeat initial dosage until control is evident. Slug additions of 10.1 to 24.7 fluid ounces of Microbiocide 15 per 1000 gallons of water should be employed. The frequency of addition depends upon the relative amount of the bacterial problem. Slug additions may be made to collection trays of the airwash system.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Do not store more than four drums high. Leaking or damaged drums should be disposed of. Drums should be cleaned in a well-ventilated area and stored upright when not in use.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product should be disposed of by approved waste disposal facility.

CONTAINER DISPOSAL:

PLASTIC: Triple rinse for equivalent. Then offer for recycling or other secondary disposal, by incineration, or, if allowed by state and local authorities.

METAL: Triple rinse for equivalent. Then offer for recycling or other secondary disposal, by incineration, or by other procedures approved by state and local authorities.

Manufactured by:

Blue Valley Laboratories of Tennessee

P.O. Box 11984

Memphis, TN 38111

EPA REG. NO.

EPA EST.

NET CONTENTS