

Reg # 52466-8

PM-19

194

PRODUCT: TPC SEVIN 5% DUST

EPA Reg. No.: 52466-8

PRODUCT No:


MIX No.:

DATE: 01/25/93

ITEM:

COMMENTS: Draft Label For Submission To EPA In Response To EPA Letter Dated December 16, 1992.

Draft Label
Front Panel



(Logo)

TPC SEVIN* 5% DUST

(Insecticide)

A Ready To Use Multi-Purpose Dust

For use on Fruits, Vegetables, Poultry, Dogs, Cats Premises & Litter
For Control of Northern Fowl Mites and Lice on Poultry
For Control of Ticks and Fleas on Dogs and Cats
A Premise Treatment For Chicken Mites and Fleas
For Control of Northern Fowl Mites, Chicken Mites,
Fleas and Bed Bugs in Litter

Active Ingredient:

Carbaryl (1-Naphthyl N-Methylcarbamate) 5.00%

Inert Ingredients: 95.00%
100.00%

* Trademark of Rhone Poulenc for the active ingredient
(1-Naphthyl N-Methylcarbamate)

Keep Out Of The Reach Of Children

CAUTION

See Side Panel For Additional
Precautionary Statements

NET CONTENTS: 1 Pound (454 g)

EPA Reg. No.: 52466-8


EPA Est. No.: 6175-LA-1

Horse Health Products, Inc.
421 East Hawley Street
Mundelein, IL 60060 USA

(lot number statements as needed)

Mfg. & Printed in U.S.A. (label revision number) (product number)

Page 1 of 4
January 1992
A:JDHHP008.DOC-1



274

Draft Label
Side/Back Panel

TPC SEVIN 5% DUST

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed. Harmful if inhaled. Avoid breathing dust. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Remove contaminated clothing and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling.

Do not use on kittens or puppies less than 4 weeks of age. Do not use on pregnant dogs. Do not use this product in commercial food areas of food handling establishments, restaurants, or other places where food is prepared or processed. During application avoid drift to adjacent food and forage crops that may be damaged or rendered unfit for sale, use or consumption. Keep out of the reach of children and animals.

Note to Physician: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.


IF IN EYES: Flush Eyes with plenty of water. Call a physician if irritation persists.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff or drift from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment of residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Additional protective information may be obtained from your Cooperative Extension Service.



394

Draft Label
Side/Back Panel

=====

TPC SEVIN 5% DUST

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Apply TPC Sevin 5% Dust when insects or their damage appear. Make applications in the morning or evening when it is cool. Treat upper and lower surfaces of foliage for complete coverage. Repeat as necessary, generally at 7 day intervals. Do not apply in excess.

FOR USE ON BEANS TO CONTROL:

MEXICAN BEAN BEETLE - Use at a rate of 1/4 lb. TPC Sevin 5% Dust per 1,000 sq. ft.

LEAFHOPPERS, BEAN LEAF BEETLE, CUCUMBER BEETLES, WESTERN BEAN CUTWORM, FLEA BEETLES, JAPANESE BEETLES - 1/2 lb. per 1000 sq. ft.

CORN EARWORM, ARMYWORMS, VELVET BEAN CATERPILLAR - 1/2 to 3/4 lb. per 1,000 sq. ft.

LIMA BEAN POT BORER, LYGUS and STINK BUGS - 1 lb. per 1,000 sq. ft.

May be used up to harvest. Apply thoroughly at first signs of insect infestation. When necessary repeat every 7 to 10 days.

FOR USE ON CUCUMBER, SUMMER SQUASH, PICKLES AND MELONS TO CONTROL:

CUCUMBER BEETLES, SQUASH BUGS, and FLEA BEETLES - 1/2 lb. per 1,000 sq. ft.

PICKLEWORM, MELONWORM - 1/4 TO 1/2 lb. per 1,000 sq. ft.


May be used up to harvest.

FOR USE ON GRAPES TO CONTROL:

GRAPE LEAF FOLDER - 1st brood: 1/2 lb. per 1,000 sq. ft. just before first brood larvae emerge from first rolls, 2nd and 3rd Broods - 1/2 to 1 lb. per 1,000 sq. ft.

GRAPE LEAF HOPPER - 1/2 lb. per 1,000 sq. ft.

May be applied up to 7-day preharvest.



434

Draft Label
Side/Back Panel

=====

TPC SEVIN 5 $\frac{1}{2}$ DUST
DIRECTIONS FOR USE (CONT'D)

FOR USE ON LETTUCE, MUSTARD GREENS, SPINACH, AND TURNIPS TO CONTROL:

FLEA BEETLES, HARLEQUIN BUG, LEAPHOPPER, - 3/4 to 1 lb per 1,000 sq. ft.

IMPORTANT: DO NOT APPLY TO HEAD LETTUCE WITHIN 3 DAYS OF HARVEST. DO NOT APPLY TO LEAF LETTUCE OR SPINACH, TURNIP OR MUSTARD GREENS WITHIN 14 DAYS OF HARVEST.

FOR USE ON DOGS AND CATS TO CONTROL:

TICKS AND FLEAS - Dust powder liberally over the entire body. Rub dust into hair and around feet, pads and between the toes so that they are well treated. Take care not to get the powder in the animals eyes, nose or mouth. Dust bedding, kennel and sleeping area regularly. Apply at least once weekly for best results. Do not use on Pregnant Dogs. Avoid treatment of kittens or puppies less than 4 weeks of age.

FOR USE ON CHICKENS, TURKEYS, DUCKS, GEESE, AND GAME BIRDS FOR CONTROL OF:

NORTHERN FOWL MITE, LICE and as a supplement to PREMISE treatment for CHICKEN MITES, and FLEAS - Dust birds at a rate of 1 lb. per 100 birds. As a litter treatment to control northern fowl mite, chicken mite, lice, fleas, and bed bugs, treat litter at a rate of 1 lb. per 40 sq. ft.

CAUTION: Do not repeat applications within 4 weeks. Do not apply to birds within 7 days of slaughter or do not house birds in treated houses within 7 days of slaughter. Do not apply directly to eggs or litter. Do not contaminate drinking water.

STORAGE AND DISPOSAL

Storage: Store in a cool, dry place.

Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

Container Disposal: Do not reuse Bottle. Rinse thoroughly before discarding in trash

NOTICE OF WARRANTY

HORSE HEALTH PRODUCTS, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR OTHERWISE, EXPRESSED OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THE PRODUCT UNDER NORMAL CONDITIONS IN ACCORD WITH THE STATEMENTS MADE ON THE LABEL.

