Mefr Kempter Product Manager (32) Disinfectants Franch

Enclosures

EPA Form 8570-6 (Rev. 5-76)

ATTACHMENT IS APPLICABLE
SIGNATURE OF APPROVING OFFICIAL

91949:I: Pringle:K-12: KENCO: 1(1/16/87: 10/27/87: aw: vo:dg: jh

Registration Division (TS-767C)

BEST AVAILABLE COPY

PREVIOUS EDITION MAY BE USED UNTIL SUPPLY IS EXHAUSTED.

RM-40 MICROBIOCIDE

ACCEPTED with COMMENTS in EPA Letter Dated

10121 1987

Under the Federal Insected Eungaste, and Rodenticute Act as neverded, for the registered under EPA

ACTIVE INGREDIENT:

Poly(oxyethylene (dimethyliminio) ethylene-

'.: '.:

(dimethyliminio) ethylene dichloride)

INERT INGREDIENT

BEST AVAILABLE COPY

This product contains 1.74 b. of active ingredient per gallon and weighs <u>8.59</u> lb, per gallon,

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful it swallowed. Avoid breathing vapors. Avoid contact with

skir, eyes, or clothing. FIRST AID: If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, cr. if these are not available, drink large quantities of water.

Avoid alcohol. Call a physiciari immediately. ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes, streams, or ponds. Permits may be required for discharges containing this pessicide into lakes, streams, ponds, of public water. For guidance, contact the regional office of เมื่อ ็ะเพ่าอาเทคิกิเลโะก็อเอเนือก Agency.

DIRECTIONS FOR

It is a violation of Federal law to use this proc its labeling.

RM-40 MICRORIOCIDE and fungi in recirculating commercial and indu

to its use, systems must be cleaned to remo slime, and other deposits. An initial slug addit fluid ounces of <u>RM-40 MICROBIOCIDE</u> of water to provide a concentration of ___24_ RM-40 MICROBIOCIDE

in the system, is recommended. Repeat initial Subsequent slug additions of ____0.73

RM-40 MICROBIOCIDE per 1000 parts per million of <u>RM-40 MICRO</u>

employed every 2 to 5 days, or as needed. The upon the relative amount of bleedoff and the se lēm. Slug additions should be made in the sur RM-40 MICROBIOCIDE is use air-washing systems that maintain effective m

to its use, systems should be cleaned to re deposits. An initial slug dose of ____11.1___ RM-40 MICROBIOCIDE mended. Repeat initial dosage until control is

tions of 7.5 to 18.3 fluid ounces of _ per 1000 gallons of water should be employed e frequency of addition depends upon the relativ of the bacterial problem. Slug additions may be

collection trays of the airwash system.

STORAGE AND D

Do not contaminate water, food, or feed by storage or disposal, STORAGE: Do not stack more than four drums high. Leaking drums for disposal. Spit's should be absorbed in sawdust or sand tainer closed when not in use,

PESTICIDE DISPOSAL: Wastes resulting from the use of this proved waste disposal facility

CONTAINER DISPOSAL: [] PLASTIC. Triple rinse for equivalent). Then offer for recycling sanitary fandfill, by incineration, or, if alloyed by state and lo

[] METAL Tuple rinse for equivalent). Then offer for recycling i sanitary fandful, or by other procedures approved by state an

	Ma	nuf	act	n Lec
CTI	MER.	D T N	10	

ROMAC ENGI 25669 BELLERIVE RD VALENCIA, CA 91355 EPA REG. NO. ___ EP

NET CONTENTS __