US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (TS-767) WASHINGTON, DC 20460

52951-2 TERM OF ISSUANCE

EPA REGISTRATION NO.

TE OF ISSUANCE

NOTICE OF PESTICIDE: REGISTRATION

AEREGISTRATION

(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)

NAME OF PESTICIDE PRODUCT

Foura-Lemon

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

1

Foural Enterprises 204 Hillcrest Avenue Somerset, NJ 08873

J

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered

This product is conditionally registered in accordance with FIFPA section 3(c)(7)(A) provided that you:

- 1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
- 2. Make the labeling changes listed below before you release the product for shipment:
 - Add the phrase "IPA Pegistration No. 52951-2."
 - Include the subheadings:
 - Pesticide Disposal
 - Container Disposal

in the appropriate places in the Storage and Disposal section.

Place the referral statement "See side panel for additional precautionary statements" to be placed directly below the statement "Call a physician immediately."

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE

EPA Form 8570-6 (Rev. 5-76)

PREVIOUS EDITION MAY BE USED UNTIL SUPPLY IS EXHAUSTED.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

John H. Lee Product Manager (31) Disinfectants Branch

Registration Division (TS-767C)

Enclosures



THIS DISINFACTANT reduces the surface tension of water, mproving its wettability. Microbes are capable of mass reproduction in sub-visible pores, scratches and cracks of surfaces. The excellent wettability of THIS PRODUCT improves the penetration of the use solution into areas that otherwise might not be reached.

THIS DISINFECTANT is effective against a wide variety of gram-positive and gram-negative bacteria. It is also an excellent deodorant and is especially effective in controlling soors that are bacterial in origin. Has a pleasant lemon xdor.

THIS DISINFECTANT mixes with water to form clear, stable use solutions. Use solutions are non-corrosive and non-staining to plastic, vinyl, synthetics, enamel, tile and most common metals.

PRECAUTIONARY STATEMENTS **HAZARDS TO HUMANS** AND DOMESTIC ANIMALS CAUTION

Do not get in eyes, on skin or on clothing. Harmful If swallowed. Avoid contamination of food. Avoid breathing spray mist.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not discharge into lakes, streams, ponds, or public water unless in accordance with an NPDES Permit. Do not contaminate water by cleaning of equipment or by disposing of wastes.

STORAGE AND DISPOSAL

Store only in tightly closed, original container in a secure area water resulting from the use of this product may be disposed of on site or at an approved waste disposed facility.

Do not read amply container. Triple rinse empty container with water. Return metal drums to reconditioner or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities. Plastic containers may be disposed of in a sanitary landfill, incinerated or, if allowed by local authorities, by burning. If burned, stay out of amake.

Directions for use

t is a violation of Federal law to use this product in a nanner inconsistent with its labeling,

APPLICATION INFORMATION

Select the proper dilution of THIS PRODUCT for your use from the directions below. Apply the use solution with a clean mop, sponge, cloth, squeegee, mechanical scrubber or mechanical spraying device.

BEST AVAILABLE COPY

FOURA-Lemon

LEMON ODOR-DISINFECTANT-DEODORANT-CLEANER

*Kills Pseudomonas, Salmonella and Staphylococcus. • • Virucidal

ACTIVE INGREDIENT:

Alkyl (C₁₄, 58%; C₁₆, 28%; C₁₂, 14%) dimethyl benzyl ammonium chloride 2.00% INERT INGREDIENTS:

This disinfectant is effective at 5 oz./gal. against Pseudomonas aeruginosa PRD-10, Salmonella choleraesuis ATCC 10708, and Staphylococcus aureus ATCC 6538 when tested by the A.O.A.C. Use Dilution Test Method for precleaned, hard non-porous, inanimate environmental surfaces.

**This disinfectant is also an effective virucide at 5 oz./gal, for influenza virus, Type A, Brazil on inanimate environmental surfaces.

This product may be used for cleaning and deodorizing, onestep cleaning, and disinfection, or disinfectant of precleaned, non-porous, Inanimate hard surfaces.

KEEP OUT OF REACH OF CHILDREN

CALITICAL

See side panel for additional precautionary statements.

FOR INDUSTRIAL, COMMERCIAL OR MEDICAL USE ONLY

Statement of Prectical Treatment. In case of contact, immediately flush eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, raw egg whites, getatin solution or, it these or not readily available, drink a large quantity of water. Avoid alcohol.

Call a physician immediately. ___ EPA Est. No. 52951 - NJ-01THIS PRODUCT EPA Reg. No.

Manufactured by:

NET CONTENTS:

FOURAL Enterprises

204 Hillcrest Ave.

I U.S. Gal.

Somerset, NJ 08873

This product may be hard, non-pocqus, (glass, metal, glaze leather, vinyl, enan walls, floors, applian

Prepare fresh use so Discard solutions wh disinfection, thoroug Surfaces mûst rema

assure proper gerinic

To Clean and De Disinfection. Depen cleaned, use 12 to 16 water.

water) for one-step with light to mediur preclean as describe PRODUCT at 5 oz. p Industrial areas, Ba

Hospitals, Veterinary

of this product per

Food and Meat Pro areas), Other non-me disinfectant with a : 1-1/3 gallons of was natively, clean and of THIS PRODUCT parts water).

For deodorizing gar waste receptacies, a especially importan form properly. The oz. of THIS PRODU

parts water),

For mildew contro disinfection. Repea new growth appears

THIS PRODUCT m scrubbing equipment

12 inches use 4 ounces of BUTYRAC 200 with 3 quarts of Alanap-L.

If applied according to directions BUTYRAC® 200 Herbicide with Alanap-L will control or suppress growth of above weeds, reducing their competition with the soybean crop.

NOTE: Applications of this tank mix should not be made when weather conditions favor drift. Do not apply this tank mix if rain is expected within 6 hours after application. Observe all precautions and limitations on the labels for BUTYRAC® 200 and Alanap-L.

BUTYRAC® 200 Broadleaf Herbicide Plus Basagran® Herbicide for Soybeans.

For improved postemergence control of annual morningglories and cocklebur.

To the label rate of Basagran Herbicide selected according to the other weed species present, add 2 fluid ounces per acre of BUTYRAC 200 broadleaf herbicide for annual morningglories and cockleburs.

Broadcast the tank mix of BUTYRAC® 200 Broadleaf Herbicide and Basagran Herbicide when weeds are actively growing and before annual morningglory vines are a maximum of 6 inches long in the South (AL, AR, FL, GA, LA, MS, NC, OK, SC, TN, TX, VA) or a maximum of 10 inches long in all other states. This is normally up to the time when soybeans have two expanded trifoliate leaves. If application is delayed until weeds are larger than maximum size stated on the Basagran herbicide label, control will be inadequate.

Use a minimum of 20 gallons of total spray mixture per acre with ground equipment and 40 psi pressure with flat fan or hollow cone nozzles spaced 20 inches apart.

Fill the spray tank half way with water, add the recommended amount of $BUTYRAC \otimes 200$ and Basagran Herbicides while the agitator is running, then finish filling.

The tank mix is effective partly through contact action, so weeds must be thoroughly covered with spray. Canopies of large crop and weed leaves shelter smaller weeds and prevent adequate spray coverage. Soybeans tolerate the tank mix of BUTYRAC 200 broadleaf herbicide and Basagran; however, under certain conditions soybean leaves may burn, crinkle and bronze.

NOTE: Observe all precautions and limitations on the labels for ${\tt BUTYRAC}\ 200$ and ${\tt Basagran}$.