

<b>US ENVIRONMENTAL PROTECTION AGENCY</b> <b>OFFICE OF PESTICIDES PROGRAMS</b> <b>REGISTRATION DIVISION (TS-767)</b> <b>WASHINGTON, DC 20460</b>	<b>EPA REGISTRATION NO.</b> 52951-3	<b>DATE OF ISSUANCE</b> <b>AUG 24 1987</b>
	<b>TERM OF ISSUANCE</b>	
	<b>NAME OF PESTICIDE PRODUCT</b> Py-Sol-Mint	

**NOTICE OF PESTICIDE:** ☒ REGISTRATION  
☐ REREGISTRATION  
(Under the Federal Insecticide, Fungicide,  
and Rodenticide Act, as amended)

**NAME AND ADDRESS OF REGISTRANT (Include ZIP code)**

Foural Enterprises  
204 Hillcrest Avenue  
Somerset, NJ 08873

**NOTE:** Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment:
  - a. Add the phrase "EPA Registration No. 52951-3."
  - b. Include the subheadings:
    - Pesticide Disposal
    - Container Disposalin the appropriate places in the Storage and Disposal section.
  - c. Add a Note to Physician statement to read:

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.


☒ ATTACHMENT IS APPLICABLE

<b>SIGNATURE OF APPROVING OFFICIAL</b>	<b>DATE</b>
--	-------------

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

  
John H. Lee  
Product Manager (31)  
Disinfectants Branch  
Registration Division (TS-767C)

Enclosures

For best results apply BUTYRAC® 200 Broadleaf Herbicide to peanuts when the weeds are 1 to 2 inches in size. For ground application, use 10 to 30 gallons of spray solution per acre. For air application use a minimum of 5 gallons of spray solution per acre.

Alabama, Arkansas, Florida, Georgia, Louisiana, North Carolina, South Carolina, Tennessee

Apply 0.8 pint to 1.0 pint (12.8 to 16 fluid ounces) of BUTYRAC® 200 Broadleaf Herbicide per acre on peanuts 2 to 12 weeks after planting. For late germinating cockleburs and morningglories, make a second application about 3 weeks after the first one.

In the Virginia-Carolina area, do not apply later than 45 days before harvest.

Oklahoma, Texas, New Mexico

Apply 0.8 to 1.6 pint (12.8 to 25.6 fluid ounces) of BUTYRAC® 200 Broadleaf Herbicide per acre. Use the low rate on morningglories and cockleburs up to 12 inches in size. Use the higher rate on other weeds as listed above. BUTYRAC® 200 Broadleaf Herbicide may be applied twice during the season. For best suppression of Sida make applications 14 days apart. The second application should be made not more than 100 days after planting or 30 days before harvest.

Do not feed treated peanut vines or peanut hay to livestock. Do not apply BUTYRAC® 200 Broadleaf Herbicide if peanut plants are under stress from drought as injury may occur.

WEED CONTROL IN SEEDLING ALFALFA, BIRDSFOOT TREFOIL, ALSIKE, LADINO OR RED CLOVER

Best results will be obtained by applying BUTYRAC® 200 Broadleaf Herbicide to spring- or fall-seeded legumes after their emergence when seedling weeds are small, no more than 2-3 inches high or rosettes less than 2 inches across and not bolting. Later application to more mature weeds or to new growth from old established weeds will not be satisfactory.

Weed and crop seedlings should be growing strongly for greatest selectivity. Control is difficult under cold or droughty conditions. Rainfall within 7-10 days after treatment can cause injury. For irrigated crops, apply BUTYRAC® 200 Broadleaf Herbicide as soon as possible after irrigation. Delay the next irrigation for at least 7 to 10 days after spraying.

WEED CONTROL IN ESTABLISHED ALFALFA

Certain weeds will emerge in the fall and over-winter in the rosette stage. Best control of these weeds will result from postemergence treatment in late fall or early winter rather than in spring when weeds are no more than 2-3 inches high or rosettes less than 2 inches across. Follow state recommendations for timing and rates to be used under local conditions.

Rainfall within 10 days after treatment can cause injury.

## Directions for use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**FOR GENERAL DISINFECTION** of hard, non-porous inanimate surfaces (glass, metal, plastic, glazed porcelain, linoleum, tile, leather, vinyl, enamel and finished wood surfaces) on walls, floors, appliances and furnishings in industrial and commercial offices, restrooms and work areas.

**CLEANING AND DEODORIZING.** Whenever practical, flush away gross soils with water. For cleaning and deodorizing, use 3 to 8 ounces of THIS PRODUCT per gallon of water. Cleaning may be accomplished with a clean mop, sponge, squeegee, wet vacuum system, or a mechanical spraying device. After cleaning remove suspended soils with a clear water rinse, vacuum pick up or by wiping with a clean mop, sponge, squeegee or cloth.

**DISINFECTING.** Before disinfecting, pre-clean surfaces as directed above. Then, prepare 3 ounces of THIS DISINFECTANT per gallon of water and thoroughly wet the surfaces to be disinfected. The disinfecting solution must contact the surface for 10 minutes for proper disinfection. If desired, the surfaces may then be rinsed with clear water or wiped with a clean damp cloth.

**FOR CLEANING AND DEODORIZING OF GARBAGE CANS, GARBAGE TRUCKS, INDUSTRIAL WASTE RECEPTACLES AND OTHER GARBAGE HANDLING EQUIPMENT.** It is especially important to pre-clean, in order for the disinfectant to perform properly. Then apply a wetting solution of 8 ounces of THIS PRODUCT per gallon of water.

**FOR CLEANING OF WALLS, FLOORS AND INDUSTRIAL WORK AREAS.** Use 3-8 ounces per gallon of water. The use of rubber gloves is recommended for personnel doing continuous hand cleaning.

## Fy-Sol-Mint

DISINFECTANT, CLEANER, DEODORANT  
Pleasantly mint scented

### ACTIVE INGREDIENT:

Alkyl (C <sub>12</sub> , 57%; C <sub>16</sub> , 28%; C <sub>18</sub> , 14%)	
dimethyl benzyl ammonium chloride	2.0%
INERT INGREDIENTS	98.0%
<b>TOTAL INGREDIENTS</b>	<b>100.0%</b>

Contains methyl salicylate

### KEEP OUT OF REACH OF CHILDREN DANGER

**Statement of Practical Treatment.** In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, raw egg whites, gelatin solution or, if these are not available, drink a large quantity of water. Avoid alcohol. Call a physician immediately.

See side panel for additional precautionary statements.

### FOR INDUSTRIAL AND COMMERCIAL USE ONLY.

EPA Reg. No.

EPA Est. No. 52951-NJ-01

Manufactured by:

FOURAL Enterprises

204 Hillcrest Ave.

Somerset, NJ 08873

NET CONTENTS:

1 U.S. Gal.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

AUG 24 1987

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

52951-3

This DISINFECTANT is effective against gram-positive and gram-negative bacteria.

Kills: *Salmonella choleraesuis* ATCC 6538, *Streptococcus aureus* ATCC 11222, *Escherichia coli* ATCC 11222, *Shigella sonnei*, *Candida albicans*.

THIS DISINFECTANT is also especially effective in controlling odors.

THIS DISINFECTANT mixes with water to form a clear stable solution. It is non-staining to plastic, vinyl, and most common metals. Has a pleasant mint odor.

THIS DISINFECTANT may be used on scrubbing equipment.

This product must not be used on food surfaces.

### STORAGE AND DISPOSAL

Store only in tightly closed, inaccessible to children. Do not use for storage or disposal.

Pesticide wastes are acutely toxic. If these wastes cannot be disposed of by the instructions, contact your State Agency, or the hazardous waste Regional Office for guidance.

Do not reuse empty container. Thoroughly rinse empty container with water. Do not reuse for other purposes approved by local authorities, or for disposal of smoke.

### PRECAUTIONS HAZARD TO HUMANS AND DOMESTIC ANIMALS

Causes eye damage and skin irritation. Wear gloves when handling. Harmful to fish and aquatic life. Avoid breathing dust or smoke.

PHYSICAL AND CHEMICAL: Flammable near heat or open flame.

ENVIRONMENTAL HAZARD: Do not discharge into lake or stream unless in accordance with local regulations.

Do not contaminate water when disposing of wastes.

BEST AVAILABLE COPY

alfalfa is in the CRP cover crop treated with BUTYRAC® 200, some temporary twisting of stems and malformation of leaves may occur.

Rainfall within 10 days after treatment can cause injury to legumes in CRP areas.

Overhead sprinkler irrigation should be delayed as long as possible (10 days or more) following application of BUTYRAC® 200 to avoid washing the chemical into the root zone and causing legume injury.

SEE ADDITIONAL USE PRECAUTIONS BELOW

BUTYRAC 200 BROADLEAF HERBICIDE DIRECTIONS FOR USE IN CRP AREAS CONTAINING GRASSES (MINIMUM 6 LEAF STAGE & TILLERING) AND SEEDLING OR ESTABLISHED ALFALFA AND SEEDLING BIRDSFOOT TREFOIL, ALSIKE, LADINO OR RED CLOVER

SEEDLING BROADLEAF WEEDS CONTROLLED:

Common lambsquarters, redroot pigweed, field pennycress, wild mustard, common ragweed, cocklebur, yellow rocket, Russian thistle, annual morningglory species:

Less than 1 inch high.....	1 to 2 quarts per acre
1 to 3 inches high.....	2 to 3 quarts per acre

Smartweed, curly dock:

Up to 3 inches high.....	3 quarts per acre
--------------------------	-------------------

This material does not adequately control over-wintered field pennycress and mustards.

Use the higher rate in dry, low humidity areas.

Apply BUTYRAC 200 broadleaf herbicide in 10 or more gallons of water per acre when using ground equipment and in at least 5 gallons per acre with aerial equipment. The volume of spray applied should provide good coverage.

SEE ADDITIONAL USE PRECAUTIONS BELOW.