

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
& DOMESTIC ANIMALS
CAUTION**

Avoid inhalation, ingestion or absorption through skin. May cause irritation of nose, throat or skin. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water or wetlands. Do not apply where runoff conditions are likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL AND CHEMICAL HAZARDS

Sulfur dust suspended in air irritates eyes. Keep away from heat, sparks and flame. Do not smoke while applying this product.

DIRECTIONS FOR USE

This violation of Federal law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

Do not enter treated area without protective clothing and respiratory protection for 24 hours following application.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

When normal warnings must be given to workers who are expected to be in a treated area within a reasonable time after application of this product, written warnings are given. Warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

CAUTION

Area treated with LX Flowable Sulfur. Do not enter within appropriate protective clothing for 24 hours following application in case of accidental exposure. See the Statement of Practical Treatment on this label.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store unopened product in original container only in cool, dry area, out of reach of children and animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse for equipment. Then offer for recycling or reuse, donate, or dispose of in a sanitary landfill, or by incineration, if allowed by state and local authorities.

PRODUCT INFORMATION

LX Flowable Sulfur is a flowable sulfur formulation that may be applied as a ground or aerial application. Consult your State Agricultural Experiment Station or Extension Specialist for advice in selecting treatments from this label to best fit local conditions.

USE ONLY ON CROPS LISTED AND ONLY AS SPECIFIED ON THIS LABEL EXCEPT ON COTTON: Do not apply when temperatures exceed or are likely to exceed 90°F. Do not use sulfur with oil or within 14 days of an oil spray (for citrus 21 days). Do not combine with emulsifiable liquids.

NOTE: Sulfur will cause severe fruit and leaf injury to sulfur sensitive crops. Do not apply or allow to drift to sprouts of Avocado and Citrus peels, cranberries, cucurbits (cucumber, cantaloupe, melons, squash, Hubbard), spinach, tung trees, walnuts or other sensitive plants. Sulfur may burn foliage and fruit during periods of high temperatures and under certain climate conditions.

LX Flowable Sulfur can be mixed and applied with water-soluble sulfur, liquid fertilizer or water. **DO NOT STOP AGITATION WITH SULFUR IN SPRAY EQUIPMENT. SULFUR IS HIGHLY CORROSIVE AND EQUIPMENT SHOULD BE CLEANED THOROUGHLY AFTER EACH DAY'S SPRAYING.**

RECOMMENDATIONS:

FRUIT TREE CROPS

Unless otherwise specified, for specific crop dosage rates are given as parts of LX Flowable Sulfur per 100 gallons water for use in a thorough cover spray. The total amount of LX Flowable Sulfur per acre should be 7 to 8 gallons depending on the size of the tree. For use of sulfur in the form of spray alone, use a single 1/2 inch high tree is treated. For high volume sprayers (output 80 to 100 gallons per acre) use the low rate per 100 gallons. For low volume sprayers (output 20 to 30 gallons per acre) use the high rate per 100 gallons.

Apply in sufficient water for thorough coverage. For aerial application use a minimum of 5 gallons of finished spray per acre.

NOT REVIEWED

In Accordance with FR Notice 92-2.
Based on Draft Labeling Dated 2/27/86

BEST AVAILABLE COPY

LX

FLOWABLE SULFUR

ACTIVE INGREDIENT:

Sulfur as S...	53.1
INERT INGREDIENTS	46.9
TOTAL	100.0

This product contains 6 lbs. of Sulfur per gallon

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

IF ON SKIN: Wash thoroughly with soap and water. Get medical attention.

IF IN EYES: Wash eyes with plenty of clear water for at least 15 minutes.

See side panel for additional precautionary statements.

EPA Est. No. 37429 GA 01
EPA Reg. No. 53285-1

NET CONTENTS



MANUFACTURED FOR
W. R. LANDIS ASSOCIATES, INC.
P. O. BOX 5126
VALDOSTA, GA 31603 5126

APPLE, PEAR

Scab
Pre-bloom through calyx sprays 1 1/2 to 3% parts
Cover Sprays 1/2 to 2 parts

Powdery Mildew

Pre-bloom through calyx sprays 3 parts
Cover Sprays 2 parts

CHERRY, PLUM, PRUNE:

Brown Rot, leafspot
Pink and Bloom Sprays 1 1/2 to 3% parts
Petiole fall, shuck & cover 1/2 to 1 1/2 parts

CITRUS:

Rust Mite, cover mist
Rust Mite sprays November thru May 1/2 to 4% parts
THRIPS (Early spring treatment for the reduction of Thrips population) Apply 1 to 3 gallons per acre in sufficient water for thorough coverage. Make application after spring flush has made 3 to 4 inch growth. Repeat if necessary. Do not apply within 21 days of an oil application.

PEACH, NECTARINE

Powdery Mildew, brown rot, scab
Pink and Bloom Sprays 1 1/2 to 3% parts
Petiole Fall, shuck and cover sprays 1/2 to 1 1/2 parts

Application to mature nectarines may cause discoloration.

FIELD AND VINE CROPS:

Unless otherwise specified for specific crops, dosage rates are given as parts of LX Flowable Sulfur per acre. This product should be applied by ground equipment or aircraft in sufficient water to provide thorough coverage.

For aerial application use a minimum of 5 gallons of finished spray per acre.

COTTON: Atlantic Mine 2 parts for early season control and up to 1 gallon for mid to late season control.

GRAPES: Powdery Mildew Use 1 to 2 parts per 100 gallons water, maximum of 8 parts LX Flowable Sulfur per acre in spring and summer. Make first application when shoots are 6 to 8 inches long, second when shoots are 12 to 15 inches long, and third at about bloom stage. The timing, rate and number of applications will vary with local conditions. Consult State Agricultural Extension Service or State Agricultural Experiment Stations in your region for specific information. Follow the USE PRECAUTIONS given on this label.

PEANUTS: For rust use 2 to 4 parts per acre of LX Flowable Sulfur in enough water for good coverage. Begin 40 to 60 days after planting. Use 7 to 14 day intervals as long as needed to control rust.

POTATOES: Powdery Mildew Use 3 to 4 parts in 8 gallons water per acre by air. Apply at the first sign of infection and repeat at 3 week intervals.

SUGAR BEETS: For control of powdery mildew apply the recommended rate in 80 gallons of water per acre by air at first sign of infection. Use 8 parts per acre in Arizona and 4 parts per acre in other states. Repeat applications should be made at 10 to 30 day intervals or as necessary throughout the season. Under most disease conditions only one or two applications will be needed.

STRAWBERRIES: Powdery Mildew Use 2 parts. Apply at first sign of infection and repeat at 3 week intervals. For Strawberries that will be canned, discontinue application of sulfur well before harvest in accordance with local canners' recommendations.

TOMATOES: Tomato rust mite Use 1/2 to 1 gallon per acre depending upon size of plants. Use full coverage sprays.

WARRANTY STATEMENT

W. R. LANDIS ASSOCIATES, INC., warrants that this product in its unopened package conforms to the chemical description on the label and is reasonably fit for the purposes set forth on the label when used according to directions under normal use conditions.

There are no other warranties, expressed or implied, concerning the use of the product other than indicated on the label. This warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to seller and buyer assumes all risk of any such use.

Calson Printing Co. 31788

3/13/86

PM 21 53285-1 Pg. 1 of 1