

53431-14

01/05/2010

1/5



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Antimicrobials Division (7510-P)
1200 Pennsylvania Avenue N.W.
Washington, D.C. 20460

EPA Reg. Number:
53431-14

Date of Issuance:
Jan. 5, 2010

Term of Issuance:
Conditional

Name of Pesticide Product:
1512

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

ChemCal Inc
635 Westport Parkway Suite 312
Grapevine, TX. 76051

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Antimicrobials Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product (OPP Decision No. 417614) is conditionally registered in accordance with FIFRA sec 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and, submit acceptable responses required for re-registration of your product under FIFRA section 4.
2. Change EPA File Symbol 53431-RU to EPA Registration Number 53431-14.

Submit one copy of the finished final printed label prior to releasing this product for sale.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e).

Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of approved label is enclosed for your records.

Signature of Approving Official:

Wanda Y. Henson
Wanda Y. Henson
Acting Product Manager 32
Antimicrobials Division (7510-P)

Date:

January 5, 2010



INCORPORATED

ISO 9001:2000

Where water and technology meet™

1512

SODIUM HYPOCHLORITE SOLUTION

ACTIVE INGREDIENT:

Sodium Hypochlorite.....	12.5%
OTHER INGREDIENTS.....	87.5%
TOTAL	100.0%

Available Chlorine: 11.9%

KEEP OUT OF REACH OF CHILDREN
Danger

First Aid

If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a Poison Control Center or doctor for treatment advice.

If Swallowed: Call a Poison Control Center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.

If Inhaled: Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a Poison Control Center or doctor for treatment advice.

"Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage."

*Have product container or label with you when calling a Poison Control Center or doctor for treatment advice. Contact the Poison Center at 1-800-222-1222 for 24 hour emergency medical treatment information.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS

Danger: CORROSIVE. Causes severe skin and eye irritation or chemical burns. Causes eye damage. May be fatal if swallowed. Wear safety glasses or goggles and rubber gloves when handling this product. Do not get in eyes, on skin or on clothing. Irritating to nose and throat. Do not breathe vapor or spray mist. Vacate poorly ventilated areas. Do not return until strong odors dissipate. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, using tobacco, or using restroom. Remove and wash contaminated clothing before reuse.

Environmental hazards: This product is toxic to fish and aquatic organisms. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

PHYSICAL AND CHEMICAL HAZARDS:

STRONG OXIDIZING AGENT: Mix only with water according to label directions. Mixing this product with chemicals (e.g. ammonia, acids, detergents, etc.) or organic matter (e.g. urine, feces, etc.) will release chlorine gas which is irritating to eyes, lungs and mucous membranes.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Note: ChemCal 1512 degrades with age. Use a chlorine test kit and increase dosage, as necessary, to obtain the required level of available chlorine.

ACCEPTED

JAN 05 2010

Under the Federal Insecticide, Fungicide, and
Rodenticide Act as amended, for the
pesticide, registered under
EPA Reg. No. 53431-04

COOLING TOWERS / EVAPORATIVE CONDENSER WATER

SLUG FEED METHOD – Initial Dose: When system is noticeably fouled, apply 52 to 104 oz. of this product per 10,000 gallons of water in the system to obtain from 5 to 10 ppm available chlorine. Repeat until control is achieved.

Subsequent Dose: When microbial control is evident, add 11 oz. of this product per 10,000 gallons of water in the system daily, or as needed to maintain control and keep the chlorine residual at 1 ppm. Badly fouled systems must be cleaned before treatment is begun.

INTERMITTENT FEED METHOD – Initial Dose: When system is noticeably fouled, apply 52 to 104 oz. of this product per 10,000 gallons of water in the system to obtain 5 to 10 ppm available chlorine. Apply half (or 1/3, 1/4, or 1/5) of this initial dose when half (or 1/3, 1/4, or 1/5) of the water in the system has been lost by blowdown.

Subsequent Dose: When microbial control is evident, add 11 oz. of this product per 10,000 gallons of water in the system to obtain a 1 ppm residual. Apply half (or 1/3, 1/4, or 1/5) of this initial dose when half (or 1/3, 1/4, or 1/5) of the water in the system has been lost by blowdown. Badly fouled systems must be cleaned before treatment is begun.

CONTINUOUS FEED METHOD – Initial Dose: When system is noticeably fouled, apply 52 to 104 oz. of this product per 10,000 gallons of water in the system to obtain 5 to 10 ppm available chlorine.

Subsequent Dose: Maintain this treatment level by starting a continuous feed of 1 oz. of this product per 1,000 gallons of water lost by lowdown to maintain a 1 ppm residual. Badly fouled systems must be cleaned before treatment is begun.

SEWAGE AND WASTEWATER TREATMENT

EFFLUENT SLIME CONTROL – Apply a 100 to 1000 ppm available chlorine solution at a location which will allow complete mixing. Prepare this solution by mixing 10 to 100 oz. of this product with 100 gallons of water. Once control is evident, apply a 15 ppm available chlorine solution. Prepare this solution by mixing 3 oz. of this product with 100 gallons of water.

FILTER BEDS – SLIME CONTROL: Remove filter from service, drain to a depth of 1 ft. above filter sand, and add 80 oz. of product per 20 sq/ft evenly over the surface. Wait 30 minutes before draining water to a level that is even with the top of the filter. Wait for 4 to 6 hours before completely draining and backwashing filter.

PUBLIC SYSTEMS: Mix a ratio of 1 oz. of this product to 100 gallons of water. Begin feeding this solution with a hypo chlorinator until a free available chlorine residual of at least 0.2 ppm and no more than 0.6 ppm is attained throughout the distribution system.

DISINFECTION OF DRINKING WATER (EMERGENCY /INDIVIDUAL SYSTEMS)

PUBLIC SYSTEMS: Mix a ratio of 1 oz. of this product to 100 gallons of water. Begin feeding this solution with a hypo chlorinator until a free available chlorine residual of at least 0.2 ppm and no more than 0.6 ppm is attained throughout the distribution system. Check water frequently with a chlorine test kit. Bacteriological sampling must be conducted at a frequency no less than that prescribed by the National Primary Drinking Water Regulation. Contact your local Health Department for further details.

PUBLIC WATER SYSTEMS

NEW WELLS – Flush the casing with a 50 ppm available chlorine solution of water containing 5 oz. of this product for each 100 gallons of water. The solution should be pumped or fed by gravity into the well after thorough mixing with agitation. The well should stand for several hours or overnight under chlorination. It may then be pumped until a representative raw water sample is obtained. Bacterial examination of the water will indicate whether further treatment is necessary.

AGRICULTURAL USES

FRUIT & VEGETABLE WASHING – Thoroughly clean all fruits and vegetables in a wash tank. Thoroughly mix 5 oz. of this product in 200 gallons of water to make a sanitizing solution of 25 ppm available chlorine. After draining the tank, submerge fruit or vegetables for 2 minutes in a second wash tank containing the recirculating sanitizing solution. Spray rinse vegetables with the sanitizing solution prior to packing. Rinse fruit with potable water only prior to packaging. Rinse fruit with potable water only prior to packaging.

COMMERCIAL LAUNDRY SANITIZER


Wet Fabrics or clothing should be spun dry prior to sanitization. Thoroughly mix 2 oz. of this product with 10 gallons of water to yield 200 ppm available chlorine. Promptly after mixing the sanitizer, add the solution into the prewash prior to washing fabrics/clothing in the regular wash cycle with good detergent. Test the level of available chlorine, if solution has been allowed to stand. Add more of this product if the available chlorine level has dropped below 200 ppm.

MILK PLANT AND DAIRY EQUIPMENT

Bottles, bottle fillers, coolers, pasteurizers, chums, separators, utensils and other equipment may be sanitized by using 1512. Surfaces or articles to be sanitized must be thoroughly pre-cleaned. Rinse all surfaces with the sanitizing solution prepared by adding 2 oz. of this product with 10 gallons of water to provide 200 ppm available chlorine by weight. Maintain contact for at least 2 minutes after sanitizing. Do not rinse equipment with water after treatment and do not soak equipment overnight.

MEAT AND POULTRY PROCESSING PLANTS

Surfaces or articles to be sanitized must be thoroughly pre cleaned. Solutions containing 600 ppm available chlorine by weight (6 1/2 oz. 1512 per 10 gallons water) may be used for sanitizing walls, floors, ceilings and similar areas. Maintain contact with sanitizer for at least 2 minutes. Do not rinse after sanitizing. Solutions containing 200 ppm available chlorine (2 oz. per 10 gallons water) may be used on edible product equipment. Maintain contact time with sanitizer for at least 2 minutes. Do not rinse after sanitizing. Do not allow the solution to come directly in contact with meat or poultry.



FOOD EGG SANITIZATION

Thoroughly clean all eggs. Thoroughly mix 2 oz. of this product with 10 gallons warm water to produce a 200 ppm available chlorine solution. The sanitizer temperature should not exceed 130°F. Spray the warm sanitizer so that the eggs are thoroughly wetted. Allow the eggs to thoroughly dry before casing or breaking. Do not apply a potable water rinse. The solution must not be reused to sanitize eggs.

RESTAURANTS AND TAVERNS

Pre-clean dinnerware and utensils with dishwasher detergent and rinse with clean water, immerse dinnerware and utensils in sanitizing solution prepared by adding 2 oz. of this product with 10 gallons of water to provide 200 ppm available chlorine by weight. Maintain contact with sanitizer for at least 2 minutes and allow sanitizer to drain. Do not rinse utensils with water after treatment.

MACHINE DISHWASHING TERMINAL RINSE SANITIZATION

As a terminal sanitizing rinse for pre-cleaning food utensils, adjust automatic dispensing equipment to provide a sanitizing use solution of 100 to 200 ppm available chlorine by weight according to requirements of public health authorities. Use solution should be tested frequently with a suitable chlorine test kit to ascertain that the rinsate strength does not fall below 50 ppm. In absence of a test kit starting concentration of 200 ppm should be used. Do not rinse utensils with water after treatment.

SANITIZATION OF NONPOROUS FOOD CONTACT SURFACES

RINSE METHOD - A solution of 100 ppm available chlorine may be used in the sanitizing solution if a chlorine test kit is available. Solutions containing an initial concentration of 100 ppm available chlorine must be tested and adjusted periodically to ensure that the available chlorine does not drop below 50 ppm. Prepare a 100 ppm sanitizing solution by thoroughly mixing 1 oz of this product with 10 gallons of water. If no test kit is available, prepare a sanitizing solution by thoroughly mixing 2 oz. of this product with 10 gallons of water to provide approximately 200 ppm available chlorine by weight.

Clean equipment surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizing solution for at least 2 minutes. If solution contains less than 50 ppm available chlorine, as determined by suitable test kit, either discard the solution or add sufficient product to reestablish a 200 ppm residual. Do not rinse equipment with water after treatment and do not soak equipment overnight.

Sanitizers used in automated systems may be used for general cleaning but may not be re-used for sanitizing purposes.

IMMERSION METHOD - A solution of 100 ppm available chlorine may be used in the sanitizing solution if a chlorine test kit is available. Solutions containing an initial concentration of 100 ppm available chlorine must be tested and adjusted periodically to ensure that the available chlorine does not drop below 50 ppm. Prepare a 100 ppm sanitizing solution by thoroughly mixing 1 oz of this product with 10 gallons of water. If no test kit is available, prepare a sanitizing solution by thoroughly mixing 2 oz. of this product with 10 gallons of water to provide approximately 200 ppm available chlorine by weight.

Clean equipment surfaces in the normal manner. Prior to use, immerse equipment in the sanitizing solution for at least 2 minutes and allow the sanitizer to drain. If solution contains less than 50 ppm available chlorine, as determined by suitable test kit, either discard the solution or add sufficient product to reestablish a 200 ppm residual. Do not rinse equipment with water after treatment and do not soak equipment overnight.

Sanitizers used in automated systems may be used for general cleaning but may not be re-used for sanitizing purposes.

FLOW PRESSURE METHOD - Disassemble equipment and thoroughly clean after use. Assemble equipment in operating position prior to use. Prepare a volume of a 200 ppm available chlorine sanitizing solution equal to 110% of volume capacity of the equipment by mixing the product in a ratio of 2 oz. per 10 gallons of water. Pump solution through the system until full flow is obtained at all extremities, the system is completely filled with the sanitizer and all air is removed from the system. Close drain valves and hold pressure for at least 2 minutes to ensure contact with all internal surfaces. Remove some cleaning solution from the drain valve and test with a chlorine test kit. Repeat entire cleaning/sanitizing process if effluent contains less than 50 ppm available chlorine.

CLEAN IN PLACE METHOD - Thoroughly clean equipment after use. Prepare a volume of a 200 ppm available chlorine sanitizing solution equal to 110% of volume capacity of the equipment by mixing the product in a ratio of 2 oz. per 10 gallons of water. Pump solution through the system until full flow is obtained at all extremities, the system is completely filled with the sanitizer and all air is removed from the system. Close drain valves and hold pressure for at least 10 minutes to ensure contact with all internal surfaces. Remove some cleaning solution from the drain valve and test with a chlorine test kit. Repeat entire cleaning/sanitizing process if effluent contains less than 50 ppm available chlorine.

SPRAY/FOG METHOD - Pre-clean all surfaces after use. Use a 200 ppm available chlorine solution to control bacteria, mold or fungi; and a 600 ppm solution to control bacteriophage. Prepare a 200 ppm sanitizing solution of sufficient size by thoroughly mixing the product in a ratio of 2 oz. product with 10 gallons of water. Prepare a 600 ppm solution by thoroughly mixing the product in a ratio of 6 oz. product with 10 gallons of water. Use spray or fogging equipment which can resist hypochlorite solutions. Always empty and rinse spray/fog equipment with potable water after use. Thoroughly spray or fog all surfaces until wet, allowing excess sanitizer to drain. Vacate area for at least 2 hours. Prior to using equipment, rinse all surfaces with a 200 ppm solution.

SANITIZATION OF POROUS FOOD CONTACT SURFACES

RINSE METHOD - Prepare a 600 ppm sanitizing solution by thoroughly mixing 6 oz of this product with 10 gallons of water. Clean equipment surfaces in the normal manner. Rinse all surfaces thoroughly with the 600 ppm solution maintaining contact for at least 2 minutes. Prepare a 200 ppm sanitizing solution of sufficient size by thoroughly mixing the product in a ratio of 2 oz. product with 10 gallons of water. Prior to using equipment, rinse all surfaces with a 200 ppm available chlorine solution. Do not rinse and do not soak equipment overnight.

IMMERSION METHOD - Prepare a 600 ppm solution by thoroughly mixing, in an immersion tank, 6 oz. of this product with 10

ACCEPTED
JAN 05 2010
53431-14

gallons of water. Clean equipment in the normal manner. Immerse equipment in the 600 ppm solution for at least 2 minutes. Prepare a 200 ppm sanitizing solution of sufficient size by thoroughly mixing the product in a ratio of 2 oz. product with 10 gallons of water. Prior to using equipment, immerse in a 200 ppm available chlorine solution. Do not rinse equipment with water after treatment and do not soak equipment overnight.

SPRAY/FOG METHOD – Pre-clean all surfaces after use. Prepare a 600 ppm solution by thoroughly mixing, in an immersion tank, 6 oz. of this product with 10 gallons of water. Use spray or fogging equipment which can resist hypochlorite solutions. Always empty and rinse spray/fog equipment with potable water after use. Thoroughly spray or fog all surfaces until wet, allowing excess sanitizer to drain. Vacate area for at least 2 hours. Prior to using equipment, rinse all surfaces with a 200 ppm available chlorine solution.

SANITIZATION OF NONPOROUS NON-FOOD CONTACT SURFACES

RINSE METHOD – Prepare a sanitizing solution by thoroughly mixing 2 oz. of this product with 10 gallons of water to provide approximately 200 ppm available chlorine by weight. Clean equipment surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizing solution for at least 2 minutes. Do not rinse equipment with water after treatment and do not soak equipment overnight.

IMMERSION METHOD – Prepare a sanitizing solution by thoroughly mixing, in an immersion tank, 2 oz. of this product with 10 gallons of water to provide approximately 200 ppm available chlorine by weight. Clean equipment in the normal manner. Prior to use, immerse equipment in the sanitizing solution for at least 2 minutes and allow the sanitizer to drain. Do not rinse equipment with water after treatment.

SPRAY/FOG METHOD – Pre-clean all surfaces after use. Prepare a 200 ppm solution by thoroughly mixing 2 oz. of this product with 10 gallons of water. Use spray or fogging equipment which can resist hypochlorite solutions. Always empty and rinse spray/fog equipment with potable water after use. Thoroughly spray or fog all surfaces until wet, allowing excess sanitizer to drain. Vacate area for at least 2 hours.

SANITIZATION OF POROUS NON-FOOD CONTACT SURFACES

RINSE METHOD - Prepare a 600 ppm sanitizing solution by thoroughly mixing 6 oz of this product with 10 gallons of water. Clean equipment surfaces in the normal manner. Prior to using equipment, rinse all surfaces with the disinfecting solution, maintaining contact with the sanitizer for at least 2 minutes. Do not rinse equipment with water and do not soak equipment overnight.

IMMERSION METHOD – Prepare a 600 ppm solution by thoroughly mixing, in an immersion tank, 6 oz. of this product with 10 gallons of water. Clean equipment in the normal manner. Prior to using equipment in the sanitizing solution for at least 2 minutes and allow the sanitizer to drain. Do not rinse equipment with water after treatment.

SPRAY/FOG METHOD – After cleaning, sanitize non-food contact surfaces with a 600 ppm available chlorine by thoroughly mixing 6 oz. of this product with 10 gallons of water. Use spray or fogging equipment which can resist hypochlorite solutions. Always empty and rinse spray/fog equipment with potable water after use. Thoroughly spray or fog all surfaces until wet, allowing excess sanitizer to drain. Vacate area for at least 2 hours.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.
PESTICIDE STORAGE: Store in original container, tightly closed away from heat or open flame. Store in a cool, dry, well ventilated place. Keep container off wet floors.
PESTICIDE DISPOSAL: Do not put spilled product, filled or partially filled containers in the trash. Dispose of waste material according to label use directions otherwise call 1-800-CLEANUP for guidance from your local solid waste agency.
CONTAINER HANDLING: Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container promptly after emptying. For containers less than or equal to 5 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or mix tank. Fill the container 1/4 full with water and recap. Shake for ten seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.
For containers over 5 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.
Offer for recycling if available, or place in trash.

EPA REG. NO. XXXXXXXXXXXX EPA EST. NO. 53431-TX-001

NET CONTENTS: Marked on Container

DOT SHIPPING INFORMATION: Hypochlorite Solution, UN-1791

Distributed by:
ChemCal, Inc.

