BIOCONTROL

FFB 0 1 200U



53575-15

PB-ROPE* L

ACTIVE INGREDIENTS:

Z, Z-7, 11-Hexadecadien-1-yl Acetate	46.7 %
Z, E-7, 11-Hexadecadien-1-yl Acetate	44.1 %
INERT INGREDIENTS	9.2 %
· · · · · · · · · · · · · · · · · · ·	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,

Keep out of reach of children CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

STATEMENT OF PRACTICAL TREATMENT

Avoid contact with eyes. In case of contact, immediately flush with water. Get medical attention if irritation persists. Wash hands with soap and water after handling.

ENVIRONMENTAL HAZARDS

Do not discharge into lakes, streams, ponds or public waters unless this product is specifically identified and addressed in a NPDES permit. For guidance contact your regional office of the Environmental Protection Agency.

PACIFIC BIOCONTROL CORPORATION 16010 NE 36th Avenue Ridgefield, WA 98642 U.S.A. Telephone (360) 574-9726 1-800-999-8805

MADE IN JAPAN

NET CONTENTS:

500 Dispenser Units

EPA Est. No: 47265-JP-01 EPA Reg. No: 53575-15 One dispenser contains 0.0056 floz or 147 mg Total content of package: 2.8 floz or 73.5 gm

*PB-Rope is a registered Trademark of Shin-Etsu Chen

Do., Ltd.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal

	•
Pesticide Storage	Store in original unopened package at temperatures below 40°F in a dry location.
Pesticide Disposal	Pesticide that cannot be used according to label instructions must be disposed of according to applicable Federal, State and local government procedures. Contact the State pesticide or EPA Hazardous Waste representative at nearest EPA regional office.
Container Disposal	Dispose of containers in a sanitary landfill, or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELLING.

It is critical that PB-ROPE L is applied as directed.

	It is chucal that PD-HOPE Lis applied as directed.
1. Crop	Cotton.
2. Pest	Pink bollworm moth (Pectinophora gossypiella).
3. Rate	100-200 dispensers per acre or 250-500 dispensers per hectare.
4. Application	 i. Twist dispensers loosely around the main stem of cotton near the bottom of the plant. ii. Apply immediately prior to moth emergence in the field or adjoining area. iii. Dispensers must be applied uniformly throughout the treated acreage to obtain a reduction in mating.
5. Timing	Attach dispensers at pinsquare.
5. Precautions	PB-Rope L suppresses mating of pink bollworm moths. However, if a major source of mated female pink bollworm moths is present in adjacent areas, migration of these moths may significantly reduce the level of control achieved. Sources are likely to be cotton (sprayed or unsprayed) or other host species within 300 yards of the treated field. This may be counteracted by: a. Treatment of entire blocks and not just sections of large conventionally treated fields. b. Treatment of infestation sources with PB-Rope L (e.g. a strip at least 250 yards wide nearest the treated field). c. Treatment of infestation sources with an effective insecticide. Supplementary applications of insecticide are advised when PB-Rope L is used to control very high populations of pink bollworm moths. Other pests must be monitored so that timely intervention with insecticides is possible.

WARRANTY AND LIMITATION OF DAMAGES

All statements concerning the use of this product apply only when used as directed. The Manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extra reyond the description on the label. Read all directions carefully.