

BIOCONTROL



PB-ROPE L

ACCEPTED

3/5/03

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 53575-15

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal

Pesticide Storage	Store in original unopened package at temperatures below 40°F in a dry location.
Pesticide Disposal	Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility.
Container Disposal	Dispose of containers in a sanitary landfill, or by incineration, or, if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

ACTIVE INGREDIENTS:

(Z, Z)-7, 11-Hexadecadien-1-yl Acetate	46.7 %
(Z, E)-7, 11-Hexadecadien-1-yl Acetate	44.1 %
OTHER INGREDIENTS	9.2 %
TOTAL	100.0 %

133.55 mg active ingredients per dispenser

Keep out of reach of children

CAUTION

FIRST AID

- If on Skin or Clothing
- Take off contaminated clothing.
 - Rinse skin immediately with plenty of water for 15-20 minutes.
 - Call a poison control center or doctor for treatment advice.
- If in Eyes
- Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
 - Call a poison control center or doctor for treatment advice.
- Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-999-8805 for further questions.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of this product. Do not discharge into lakes, streams, ponds or public waters unless in accordance with NPDES permit. For guidance contact your regional office of the Environmental Protection Agency.

PACIFIC BIOCONTROL CORPORATION

14615 NE 13th Court, Suite A, Vancouver, WA 98685 U.S.A.
Telephone (360) 571-2247 or 1-800-999-8805

MADE IN JAPAN
EPA Est. No: 47265-JP-01
EPA Reg. No: 53575-15

NET CONTENTS: 500 Dispenser Units
One dispenser contains 0.0056 fl oz or 147 mg
Total content of package: 2.8 fl oz or 73.5 gm

® is a registered Trademark of Pacific Biocontrol Corporation.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
It is *critical* that PB-ROPE L is applied as directed.

- Crop: Cotton.
- Pest: Pink bollworm (*Pectinophora gossypiella*).
- Rate: 100-200 dispensers per acre or 250-500 dispensers per hectare (0.51-1.02 fl oz or 13.36-26.71 gm a.i. per acre). Do not exceed 150 gm a.i. per acre per year.
- Application:
 - Twist dispensers securely around the main stem of cotton near the bottom of the plant or to stakes placed uniformly within the treated field. Dispensers twisted too lightly can constrict stem growth. Avoid placing dispensers in contact with the soil.
 - Apply immediately prior to moth emergence in the field or adjoining area.
 - Dispensers must be applied uniformly throughout the treated acreage to obtain a reduction in mating.
- Timing: Attach dispensers at pinsquare. Dispensers release pheromone for up to 90 days depending on temperature. Consult a local pest control advisor for proper timing.
- Note: PB-Rope L suppresses mating of pink bollworms. However, if a major source of mated female pink bollworm moths is present in adjacent areas, migration of these moths may significantly reduce the level of control achieved. Sources are likely to be cotton (sprayed or unsprayed) or other host species within 300 yards of the treated field. This can be overcome by:
 - Treatment of entire blocks and not just sections of large conventionally treated fields.
 - Treatment of infestation sources with PB-Rope L (i.e., a strip at least 250 yards wide nearest the treated field).
 - Treatment of infestation source with an effective insecticide.
 Supplementary applications of insecticide are advised when PB-Rope L is used to control high populations of pink bollworms. All pests must be monitored so that timely intervention with insecticides is possible.

WARRANTY AND LIMITATION OF DAMAGES

All statements concerning the use of this product apply only when used as directed. The Manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.

53575-15

03/05/2003

1/1