

PM 9/

53575-17

5/24/95

4/2

# BIOCONTROL



## ISOMATE\*-P PHEROMONE

<b>ACTIVE INGREDIENTS:</b>	
3,13 Octadecadien-1-yl Acetate.....	90.0 %
<b>INERT INGREDIENTS.....</b>	<b>10.0 %</b>
<b>TOTAL</b>	<b>100.0 %</b>

**Keep out of reach of children**  
**WARNING**

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**WARNING**

Avoid inhalation of vapors or contact with liquid ingredients. Harmful if swallowed. Avoid contact with eyes. In case of eye contact, immediately flush with water. Get medical treatment if irritation persists. Wash hands thoroughly with soap and water after handling.

**ENVIRONMENTAL HAZARDS**

Do not discharge into lakes, streams, ponds or public waters unless in accordance with NPDES permit. For guidance contact your regional office of the Environmental Protection Agency.

BIOCONTROL LIMITED  
 719 Second Street, Suite 12  
 Davis, California 95616 U.S.A.  
 Telephone (916) 757-2307


MADE IN JAPAN

EPA Est. No: 47255-JP-01  
 EPA EUP No: 53575-EUP-4

NET CONTENTS:  
500 Dispenser Units

One dispenser contains 0.0014 fl oz or 37.8 mg  
 Total content of package: 0.7 fl oz or 18.9 gm

\*ISOMATE is a registered Trade Mark of Biocontrol Limited



2/2

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal

<b>Storage</b>	Store unopened package at temperatures below 40°F in a dry location. Product may be stored in cold storage facilities used for food storage.
<b>Pesticide Disposal</b>	Pesticide that cannot be used according to label instructions must be disposed of according to applicable Federal, State and local procedures.
<b>Container Disposal</b>	Dispose of empty dispensers by burning or burying with prunings in winter. If burned, stay out of smoke. Dispose of dispensers according to applicable Federal, State and local procedures. Foil envelopes can be disposed of as household refuse.
<b>General</b>	Consult Federal, State or local disposal authorities for approved alternative procedures.

## DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

It is critical that ISOMATE-P be applied as directed.

<b>1. Crop</b>	Peach, almond, nectarine, cherry, prune, plum, apricot.
<b>2. Pest</b>	Peach Tree Borer ( <i>Synanthedon scitella</i> ).
<b>3. Rate</b>	Minimum of 100 dispensers per acre or 250 dispensers per hectare. Attach one dispenser per tree. Maximum of 250 dispensers per acre or 625 dispensers per hectare.
<b>4. Application</b>	Attach the dispensers onto branches above the mid point of the tree.
<b>5. Timing</b>	Apply prior to moth emergence in spring.
<b>6. Precautions</b>	Isomate-P prevents Peach Tree Borer from mating. However, if a major source of mated female moths of these species is present adjacent to the treated field, migration of these moths may significantly reduce the level of control achieved. Sources are likely to be untreated host trees within 300 yards of the treated field. This can be overcome by: <ul style="list-style-type: none"> <li>a. Treatment of entire blocks and not just sections of large conventionally treated fields which frequently serve as sources of mated females.</li> <li>b. Treatment of infestation sources with Isomate-P (e.g. a strip at least 50 yards wide nearest the treated field).</li> <li>c. Treatment of infestation source with an effective insecticide.</li> </ul> Other pests must be monitored and controlled by timely intervention with conventional insecticides.

## WARRANTY AND LIMITATION OF DAMAGES

All statements concerning the use of this product apply only when used as directed. The Manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.