53575-19



NOV - 7 2008

Ms. Kathy Bolan Pacific Biocontrol Corporation 575 Viewridge Drive Angwin, CA 94508

 Subject: Product Name: Isomate-M 100 EPA Registration No. 53575-19 Label Amendment to update ingredient statement, net contents and rate; addition of mandatory and advisory statements; adoption of current container disposal statements; update pesticide storage; maximum rate in dispensers under Directions for Use; updated "Application" under Direction for Use; updated "Note" under Directions for Use; revision warranty statement. Application Dated: September 3, 2008

Dear Ms. Bolan:

The amendment referred to above, submitted in connection with registration under FIFRA section 3(c)(5), is acceptable provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.

2. Submit three (3) copies of your final printed labeling before you release the product for shipment. Final printed labeling means the label or labeling of the product when distributed or sold. Clearly legible reproductions or photo reductions will be accepted for unusual labels, such as those silk-screened directly onto glass or metal containers or large bags or drum labels.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(b). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

If you have any questions contact Andrew Bryceland at 703-305-6928 or by email at: <u>bryceland.andrew@epa.gov</u>. A stamped copy of the label is enclosed for your records.

Sincerely,

Linda A. Hollis, Chief Biochemical Pesticides Branch

		21001101110				
	cobiopestivides and Pollution					
SYMBOL	75118	Prevention	Division (75	l 1P)		
	Bricaland					
DATE	10/28/04					
EPA Form 1320-1A (1/90)		Drimtad on Regulad	Danor			

\* U.S. Government Printing Office: 2005

## ACCEPTED LOT #

NOV - 7 2008 BIOCONTROL

Under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, for the pesticide registered under

EPA Reg. No. 53575-19 ISOMATE®-M 100

A MATING DISRUPTION FORMULATION FOR ORIENTAL FRUIT MOTH (Grapholita molesta). MACADAMIA NUT BORER (Cryptophiebia ombrodelta), KOA SEEDWORM (Cryptophiebia Illepida)

ACTIVE INGREDIENTS:	1
Z-8-Dodecen-1-yl Acetate	
E-8-Dodecen-1-yl Acetate	5.68 %
Z-8-Dodecen-1-ol	
OTHER INGREDIENTS	4.79 %
TOTAL	
232.13 mg active ingredients per dispen	ser

### **KEEP OUT OF REACH OF CHILDREN** CAUTION

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Avoid contact with skin, eyes or clothing: Harmful if absorbed through skin. Causes moderate eye irritation. After handling, wash thoroughly with soap and water before eating, drinking, chewing gum, using tobacco or using the tollet. Remove and wash contaminated clothing before reuse.

#### EIDST AID STATEMENT

	FIRSTAIDSTATEMENT
IF ON SKIN OR CLOTHING:	<ul> <li>Remove contaminated clothing.</li> <li>Immediately rinse skin for 15-20 minutes using plenty of water.</li> <li>Contact a poison control center or doctor for treatment advice.</li> </ul>
IF IN EYES:	<ul> <li>Hold eyes open, slowly and gently rinse with water for 15-20 minutes. If wearing contact lenses, rinse eyes for 5 minutes, remove contact lenses, then continue rinsing.</li> <li>Contact a poison control center or doctor for treatment advice.</li> </ul>
center or door Pesticides in week. During	UMBER: Have the product container or label with you when contacting a poison control tor, or going for treatment. For emergency information on this product, call the National formation Center at 1-800-858-7878, 6:30AM to 4:30PM Pacific time (PT), seven days a other times, call the poison control center at 1-800-222-1222. You may also contact Pacific 1-800-999-8805 for additional information.
	ENVIRONMENTAL HAZARDS
	uses Do not apply directly to water, to areas where surface water is present or to intertidal he mean high water-mark Do not contaminate water when disposing of this product.
	PACIFIC BIOCONTROL CORPORATION

·	
PACIFIC BIOCONT	ROL CORPORATION
146 5 NE 13th Court; Suite A	Vancouver, WA 98685 U.S.A.
	2247 or 1-800-999-8805
MADE IN APAN	NET CONTENTS: 400 Dispenser Units
EPA Est. No: 47265-JP-01	One dispenser contains 0.0094 fl oz or 243.79 mg
EPA Reg. No: 53575-19	Total content of package: 3.75 fl oz or 97.52 gm

43.79 mg Total content of package: 3.75 fl oz or 97.52 gm

ISOMATE® is a registered Trademark of Pacific Biocontrol Corporation

and other pome fruits; almond, macadamia and other tree nut crops.         Target Pests       Oriental fruit moth (Grapholita molesta), macadamia nut borer (Cryptophlebia ombrod koa seedworm (Cryptophlebia illepida).         Rate       100-150 dispensers per acre or 250-375 dispensers per hectare (0.89-1.34 fl oz or 2 34.82 gm a.i. per application). Do not exceed 150 grams (or 647 dispensers) a.i. per per year.         Application       Place dispensers securely to lateral branches as high as can be reached in canopy. Apply dispensers uniformly in the treated area. Dispensers twisted too tightly girdle branches.         Timing       Apply prior to moth emergence in the spring. In crops with long field seasons, a sea application can provide better control. If subsequent applications are required, apply prite start of subsequent flights. Consult your local pest control advisor or Pacific Bioc representative for proper timing or estimated formulation longevity in your area.         Note       Isomate-M 100 suppresses mating of target pests. Immigration of mated female mot these species from adjacent external sources of infestation will reduce the level of co Manage by: <ul> <li>a. Treatment of external sources of infestation with long the advised.</li> <li>c. Treatment of pheromone treated orchard with insecticide.</li> <li>c. Treatment of pheromone treated orchard with insecticide.</li> </ul> Areawide application is most effective. Supplementary applications of insecticide are advised Isomate-M 100 is used in orchards with high pest populations. All pests must be monitored so that to somate-M 100 is used in orchards with high pest populations. All pests must be monitored so that to somate-M 100 is used in orchards with hi		DIRECTIONS FOR USE
and other pome fruits; almond, 'macadamia and other tree nut crops.         Target Pests       Oriental fruit moth (Grapholita molesta), macadamia nut borer (Cryptophlebia ombrod koa seedworm (Cryptophlebia illepida).         Rate       100-150 dispensers per acre or 250-375 dispensers per hectare (0.89-1.34 fl oz or 2 34.82 gm a.i. per application). Do not exceed 150 grams (or 647 dispensers) a.i. per per year.         Application       Place dispensers securely to lateral branches as high as can be reached in canopy. Apply dispensers uniformly in the treated area. Dispensers twisted too tightly girdle branches.         Timing       Apply prior to moth emergence in the spring. In crops with long field seasons, a se application can provide better control. If subsequent applications are required, apply prite start of subsequent flights. Consult your local pest control advisor or Pacific Bioc representative for proper timing or estimated formulation longevity in your area.         Note       Isomate-M 100 suppresses mating of target pests. Immigration of mated female mot these species from adjacent external sources of infestation will reduce the level of co Manage by: <ul> <li>a. Treatment of external sources of infestation with long.</li> <li>b. Treatment of external sources of infestation with insecticide.</li> <li>c. Treatment of pheromone treated orchard with insecticide.</li> </ul> Areawide application is most effective. Supplementary applications of insecticide are advised Isomate-M 100 is used in orchards with high pest populations. All pests must be monitored so that to somate-M 100 is used in orchards with high pest populations. All pests must be monitored so that to somate-M 100 is used in orchards with high pes	IT IS A	INCONSISTENT WITH ITS LABELING.
koa seedworm (Cryptophlebia Illepida).         Rate       100-150 dispensers per acre or 250-375 dispensers per hectare (0.89-1.34 fl oz or 2 34.82 gm a.i. per application). Do not exceed 150 grams (or 647 dispensers) a.i. per per year.         Application       Place dispensers securely to lateral branches as high as can be reached in canopy. Apply dispensers uniformly in the treated area. Dispensers twisted too tightly girdle branches.         Timing       Apply prior to moth emergence in the spring. In crops with long field seasons, a se application can provide better control. If subsequent applications are required, apply pr the start of subsequent flights. Consult your local pest control advisor or Pacific Bloc representative for proper timing or estimated formulation longevity in your area.         Note       Isomate-M 100 suppresses mating of target pests. Immigration of mated female mot these species from adjacent external sources of infestation will reduce the level of co Manage by: a. Treatment of external sources of infestation with an effective insecticide. c. Treatment of external sources of infestation with an effective insecticide.         Areawide application is most effective. Supplementary applications of insecticide are advised Isomate-M 100 is used in orchards with high pest populations. All pests must be monitored so that t	Сгорз	Peaches, nectarines, apricots, plums, cherries and other stone fruits; apples, pears, quinc and other pome fruits; almond, macadamia and other tree nut crops.
34.82 gm a.i. per application). Do not exceed 150 grams (or 647 dispensers) a.i. per per year.         Application       Place dispensers securely to lateral branches as high as can be reached in canopy. Apply dispensers uniformly in the treated area. Dispensers twisted too tightly girdle branches.         Timing       Apply prior to moth emergence in the spring. In crops with long field seasons, a se application can provide better control. If subsequent applications are required, apply prite start of subsequent flights. Consult your local pest control advisor or Pacific Block representative for proper timing or estimated formulation longevity in your area.         Note       Isomate-M 100 suppresses mating of target pests. Immigration of mated female moth these species from adjacent external sources of infestation will reduce the level of comparison of external sources of infestation with somate-M 100.         b. Treatment of external sources of infestation with an effective insecticide.         c. Treatment of external sources of infestation with an effective insecticide.         c. Treatment of external sources of infestation with an effective insecticide.         c. Treatment of external sources of infestation so f insecticide.         c. Treatment of external sources of infestation so f insecticide.         c. Treatment of external sources of infestation so f insecticide.         c. Treatment of external sources of infestation so f insecticide are advised lsomate-M 100 is used in orchards with high pest populations. All pests must be monitored so that the sources of infestation so f insecticide are advised	Target Pests	Oriental fruit moth (Grapholita molesta), macadamia nut borer (Cryptophlebla ombrodelta koa seedworm (Cryptophlebia illepida).
<ul> <li>canopy. Apply dispensers uniformly in the treated area. Dispensers twisted too tightly girdle branches.</li> <li>Timing Apply prior to moth emergence in the spring. In crops with long field seasons, a sea application can provide better control. If subsequent applications are required, apply prior to subsequent flights. Consult your local pest control advisor or Pacific Biocor representative for proper timing or estimated formulation longevity in your area.</li> <li>Note Isomate-M 100 suppresses mating of target pests. Immigration of mated female moth these species from adjacent external sources of infestation will reduce the level of comparison of the even of external sources of infestation will reduce the level of comparison. Treatment of external sources of infestation with an effective insecticide.</li> <li>Areawide application is most effective. Supplementary applications of insecticide are advised Isomate-M 100 is used in orchards with high pest populations. All pests must be monitored so that the sources of infestation with an effective insecticide.</li> </ul>	Rate	100-150 dispensers per acre or 250-375 dispensers per hectare (0.89-1.34 fl oz or 23.2 34.82 gm a.i. per application). Do not exceed 150 grams (or 647 dispensers) a.i. per acr per year.
application can provide better control. If subsequent applications are required, apply pr the start of subsequent flights. Consult your local pest control advisor or Pacific Biocon representative for proper timing or estimated formulation longevity in your area. Note Isomate-M 100 suppresses mating of target pests. Immigration of mated female mot these species from adjacent external sources of infestation will reduce the level of con Manage by: a. Treatment of external sources of infestation with Isomate-M 100. b. Treatment of external sources of infestation with an effective insecticide. c. Treatment of pheromone treated orchard with insecticide. Areawide application is most effective. Supplementary applications of insecticide are advised Isomate-M 100 is used in orchards with high pest populations. All pests must be monitored so that t	Application	Place dispensers securely to lateral branches as high as can be reached in the canopy. Apply dispensers uniformly in the treated area. Dispensers twisted too tightly ma girdle branches.
these species from adjacent external sources of infestation will reduce the level of co Manage by: a. Treatment of external sources of infestation with Isomate-M 100. b. Treatment of external sources of infestation with an effective insecticide. c. Treatment of pheromone treated orchard with insecticide. Areawide application is most effective. Supplementary applications of insecticide are advised Isomate-M 100 is used in orchards with high pest populations. All pests must be monitored so that t	Timing	Apply prior to moth emergence in the spring. In crops with long field seasons, a secon application can provide better control. If subsequent applications are required, apply prior to the start of subsequent flights. Consult your local pest control advisor or Pacific Blocontro representative for proper timing or estimated formulation longevity in your area.
intervention with insecticides is possible.	Areawide app Isomate-M 10	<ul> <li>a. Treatment of external sources of infestation with Isomate-M 100.</li> <li>b. Treatment of external sources of infestation with an effective insecticide.</li> <li>c. Treatment of pheromone treated orchard with insecticide.</li> <li>lication is most effective. Supplementary applications of insecticide are advised when the sources of insecticide in orchards with high pest populations. All pests must be monitored so that time</li> </ul>
STORAGE AND DISPOSAL Do not contaminate water, food or feed by storage or disposal.		

Pesticide Storage	<ul> <li>Store in original unopened package in a dry location at temperatures below Do not store in cold facilities used for food.</li> </ul>		
Pesticide Disposal	Wastes resulting from this product may be disposed of waste disposal facility	on site or at an approved	
Container Disposal	Nonrefillable container. Do not reuse or refill this contain packet in the trash.	iner. Dispose of empty foil	

# WARRANTY AND LIMITATION OF DAMAGES

All statements concerning the use of this product apply only when used as directed. Pacific Biocontrol Corporation warrants that the product complies with the specifications expressed in this label. To the extent consistent with applicable law, Pacific Biocontrol Corporation makes no other warranties, and disclaims all other warranties, expressed or implied, including but not limited to warranties of merchantability and fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. To the extent consistent with applicable law, Pacific Biocontrol Corporation's liability or default, breach or failure under this label shall be limited to the amount of the purchase price. To the extent consistent with applicable law, Pacific Biocontrol Corporation shall have no liability for damages (special, consequential or incidental) that result from handling, storage and use of this product, which is not in compliance with the label. Use of this product implies all directions of this label have been ready carefully. Read all directions carefully.

Revision Date: September 2008