

BIOCONTROL



STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal

Pesticide Storage	Store in original unopened package at temperatures below 40°F in a dry location. Product may be stored in cold storage facilities used for food storage.
Pesticide Disposal	Pesticide that cannot be used according to label instructions must be disposed of according to applicable Federal, State and local government procedures. Contact the State pesticide or EPA Hazardous Waste representative at nearest EPA regional office.
Container Disposal	Dispose of dispensers in sanitary landfill or incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke. Foil envelopes can be disposed of as household refuse.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELLING.

It is *critical* that ISOMATE-CMLR is applied as directed.

- 1. Crop** Apple and pear.
- 2. Pest** Codling moth (*Cydia pomonella*), oblique banded leafroller (*Choristoneura rosaceana*), pandemis leafroller (*Pandemis pyrusana*).
- 3. Rate** 400 dispensers per acre or 1000 dispensers per hectare (3.9 fl oz or 102 gm a.l. per application). Apply double rate of dispensers to edges of orchard.
- 4. Application** Apply dispensers in at least upper third of tree, preferably within 2-3 feet of treetop. Apply dispensers within canopy and on branches to maximize shade protection. Apply dispensers securely on lateral branches. Dispensers twisted too tightly may girdle branches. Can be applied efficiently from moving trailer or with a pole applicator.
- 5. Timing** Apply prior to codling moth emergence in the spring. Dispensers release pheromone for 100 -120 days. In crops with long field seasons (i.e. more than 120 days) or in orchards with high pest populations, a second application is recommended. If subsequent applications are required, apply prior to the start of subsequent flights.
- 6. Precautions** Isomate-CMLR suppresses codling moth and leafroller moth from mating. However, if a major source of mated female moths of these species is present adjacent to the treated field, migration of these moths may significantly reduce the level of control achieved. Sources are likely to be unsprayed apple, pear, peach, apricot, plum, and cherry trees or other host species within 300 yards of the treated field. This can be overcome by:
 - Treatment of **entire blocks** and not just sections of large conventionally treated fields that frequently serve as sources of mated females.
 - Treatment of infestation sources with Isomate-CMLR (e.g. a strip at least 50 yards wide nearest the treated field).
 - Treatment of infestation source with an effective insecticide.
 - Treatment of 4-6 rows along border of pheromone treated orchard with insecticide.
 Other pests must be monitored on a regular schedule so that timely intervention with conventional insecticides is possible.

WARRANTY AND LIMITATION OF DAMAGES

All statements concerning the use of this product apply only when used as directed. The Manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.

53575-20
2/1/2000

ACCEPTED

FEB 01 2000

...icide.
... Act.
... pesticide
... under
EPA Reg. No. 53575-20

ISOMATE*-CM/LR PHEROMONE

ACTIVE INGREDIENTS:

Z-11-Tetradecenyl acetate.....	45.8%
E, E-8, 10-Dodecadien-1-ol.....	39.2%
Dodecanol.....	6.6%
Tetradecanol.....	1.0%

INERT INGREDIENTS.....7.4%

TOTAL.....100.0%

Keep out of reach of children
CAUTION

STATEMENT OF PRACTICAL TREATMENT:

- IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.
- IF IN EYES:** Flush eyes with plenty of water. Get medical attention if irritation persists.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of this product. Do not discharge into lakes, streams, ponds or public waters unless in accordance with NPDES permit. For guidance contact your regional office of the Environmental Protection Agency.

PACIFIC BIOCONTROL CORPORATION

16010 NE 36th Avenue
Ridgefield, WA 98642 U.S.A.
Telephone (360) 574-9726
1-800-999-8805

MADE IN JAPAN

EPA Est. No: 47265-JP-01
EPA Reg. No: 53575-20

NET CONTENTS:

400 Dispenser Units
One dispenser contains 0.01 fl oz or 264 mg
Total content of package: 4.1 fl oz or 105 gm

*ISOMATE is a registered Trademark of Biocontrol Ltd.