

ACCEPTED

AUG 18 2002

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 53575-21

BIOCONTROL



ISOMATE®-BAW PHEROMONE

ACTIVE INGREDIENTS:	
(Z,E)-9,12-Tetradecadien-1-yl Acetate.....	55.0 %
Z-9-Tetradecen-1-ol.....	24.8 %
OTHER INGREDIENTS..... 20.2 %	
TOTAL..... 100.0 %	
201.38 mg active ingredients per dispenser	

**Keep out of reach of children
CAUTION**

FIRST AID

- If in Eyes**
- Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
 - Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-999-8805 for further questions.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of this product. Do not discharge into lakes, streams, ponds, or public waters unless in accordance with NPDES permit. For guidance contact your regional office of the Environmental Protection Agency.

PACIFIC BIOCONTROL CORPORATION

14615 NE 13th Court, Suite A, Vancouver, WA 98685 U.S.A.
Telephone (360) 571-2247 or 1-800-999-8805

MADE IN JAPAN
EPA Est. No. 47265-JP-01
EPA Reg. No. 53575-21

NET CONTENTS: 500 Dispenser Units
One dispenser contains 0.0097 fl oz or 252.65 mg
Total content of package: 4.86 fl oz or 126.3 gm

ISOMATE® is a registered Trademark of Pacific Biocontrol Corporation.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal

Pesticide Storage	Store in original unopened package at temperatures below 40°F in a dry location. Only unopened or unbroken dispenser packages may be stored in cold storage facilities used for food storage. Care must be taken to avoid contamination of food or feed items.
Pesticide Disposal	Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility.
Container Disposal	Dispose of dispensers in sanitary landfill or by incineration, or if allowed by State and Local authorities, by burning. If burned, stay out of smoke. Foil envelopes can be disposed of as household refuse.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

It is critical that ISOMATE-BAW is applied as directed.

- 1. Crop** Alfalfa, asparagus, beans, beets, cabbage, celery, cole crops, cotton, cucumbers, lettuce, onions, peanuts, peas, peppers, soybeans, strawberries, sweet potatoes, tomatoes, tobacco.
- 2. Pest** Beet armyworm (*Spodoptera exigua*).
- 3. Rate** Minimum of 100 dispensers per acre or 250 dispensers per hectare (0.77 fl oz or 20.1 gm a.i. per acre). Maximum of 200 dispensers per acre or 500 dispensers per hectare (1.5 fl oz or 40.3 gm a.i. per acre). Do not exceed 150 gm a.i. per acre per year.
- 4. Application** Attach dispensers to stakes placed uniformly within the treated field at the canopy level of the crop. Avoid placing dispensers in contact with the soil. At 100 dispensers per acre, stakes are to be placed approximately every 20 feet apart. Treat all border rows. Increase the number of dispensers on upwind side of the fields or along borders adjacent to other beet armyworm hosts. Higher rates may be needed in smaller fields or in windy conditions.
- 5. Timing** Monitor with pheromone traps and crop inspection. Apply product prior to adult flight, early enough to disrupt mating communication within the field. Consult your local pest control advisor for proper timing. Dispenser releases pheromone for up to 60-90 days depending on temperature.
- 6. Note** Isomate-BAW suppresses mating between beet armyworms. However, if a major source of mated female moths of this species is present adjacent to the treated field, migration of these moths may significantly reduce the level of control achieved. A common source of mated females is unsprayed host species within 300 yards of the treated field. This can be overcome by:
 - a. Treatment of **entire blocks** and not just sections of large conventionally treated fields that frequently serve as sources of mated females.
 - b. Treatment of infestation sources with Isomate-BAW (e.g. a strip at least 50 yards wide nearest the treated field).
 - c. Treatment of infestation source with an effective insecticide.
 Supplementary applications of insecticide are advised when Isomate-BAW is used to control high populations of beet armyworm, and when beet armyworm larvae are imported into the field on transplants. All pests must be monitored so that timely intervention with insecticides is possible.

WARRANTY AND LIMITATION OF DAMAGES

All statements concerning the use of this product apply only when used as directed. The Manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.

53575-21

8/18/2002

1/1