
 <p>U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Biopesticides and Pollution Prevention Division (7511C) 1200 Pennsylvania Avenue NW Washington, DC 20460</p> <p>NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration (under FIFRA, as amended)</p>	EPA Reg. Number: 53575-25	Date of issuance: MAR 23 2001
	Term of Issuance: Unconditional	
	Name of Pesticide Product: Isomate-C TT	
Name and Address of Registrant (include ZIP Code): Pacific Biocontrol Corporation 14615 NE 13 th Court Vancouver, WA 98685		
<p>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Biopesticides and Pollution Prevention Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p> <p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This registration does not eliminate the need for continual reassessment of the pesticide. If EPA determines at any time, that additional data are required to maintain in effect an existing registration, the Agency will require submission of such data under section 3(c)(2)(B) of FIFRA.</p> <p>This product is registered in accordance with FIFRA section 3(c)(5) and is subject to the following terms and conditions:</p> <ol style="list-style-type: none"> 1. Modify your label by inserting the new EPA Reg. No. 53575-25 before the release of your product for shipment. 2. Submit five (5) copies of the revised final printed labeling before you release the product for shipment. 		
Signature of Approving Official: 	Date: March 23, 2001	

ACCEPTED

MAR 23 2001

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 43575-25

BIOCONTROL



ISOMATE®-C TT

ACTIVE INGREDIENTS:

(E, E)-8, 10-Dodecadien-1-ol.....	53.0 %
Dodecanol.....	29.7 %
Tetradecanol.....	6.0 %
OTHER INGREDIENTS.....	11.3 %

TOTAL.....	100.0 %
382.4 mg active ingredients per dispenser	

Keep out of reach of children
WARNING

FIRST AID

- | | |
|-------------------------------|--|
| If on Skin or Clothing | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |
| If in Eyes | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-999-8805 for further questions.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

WARNING

Harmful if absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of this product. Do not discharge into lakes, streams, ponds or public waters unless in accordance with NPDES permit. For guidance contact your regional office of the Environmental Protection Agency.

PACIFIC BIOCONTROL CORPORATION
14615 NE 13th Court, Vancouver, WA 98685 U.S.A.
Telephone (360) 571-2247 or 1-800-999-8805

MADE IN JAPAN	NET CONTENTS: 400 Dispenser Units
EPA Est. No: 47265-JP-01	One dispenser contains 0.017 fl oz or 431.4 mg
EPA Reg. No: 53575-EL	Total content of package: 6.81 fl oz or 172.56 gm

ISOMATE[®] is a registered trademark of Pacific Biocontrol Corporation

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal

- | | |
|---------------------------|---|
| Pesticide Storage | Store in original unopened package at temperatures below 40°F in a dry location. Product may be stored in cold storage facilities used for food storage. |
| Pesticide Disposal | Pesticide that cannot be used according to label instructions must be disposed of according to applicable Federal, State and Local government procedures. Contact the State pesticide or EPA Hazardous Waste representative at nearest EPA regional office. |
| Container Disposal | Dispose of dispensers in sanitary landfill or by incineration, or if allowed by State and Local authorities, by burning. If burned, stay out of smoke. Foil envelopes can be disposed of as household refuse. |

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
It is *critical* that ISOMATE-C TT is applied as directed.

- | | |
|-----------------------|---|
| 1. Crop | Apple, pear, walnut, quince, peach, prune, plum, peach, pecan and nectarine. |
| 2. Pest | Codling moth (<i>Cydia pomonella</i>), hickory shuckworm (<i>Cydia caryana</i>). |
| 3. Rate | 200 dispensers per acre or 500 dispensers per hectare (3.0 fl oz or 76.5 gm a.l. per application). Apply double rate of dispensers to edges of orchard. Do not exceed 150 gm a.l. per acre per year. |
| 4. Application | Apply dispensers in upper third of tree, preferably within 2-3 feet of treetop. Apply dispensers within canopy. Apply dispensers securely on branches. Can be applied efficiently from moving trailer or with a pole applicator. |
| 5. Timing | Apply prior to codling moth emergence in the spring. Dispensers release pheromone for 120-140 days depending on temperature. In crops with long field seasons (i.e. more than 120 days), a second application is recommended. If subsequent applications are required, apply prior to the start of subsequent flights. Consult your local pest control advisor for proper timing. |
| 6. Note | Isomate-C TT suppresses codling moth and hickory shuckworm from mating. However, if a major source of mated female moths of these species is present adjacent to the treated field, migration of these moths may significantly reduce the level of control achieved. Sources are likely to be unsprayed apple, pear, quince, walnut, plum, crabapple trees or other host species within 300 yards of the treated field. This can be overcome by: <ul style="list-style-type: none"> a. Treatment of entire blocks and not just sections of large conventionally treated fields that frequently serve as sources of mated females. b. Treatment of infestation sources with Isomate-C TT (e.g. a strip at least 50 yards wide nearest the treated field). c. Treatment of infestation source with an effective insecticide. d. Treatment of 4-6 rows along border of pheromone treated orchard with insecticide. Supplementary applications of insecticide are advised when Isomate-C TT is used to control very high populations of codling moth or hickory shuckworm. All pests must be monitored so that timely intervention with insecticides is possible. |

WARRANTY AND LIMITATION OF DAMAGES

All statements concerning the use of this product apply only when used as directed. The Manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.

2/a