

MICROBIOCIDES #2

BEST AVAILABLE COPY

ACCEPTED
with COMMENTS
in EPA Letter Dated:

AUG 6 1985

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

53793-1

ACTIVE INGREDIENT:

Poly(oxyethylene (dimethylamino) ethylene-
(dimethylamino) ethylene dichloride) 10.0 %

INERT INGREDIENT 90.0 %

This product contains 0.85 lb. of active ingredient per gallon and weighs
8.49 lb. per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing vapors. Avoid contact with skin, eyes, or clothing.

FIRST-AID: If swallowed, drink promptly a large quantity of milk, eg. j whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes, streams, or ponds. Permits may be required for discharges containing this pesticide into lakes, streams, ponds, or public water. For guidance, contact the regional office of the Environmental Protection Agency.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

MICROBIOCIDES #2 is used to control bacteria and fungi in recirculating commercial and industrial air wash systems. To its use, systems must be cleaned to remove sludge, lime, and other deposits. An initial slug addition of fluid ounces of MICROBIOCIDES #2 per 1000 gallons of water to provide a concentration of 48 to MICROBIOCIDES #2, based on the system, is recommended. Repeat initial dosage

Subsequent slug additions of 1.53 to MICROBIOCIDES #2 per 1000 gallons

120 parts per million of MICROBIOCIDES #2

employed every 2 to 5 days, or as needed. The frequency of addition depends upon the relative amount of bleedoff and the severity of the problem. Slug additions should be made in the sump of MICROBIOCIDES #2

is used to control bacteria in air-washing systems that maintain effective mist air wash systems. To its use, systems should be cleaned to remove

deposits. An initial slug dose of 24.7 to MICROBIOCIDES #2 per 1000 gallons

is recommended. Repeat initial dosage until control is evident. Additions of 16.9 to 41.2 fluid ounces of MICROBIOCIDES #2

per 1000 gallons of water should be employed each 1 to 2 days. The frequency of addition depends upon the relative amount of the bacterial problem. Slug additions may be made in the sump or collection trays of the airwash system.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Do not stack more than four drums high. Leaking or damaged drums for disposal. Spills should be absorbed in vermiculite or sand and disposed of in a sanitary landfill, by incineration, or, if allowed by state and local authorities, by other procedures approved by state and local authorities.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product should be disposed of in a approved waste disposal facility.

CONTAINER DISPOSAL:

PLASTIC: Triple rinse (or equivalent). Then offer for recycling or reuse, or by other procedures approved by state and local authorities.

METAL: Triple rinse (or equivalent). Then offer for recycling or reuse, or by other procedures approved by state and local authorities.

Manufactured by:

Hi-Valley Chemical, Inc.
1151 South Redwood Road Suite 100
Salt Lake City, UT 84104

EPA REG. NO. _____ EPA EST. _____

NET CONTENTS _____