

- Sale of yellow (or any other unacceptable color) fly baits by registrants will be allowed until September 31, 1996.

- Sale of yellow (or any other unacceptable color) fly bait by anyone other than the registrants will be allowed only until December 31, 1997.

A stamped copy of the label, approved with comments, is enclosed for your records. You must submit the revised CSF, labels, and cancellation letter for all other formulas prior to shipment of the revised product. If you have any questions, please contact Teung F. Chin of my Team, at (703) 305-5412.

Sincerely yours,

DHE

Dennis H. Edwards, Jr.
Product Manager 19
Insecticide-Rodenticide Branch
Registration Division (7505C)

Enclosure

53871-3.BFF

Front Panel

STIMUKIL®

FLY BAIT

ACTIVE INGREDIENTS:

Z-9-tricosene.....	0.025%
Methomyl (S-Methyl-N-[(Methylcarbamoyl)oxy]thioacetimidate.....	1.000%
INERT INGREDIENTS.....	98.975%
TOTAL	100.000%

CAUTION: KEEP OUT OF REACH OF CHILDREN

Read Entire Label for Additional Cautions

EPA REG. NO. 53871-3
EPA ESTABLISHMENT NO. 53871-AZ-01

Troy BioSciences Inc.
2620 N. 37th Drive
Phoenix, Arizona 85009

NET WEIGHT
LOT NO.

[Note: This is a master label. Label language that will appear only on STIMUKIL Fly Bait intended for bait station use has been highlighted. Label language that will appear only on STIMUKIL Fly Bait intended for commercial scatter bait use has been *italicized*. All other label language will appear on both labels.]

ACCEPTED
with COMMENTS
in EPA Letter Dated

FEB 09 1996

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under FFA Reg. No.

53871-3

SIDE PANEL

STATEMENT OF PRACTICAL TREATMENT

In case of poisoning call a physician immediately.

If in eyes: Flush eyes with water for 15 minutes. Get medical assistance if irritation persists.

If on skin: Remove contaminated clothing and wash skin immediately with soap and water. Wash contaminated clothing before reuse.

If swallowed: Drink 1 or 2 glasses of water and induce vomiting by touching the back of the throat with finger. Never induce vomiting or give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN: METHOMYL is a cholinesterase inhibitor. ATROPINE is an antidote.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling and before eating, drinking, use of tobacco products or using the toilet. **Do not use this product around or inside homes, or any other place where children are likely to be present.**

For commercial use only. Use only in areas inaccessible to food producing animals. Do not contaminate feed and foodstuffs. Do not apply where poultry or animals, especially dogs and young calves can pick it up or lick it. Wear long sleeved shirt, long pants, shoes, socks, and gloves during loading, application, repairing, and cleaning of application equipment and disposal of the pesticide.

Do not use in edible product areas of food processing plants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

IMPORTANT! If pesticide comes in contact with skin, wash off with soap and water.

ENVIRONMENTAL HAZARDS: *Accessible areas treated with this bait are hazardous to birds and other wildlife. Use with care when applying in areas frequented by wildlife or adjacent to any body of water. Collect, cover or incorporate granules spilled on the soil surface. This pesticide is toxic to birds. Do not apply directly to water, or to areas where surface water is present or intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment and when disposing of equipment washwaters. Do not contaminate water by disposal of wastes.*

5 of 5

WARRANTY- LIMITATION OF DAMAGES: Troy BioSciences Inc. warrants that this material conforms to the chemical description and guaranteed limits of active ingredients as stated on this label. Troy BioSciences Inc. makes no other express or implied warranty of fitness or merchantability of any kind. Troy BioSciences Inc. maximum breach of liability for breach of warranty shall not exceed the purchase price of this product. In no event shall Troy BioSciences Inc. be liable for indirect or consequential damages. This warranty shall not be changed by oral or written agreement unless signed by a duly authorized officer of Troy BioSciences Inc.. Buyer assumes all risk of use or handling whether in accordance with directions or not.

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. STIMUKIL Fly Bait is an improved fly bait formula containing a sex-attractant and a feeding synergist which encourages male and female flies to remain in the treated area and stimulates vigorous feeding on the bait. **STIMUKIL Fly Bait should not be used around or inside homes, or any other place where children are likely to be present.** Specific uses are described in more detail below:

Bait Station Use: Do not use around or inside of homes or other places where children are likely to be present. STIMUKIL Fly Bait may only be used in conjunction with the STIMUKIL Fly Bait Station. Pour STIMUKIL Fly Bait into Fly Bait Station. Replace cover carefully to avoid spillage of bait. Use the product at a rate of approximately two bait stations per 500 square feet of fly feeding area. Hang Fly Bait stations out of reach of food producing animals or pets (at least 4 feet off the ground). Do not hang bait station within reach of children. Use STIMUKIL Fly Bait Station around the outside of broiler houses, kennels, canneries, beverage processing plants, meat and poultry processing plants, food processing plants and refuse dumpsters associated with these establishments, and inside of caged layer houses. May also be in fenced or otherwise enclosed areas inaccessible to children around the outside of restaurants, supermarkets, commissaries, and bakeries.

Scatter Bait Use: Use of scatter bait is limited solely to commercial establishments where children will not be present, such as, feed lots, poultry houses, livestock barns, canneries, beverage plants, meat and poultry processing establishments, and food processing plants. Distribute bait from container or use other device to avoid handling bait directly. Scatter bait (do not apply in piles) at a rate of 1/4 pound per 500 square feet of area. Do not treat areas where animals can feed on bait. Do not apply directly to water. Keep children and pets clear from scatter bait treatment areas.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal. **Pesticide Disposal:** Pesticide or Rinsate that cannot be used according to label instructions must be disposed of according to applicable Federal, State or local governmental procedures. **Waste Disposal:** Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Disposal:** Triple rinse and dispose of according to local governmental procedures.