

2/3

STIMUKIL[®] FLY BAIT

ACTIVE INGREDIENTS:

(Z)-9-tricosene.....	0.040%
Methomyl (S-Methyl-N-[(Methylcarbamoyl)oxy]thioacetimidate).....	1.000%
INERT INGREDIENTS.....	98.960%
TOTAL.....	100.000%

CAUTION KEEP OUT OF REACH OF CHILDREN

Read Entire Label for Additional Cautions

EPA REGISTRATION NO. 53871-3 EPA ESTABLISHMENT NO. 53871-AZ-01

Manufactured By: TROY BIOSCIENCES INCORPORATED, 113 SOUTH 47TH AVENUE, PHOENIX, AZ 85043

FIRST AID	
If swallowed	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow Do not induce vomiting unless told to do so by the poison control center or doctor Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing the eye. Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
NOTE TO PHYSICIAN	
METHOMYL is a cholinesterase inhibitor. ATROPINE is an antidote.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing.

Personal Protective Equipment (PPE)

Applicators and all other handlers must wear:

- Long-sleeved shirt and long pants
- Socks and shoes
- Chemical resistant gloves made of any waterproof material

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, the user should wash thoroughly and change into clean clothing.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds, fish, aquatic invertebrates and mammals. Accessible areas treated with this bait are hazardous to birds and other wildlife. Collect, cover or incorporate granules spilled on the soil surface. Do not apply directly to water, or to areas where surface water is present or intertidal areas below the mean highwater mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. STIMUKIL Fly Bait is an improved fly bait formula containing a sex-attractant and a feeding synergist, which encourages male and female flies to remain in the treated area and stimulates vigorous feeding on the bait. **STIMUKIL Fly Bait should not be used around or inside homes, or any other place where children are likely to be present.** Do not apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Specific uses are described in more detail below:

BAIT STATION USE: Do not use around or inside homes or other places where children are likely to be present. Pour 25 grams or 2 Tablespoons STIMUKIL Fly Bait into Fly Bait Station. Replace cover carefully to avoid spillage of bait. Use the product at a rate of approximately two bait stations per 500 square feet of fly feeding area. Hang Fly Bait Stations out of reach of food producing animals or pets (at least 4 feet off the ground). Do not place bait station within reach of children. When fly populations are high, the STIMUKIL Fly Bait Station should be replenished with STIMUKIL Fly Bait every 3-5 days or when the bait is no longer visible in the bait station. The dead flies and any remaining bait should be disposed of as recommended below and STIMUKIL Fly Bait added to the Fly Bait Station. Use STIMUKIL Fly Bait Station around the outside of broiler houses, kennels, canneries, beverage processing plants, meat and poultry processing plants, food processing plants and refuse dumpsters[®] associated with these establishments, and inside of caged layer houses. May also be used in fenced or otherwise enclosed areas inaccessible to children, and around the outside of restaurants, supermarkets, commissaries, and bakeries.

SCATTER BAIT USE: Use of scatter bait is limited solely to the outside of certain commercial establishments where children will not be present, such as feed lots, poultry houses, livestock barns, canneries, beverage plants, meat and poultry processing establishments, commercial refuse dumpsters which are fenced or otherwise enclosed, and

ACCEPTED

AUG 17 2004

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 53871-3

food processing plants. Do not use around commercial dumpsters that are not enclosed. Do not use around dumpsters in residential settings. Distribute bait from container or use other device to avoid handling bait directly. Scatter bait (do not apply in piles) at a rate of 1/4 pound per 500 square feet of area. Do not treat areas where animals can feed on bait. Do not apply directly to water. Keep children and pets clear from scatter bait treatment areas. When fly populations are high the bait should be replenished every 3-5 days or when the bait is no longer visible in the treated area.

BRUSH ON USE: Mix 1/4 lb STIMUKIL Fly Bait with 4.0 oz of water to form a paste, stir well. Apply paste with brush on walls, windowsills, and support beams where flies usually land, such that children, poultry, cattle or other animals cannot come into contact with the paste. When the fly population is high, the bait should be reapplied at a 3-5 day interval or when the flies no appear to be affected by landing on the treated area.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or by disposal.

Pesticide Storage: Store in a tightly closed container in a cool, dry place out of the reach of children and pets.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of container in a sanitary landfill, or by other procedures approved by state and local authorities.

Bait Station Disposal: The bait station is reusable. When the container becomes cracked or broken, triple rinse (or equivalent) and dispose of according to local governmental procedures.

WARRANTY

Troy Biosciences Incorporated makes no warranty, express or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risks of use and handling of this material when such use and handling are contrary to label instructions.

STIMUKIL IS A REGISTERED TRADEMARK OF TROY BIOSCIENCES INCORPORATED



TROY BIOSCIENCES
INCORPORATED

