

53883-40


5/19/2003

1/3

Martin's

50% MALATHION
CONCENTRATE
4.3 Lbs. Malathion per Gallon

For Use On:
Vegetables ◊ Trees ◊ Ornamentals



ACTIVE INGREDIENT:
Malathion (O,O-dimethyl phosphorodithioate of diethyl mercaptosuccinate) 50%
INERT INGREDIENTS*: 50%
100%

*This product contains petroleum distillates.

KEEP OUT OF REACH OF CHILDREN
CAUTION

Net Contents: _____

FIRST AID	
Have the product container or label with you when calling the poison control center or doctor, or going for treatment.	
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.
<p>Note to Physician: This product may pose an aspiration pneumonia hazard. Contains petroleum distillate. This product is an organophosphate insecticide. If symptoms of cholinesterase are present atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.</p> <p>Emergency Phone Numbers: Transportation or spill, call CHEMTREC 800-424-9500.</p>	

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing of spray mist. Avoid contact with skin, eyes or clothing. Causes moderate eye irritation. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not contaminate food or feed products.

Malathion is a cholinesterase inhibitor. Do not use this product simultaneously with other drugs, insecticides, pesticides or chemicals having cholinesterase inhibiting activity nor within a few days before or after treating with Malathion.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates and aquatic life stages of amphibians. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.
For outdoor use around the home only.

Fly Control: For outdoor use only, mix 2 ¼ qts. per 100 gals. of water. Spray on walls, fences, garbage containers, and wherever flies congregate and breed. Wet all surfaces thoroughly. The addition of 8 Tbsp. of sugar per gal. will extend the life of the bait residue.

Peaches: For control of Spider Mites, Mealy Bugs, Woolly Aphids, Japanese Beetles (adults) and Oriental Fruit Moths, use 1 ½ to 2 Tbsp. of 50% MALATHION CONCENTRATE to 1 gal. water. For Oriental Fruit Moths, apply at petal fall and every 10 to 14 days thereafter until control is achieved. Apply as needed for other pests. Do not apply within 7 days of harvest.

Pecan Trees: For control of Pecan Nut Casebearers, Pecan Phylloxeras, Pecan Budmoths, Spider Mites and Aphids, use 2 Tbsp. of 50% MALATHION CONCENTRATE to 1 gal. of water.

Ornamentals: (Roses, Camellias, Azaleas, Chrysanthemums, Evergreens and many other flowers and shrubs) For control of Aphids, Birch Leaf Miners, Japanese Beetles (adults), Lace Bugs, Leafhoppers, Mealy Bugs, Tarnished Plant Bugs, Thrips and Whiteflies, use 1 tsp. per gal. of water. For control of Bagworms, Tent Caterpillars, Black Scale Crawlers, Leaf Scale, Oyster Shell Scale, Soft Brown Scale and Spider mites, use 2 tsp. per gal. of water. Do not use on certain ferns, including Boston, Maidenhair and Pteris, as well as some species of Crassula.

Vegetables: For Aphids and Spider mites on Tomatoes and Cucumbers; Aphids, Leafhoppers and Mealy Bugs on Potatoes; Mexican Bean Beetles, Leafhoppers and Spider Mites on Beans; Aphids on Peppers, Cabbage and Broccoli, use on 1 tsp. per gal. of water. For control of Onion Thrips on Onions, use 2 tsp. per gal. of water.

DO NOT USE later than 3 days before harvest on the above vegetables with the exception of Cabbage, and this should not be applied later than 7 days before harvest.

3/13

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in cool, dry area away from children and domestic animals.

Disposal: If empty: Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Control Solutions, Inc.
5903 Genoa-Red Bluff
Pasadena, TX 77507-1041
EPA Reg. No. 53883-40
EPA Est. No. 53883-TX-002

Amend to EPA 02-12-03