

53883-72

6.24.2003

1/21

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 2-28-95



United States  
Environmental Protection Agency  
Washington, DC 20460

Registration  
 Amendment  
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 53883 - 72	2. EPA Product Manager George LaRocca	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Permethrin Martin's Horse & Stable Spray	PM# 13	
5. Name and Address of Applicant (Include ZIP Code) Control Solutions, Inc. 5903 Genoa-Red Bluff Pasadena, TX 77507-1041 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

**NOTIFICATION**  
**NOTIFICATION**  
JUN 24 2003

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of Alternate Brand Name per PR Notice 98-10. The name of the product as referenced in the body of the label only is changed to "this product". This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 USC Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
				<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 2.5 gallon		5. Location of Label Directions <input type="checkbox"/> on label	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Joe Blake	Title Director, Regulatory Affairs	Telephone No. (Include Area Code) 800-242-5562
<b>Certification</b> I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received  (Stamped)
2. Signature 	3. Title Director, Regulatory Affairs	
4. Typed Name Joe Blake	5. Date 06-16-03	

**CONTROL  
SOLUTIONS  
INC.**

June 16, 2003

Document Processing Desk (NOTIF)  
Office of Pesticide Programs (7504C)  
U.S. Environmental Protection Agency  
Ariel Rios Building  
1200 Pennsylvania Ave., N.W.  
Washington, DC 20460

Re: Notification per (PR) Notice 98-10, Wording change in body of label  
EPA Reg. No. 53883-72 Permethrin 10 E.W.C.


Dear Sir or Madam:

Enclosed is a Notification per (PR) Notice 98-10. The name of the product in the body of the label is to be changed to "this product". See highlighted text on pages 14 and 15.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 USC Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

If you have any questions or need additional information, please feel free to contact me.

Sincerely,



Joe Blake  
Director, Regulatory Affairs

3/21

[FRONT LABEL PANEL]

**NOTIFICATION**

**JUN 24 2003**

# Permethrin 10 E.W.C.

For Use by Licensed Pest Control Operators or Commercial Applicators Only

- Kills Termites\*[including dry-wood termites](for localized control only)[Optional Statement]
- For Use around foundations, outside buildings, porches, wood piles, wooden fences and decks, patios [Optional Statement]
- Kills troublesome outdoor insects which destroy Lawns, Roses, Flowers, Trees and Shrubs, Vegetables, Fruits, and Nuts [Optional Statement]
- Provides contact and residual control for up to 4 weeks [Optional Statement]
- **KILLS OVER 100 DIFFERENT INSECTS** [Optional Statement]
- Indoor/Outdoor Insecticide [Optional Statement]
- Subterranean Termite\* Control
- Protects Lawns, Trees & Shrubs, Roses & Flowers, Fruits & Nuts, and Vegetables from: Alfalfa loopers, Ants, Aphids, Armyworms, Bagworms, Beetles, Borers, Brown dog ticks, Budworms, Cabbageworms, Chinch bugs, Cicadas, Citrus black flies, Cluster flies, Codling moths, Corn earworms, Crickets, Cutworms, Earwigs, Elm spanworms, Fall cankerworms, Fire Ants, Fleas, Green fruitworms, Gnats, Grasshoppers, Houseflies, Inchworms, Japanese beetles, Leafhoppers, Leafminers, Leafrollers, Lone Star ticks, Lygus bugs, Mealybugs, Millipedes, Mole crickets, Mosquitoes, Moths, Navel orangeworms, Scales, Sod webworms, Sowbugs (pillbugs), Spider mites, Spring cankerworms, Tent caterpillars, Thrips, Tobacco budworms, Weevils, Whiteflies and many others.

**ACTIVE INGREDIENT:**

Permethrin ..... 10.0%

**INERT INGREDIENTS:** ..... 90.0%

**TOTAL:** ..... 100.0%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**See Back/Side Label Panel For Additional Precautionary Statements**

Net Contents \_\_\_\_\_

**[BACK/SIDE LABEL PANEL]**

- **READ ENTIRE LABEL BEFORE USE**
- **FOR PET USE - USE ONLY ON DOGS**

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

<b>FIRST AID</b>	
<b>Have the product container or label with you when calling a poison control center or doctor, or going for treatment.</b>	
If swallowed	<ul style="list-style-type: none"> <li>• Call poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
If inhaled	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
If in eyes	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
If on skin or clothing	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>NOTE TO PHYSICIAN AND VETERINARIANS:</b> Do not administer or instill milk, cream or other substances containing vegetable or animal fats which will enhance absorption of lipophilic substances.	

**[BACK/SIDE LABEL PANEL]**

**Personal Protective Equipment:**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category section chart.

**Applicators and other handlers must wear:** Long-sleeved shirt and long pants, chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, or viton, and shoes plus socks. For exposure in enclosed areas, a respirator a respirator with a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G).

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Controls Statements:**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)] the handler PPE requirements may be reduced or modified as specified in the WPS.

When using in an enclosed area, do not remain in the treated area. Ventilate the area after treatment is completed. Food should be removed or covered during treatments. All food processing surfaces should be covered during treatment or thoroughly cleaned with an effective cleaning compound followed by a potable water rinse before use. When using this product in these areas, apply only when facility is not in operation.

Do not apply as a space spray while food processing is underway. Except in Federally inspected meat and poultry plants, food processing operations may continue when this product is applied as a surface spray with care and in accordance with the directions and precautions given below.

**USER SAFETY RECOMMENDATIONS**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This pesticide is extremely toxic to fish. Do not apply directly to water. Avoid contamination of ornamental fish ponds. Drifts and runoff from treated areas may be hazardous to aquatic organisms in treated areas. Do not contaminate water when cleaning equipment or disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively, visiting the treatment area.

[BACK/SIDE LABEL PANEL]

**PHYSICAL OR CHEMICAL HAZARDS**

Do not use or store near heat or open flame. Do not use in undiluted form.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**Keep people and pets off treated areas until the spray has dried.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear:

- Coveralls
- Chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber or viton.
- Shoes plus socks

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not allow children or pets to contact treated surfaces until spray has dried.

7/21

**[BACK/SIDE LABEL PANEL]**

Do not apply this product through any type of irrigation system.

**SHAKE WELL BEFORE USING**

**THREE STEPS TO MIX AND APPLY:**

1. For lawns and vegetables, measure the area to be sprayed. Calculate square feet by multiplying length by width. For all other applications, apply as a thorough cover spray.
2. Use a clean sprayer. Carefully measure and mix the amount of concentrate and water as directed.
3. Spray as directed. Thorough coverage is important. Flush sprayer with clean water after each use. Make applications when insects first appear. Repeat applications as directed.

**Use Dilution rate as indicated:**

1 fl.oz. = 6 tsp. or 2 Tbsp.

8 fl.oz. = ½ pt. or 1 cup

[BACK/SIDE LABEL PANEL]

**DIRECTIONS FOR APPLYING**

**HOME LAWNS**

<b>Turf</b>	<b>Pests</b>	<b>How to Use</b>
Bentgrass, Bermudagrass, Bluegrass, Dichondra, Fescue, Irish Moss, Merion, St. Augustine Grass	Ants, Armyworms, Brown dog ticks, Chinch bugs, Crickets, Cutworms, Fleas, Grasshoppers, Japanese beetle grubs, Mole crickets, Sod webworms, Ticks that carry Lyme Disease, Leafhoppers	Thoroughly wet down grass a few hours before applying. Home lawns should be no taller than 3 inches at time of application. Slightly more water may be used as long as the amount of product per area is as listed below. For heavy infestations, repeat application after 2 weeks. For ants, crickets and grasshoppers, use ½ fl.oz. in 3 gals. of water to cover 1,000 sq.ft. For other listed insects, use 1½ fl.ozs. in 10 gals. of water to cover 1,000 sq.ft.



**HOUSE PLANTS**

**INDOORS: TO PROTECT PLANTS IN RESIDENTIAL GREENHOUSES AND LATHHOUSES AND FOR INDOOR LANDSCAPING:**

House Plants	Pests	How to Use
Ageratum, Ardisia, Aster, Baby's Breath, Begonia, Birdsnest Fern, Bleeding Heart, Carnation, Chrysanthemum, Coleus, Crown of Thorns, Cockscomb, Cyclamen, Dracaena, Dumbcane, English Ivy, Fuchsia, Gladiolus, Grape Ivy, Juniper, Lily, Marigold, Orchid, Palm, Pansy, Peperomia, Petunia, Philodendron, Piggy-Back Plant, Poinsettia, Portulaca, Pothos, Prayer Plant, Purple Passion, Rabbits Foot Fern, Rose, Schefflera, Snapdragon, Snake Plant, Statice, Velvet Plant, Verbena, Zinnia	Aphids, Armyworms, Corn earworms, Exposed thrips, Fall cankerworms, Inchworms, Japanese beetles, Leafminers, Leafrollers, Mealybugs, Scales, Solder mites, Spring cankerworms, Whiteflies	Mix ¼ fl.oz. in 1 gal. of water. Contact and residual: Use sufficient spray to obtain full coverage of all foliage. If application is to be made indoors, remove infested plants from living areas prior to application. Use in an area not likely to be occupied by people or pets (particularly fish). After spraying, plants can be returned to original location. Do not use more product than recommended, as this may cause some plant injury. Use with care on ferns, palms and fuchsia. As with all plants listed, especially carnations, chrysanthemums, and roses, there exists a wide variety of species and cultivars, and therefore, a wide variety of susceptibility to injury. To ascertain possible phytotoxicity, treat a few plants as described above, wait 24 hours, and check for possible injury.
Velvet Plants	Fungus Gnats	

**Fire Ant Control**

**Individual Mound and Plant Container Treatment**

Mix ½ fl.oz. in 3 gals. of water and apply as a drench using a sprinkler can. Gently apply 1 gallon of the mixture over the surface of each mound. Thoroughly wet mound and surrounding area of 3x4 ft. (12 sq.ft.). For best results apply in cool weather 65<sup>0</sup>-80<sup>0</sup> F early or late in the day. Treat new mounds as they appear. Pressurized sprays may disturb ants and cause migration which will reduce product effectiveness. **Use the same method as above for the control of Fire Ants that may nest in permanently placed containerized ornamentals plants.**

**[BACK/SIDE LABEL PANEL]**

### TREES & SHRUBS

Trees & Shrubs	Pests	How to Use
Arizona, Cypress, Azalea, Birch, Cherry, Non-bearing Citrus, Conifers, Elm, English Ivy, Euonymus, Weeping Fig, Fir, Honeysuckle, Lilac, Mock-orange, Oak, Palm, Philodendron, Pine, Poinsettia, Tulip Poplar, Rhododendron, Taxus	Aphids, Bagworms, Cicadas, Exposed thrips, Fall cankerworms, Inchworms, Leafminers, Leafrollers, Mealybugs, Spider mites, Whiteflies, Gypsy Moth, Tent Caterpillar, Japanese Beetles	Use ¼ fl.oz. in 1 gal. of water. Apply as a thorough spray, wetting leaves and branches to the dripping point. Try to penetrate dense foliage. Spray in the late afternoon or evening, when the temperature ranges from 50 <sup>0</sup> -75 <sup>0</sup> F and when there is little or no wind. Spray at the first sign of insects. Application can be made up to the day of collection.
Cherry, Honeysuckle	Fall webworms	
Conifer	Budworms	
Elm	Elm leaf beetles, Elm spanworms, Fall webworms	
Douglas and Grand Firs	Tussock moths	
Non-bearing Citrus	Citrus black flies	
Oak	Oakworms	

Pine	Pine beetles, Pine moths, Pine needle miners, Needle scales	
------	---	--

On Tent caterpillars, application should be made when caterpillars are young and tents are first noticed. For best results, apply in the late afternoon or evening, when caterpillars have returned to their tents. Wet tents with spray on all sides. A few caterpillars may be seen escaping from the tent; however, they should die within several hours.

**[BACK/SIDE LABEL PANEL]**

**ROSES & FLOWERS (ORNAMENTALS)**

<b>Roses &amp; Flowers (Ornamentals)</b>	<b>Pests</b>	<b>How to Use</b>
Ageratum, Aster, Ardisia, Azalea, Baby's breath, Begonia, Birds-nest fern, Bleeding heart, Carnation, Chrysanthemum, Coleus, Common Ninebark and Snowberry, Crown of thorns, Cockscomb, Cyclamen. Dracaena, Dumbcane, English Ivy, Exacum, Fuchsia, Gladiolus, Grape Ivy, Gold bells, Hypoestes, Ivy, Juniper, Lilac, Marigold, Mock-Orange, Nannyberry, Orchid, Palm, Pansy, Pea shrub, Peperomia, Petunia, Philodendron, Piggy- back Plant, Poinsettia, Portulaca, Pothos, Prayer Plant, Purple Passion, Rabbits foot fern, Rhododendron, Rose, Schefflera, Snapdragon, Snake Plant, Statice Velvet Plant, Verbena, Zinnia	Aphids, Armyworms, Bagworms, Cabbage worms, Cicadas, Cornworms, Cutworms, Exposed thrips, Fall cankerworms, Fall webworms, Fungus gnat, Inchworms, Japanese beetles, Leafminers, Leafrollers, Mealybugs, Scale crawlers, Spring cankerworms, Spider mites, Whiteflies, Rose Chafer, Psyllids, Scale Insects	Use ¼ fl.oz. in 1 gal.of water. Apply as a thorough spray, wetting the plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage. Spray in the later afternoon or evening when the temperature ranges from 50 <sup>0</sup> - 75 <sup>0</sup> F and when there is little or no wind. Spray at the first sign of insects. Application can be made up to the day of collection.

12/21

### FRUITS & NUTS TREES

#### GENERAL DIRECTIONS

Use 1/2 fl.oz. in 1 gal. of water. Apply when insects appear. Wet the plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage.

Fruits & Nuts Trees	Pests	How to Use
Almonds, Pecans	Navel orangeworms, Peach twig borers	Do not make more than 2 applications during hull split and 5 applications per season. Can be applied up to 7 days prior to harvest.
Apples	Green fruitworms, Aphids, Leafrollers(Oblique-banded, Redbanded), Plum curculio, Rosy apple aphids, Spotted tentiform leafminers, Tarnished plant bugs, White apple leafhoppers, Japanese Beetles	Repeat application only as needed to maintain insect control. Do not make more than 3 applications per growing season. Do not apply after petal fall.
Peaches	Borers(Lesser peachtree, Peach twig), Green fruitworms, Plum curculio, Oriental fruit moths, Tarnished plant bugs	Repeat application only as needed to maintain insect control. Do not apply within 7 days of harvest. Do not make more than 8 applications per season.
Pears (dormant through delayed dormant)	Pear psylla	Repeat application only as needed to maintain insect control. Apply during the dormant through delayed dormant growth period only. Do not make more than 2 applications per season.
Pears (summer)	Codling Moths, Green fruitworms, Pear psylla, Aphids	Repeat application only as needed to maintain insect control. Do not make more than 3 applications per season. Can be applied up to 14 days prior to harvest.

### VEGETABLES

Prepare a spray solution at the dilution rates listed for each crop in the chart below. Apply the spray solution at the rate of 1 gallon to cover 440 square feet surface area.

Vegetables	Pests	How to Use
Asparagus (pre-harvest)	Asparagus beetles, Cutworms	Use ¼ fl. oz. in 2 gals. of water. Apply to ensure uniform coverage. Repeat application only as needed to maintain insect control. Do not apply more than 4 applications per crop. Can be applied up to 3 days prior to harvest.
Asparagus (post-harvest)	Larval stages of asparagus beetle; Tarnished plant bugs, Lygus bugs, Adult Japanese beetle	Use ¼ fl. oz. in 1 gal. of water. Treat ferns or bush growth after spear harvest when insects are present. Repeat application only as needed to maintain insect control.
Beans	Mexican bean beetle, Leafhoppers, Flea beetle, Corn earworm, Lygus and Stinkbugs	Use ½ fl. oz. in 1 gal. of water. Repeat application only as needed to maintain insect control. Do not make more than 8 applications per season. Can be applied up to 3 days prior to harvest.
Celery-see Leafy Vegetables		
Cole Crops, Broccoli, Brussels Sprouts, Cabbage, Cauliflower	Beet armyworms, Cabbage aphids and loopers, Diamondback moths, Imported cabbageworms	Use ¼ fl. oz. in 1 gal. of water. Apply every 5 days or as needed. Can be applied up to 1 day prior to harvest. Do not make more than 8 applications on Cauliflower, Brussels Sprouts and Broccoli and 10 applications on Cabbage.
Corn-see Sweet Corn		
Cucumber, Melons, Pumpkins, Squash	Flea beetle, Leafhopper	Use ½ fl. oz. in 1 gal. water. Repeat application only as needed to maintain insect control. Do not make more than 8 applications per season. Can be applied up to 3 days prior to harvest.

<p><u>Leafy Vegetables:</u> Amaranth, Arrugula, Celery, Celtuce, Chervil, Corn salad, Chrysanthemum(edible-leaved and garland), cress(garden and upland), Dandelion, Dock, Endive, Fennel, Lettuce(head and leaf), Okra, Parsley, Purslane(garden and winter), Rhubarb, Spinach, Swiss chard</p>	<p>Alfalfa looper, Aphids, Beet armyworm, Cabbage looper, Corn earworm, Cutworms, European corn borer, Fall armyworm, Green cloverworm, Southern armyworm, Tobacco budworm, Vegetable leafminer.</p>	<p>Use 1/3 fl. oz. in 1 gal. of water. Apply every 5 to 10 days as needed. Can be applied up to 1 day prior to harvest.</p>
<p>Melons-see Cucurbits</p>		
<p>Potatoes</p>	<p>Beet armyworm, Colorado potato beetle, Potato leafhopper, Potato tuberworm, Cabbage looper, Cutworms, European corn borer, Potato psyllid, Tarnished plant bug</p>	<p>Use ¼ fl. oz. in 2 gals. of water. Repeat application only as needed to maintain insect control. Do not make more than 12 applications. Can be applied up to 7 days prior to harvest.</p>
<p>Peppers</p>	<p>Vegetable leafminers, Cabbage looper, Flea beetle, Corn earworm, Pepper weevil, European corn borer, Green cloverworms, Leafhoppers</p>	<p>Use ½ fl. oz. in 1 gal. of water. Repeat application only as needed to maintain insect control. Do not make more than 8 applications per season. Can be applied up to 3 days prior to harvest.</p>
<p>Strawberries, Raspberries, Blueberries</p>	<p>Strawberry weevils, Aphids, Spider mites, Lygus bugs, Flea beetles, Omnivorous leaftier, Spittlebugs, Japanese beetles, Field crickets, Red Spider, Leafroller, weevils</p>	<p>Use ½ fl. oz. in 1 gal. of water. Allow 14 days between last application and harvest. Repeat application only as needed to maintain insect control. Do not make more than 8 foliar applications per growing season.</p>
<p>Sweet Corn</p>	<p>Armyworms(Fall, Southern), Corn earworms, European corn borers</p>	<p>Use ¼ fl. oz. in 2 gals. of water. Apply every 5 days, but not more than 6 applications per season. Can be applied up to 1 day prior to harvest.</p>

Spray infested areas and areas likely to be infested, such as wood surfaces, voids and channels in damaged wood, spaces between wooden parts of a structure, and between wood and foundations where infestation is likely to occur, also around fenceposts, wood piles (do not use on firewood) and tree stumps.

Suitable protective clothing should be worn in confined or overhead areas to prevent exposure to eyes and skin. The use of eye protection, protective head covering, chemical resistant gloves and footwear, a long-sleeved shirt and long pants or coveralls are recommended. Do not permit contact with treated surfaces until spray has dried. Treated confined areas, such as detached garages, under porches and decks, should be ventilated during and immediately following application before the areas are re-occupied.

**For carpenter ants**, spray around doors and windows and other places where ants may enter or crawl and hide. Also treat cracks and crevices or openings into wall voids or other areas where these ants may be present. **For carpenter bees**, spray or inject into holes and tunnels already made by the bees with a tank sprayer, and spray areas where they are beginning to build tunnels into the wood. **For Termites\* [including dry-wood termites]**: treat as for other wood-infesting insects. This product will kill workers and winged reproductive forms of termites in the treated areas.

\*The purpose of such applications of this product for termites is to kill workers or winged reproductive forms which may be present in treated channels at the time of treatment. Such applications are not a substitute for mechanical alteration, soil treatment, or foundation treatment but are merely a supplement. This product is not recommended as sole protection against termites. For active indoor termite infestations, get a professional inspection.

**Soil Termite Treatment Directions**

**Subterranean Termites:** DO NOT USE INSIDE DWELLINGS. For use around existing structures, not for pre-construction use. For post-construction treatment around residential buildings. **Mix 6.7 oz of this product per 1 gallon of water** (makes 0.5% water emulsion). Apply as specified below. One treatment should last up to 5 years if treated soil is left undisturbed. Kills and controls subterranean termites. Chemicals for soil treatment are used to establish a barrier against termites. The chemical emulsion must be adequately dispersed in the soil to provide a barrier between the wood in the structure and the termite colonies in the soil. Barriers may be established in the soil on the outside perimeter of the house by trenching using a rate of 4 gallons of emulsion per 10 linear feet per foot of depth. A complete treatment may require treatment with another EPA registered product to the inside perimeter of the foundation and to other crucial areas. For these situations, contact a licensed pest control operator for treatment. Do not use under slabs, in crawl spaces, or indoors. If termites are found within the house or structure, contact a licensed pest control operator for treatment.

[BACK/SIDE LABEL PANEL]

Suitable protective clothing should be worn in confined or overhead areas to prevent exposure to eyes and skin. The use of eye protection, protective head covering, chemical resistant gloves and footwear, a long-sleeved shirt and long pants or coveralls are recommended. Do not permit contact with treated surfaces until spray has dried. Treated confined areas, such as detached garages, under porches and decks, should be ventilated during and immediately following application before the areas are re-occupied.

Remove all wood debris and wood forms. Around piers, pipes, chimney bases and along foundations of poured concrete, dig a trench 6 inches wide and 4 inches deep. For brick and block foundations, dig the trench about 12 inches deep. If the footing is more than 12 inches deep, make holes with a crowbar, pipe or rod about 1 foot apart that extend from the trench bottom to the top of the footing. Make holes closer in hard-packed clay soils. NEVER DIG BELOW THE TOP OF THE FOOTING. Apply the prepared emulsion (or solution) in the trench and to the backfill at the rate of 4 gallons per 10 linear feet per foot of depth. Cover the treated soil with a thin layer of untreated soil.

For raised porches, terraces and entrance slabs outdoors, drill holes at one foot intervals, 6 inches from the foundation, and pour one-half gallon of emulsion (or solution) into each hole. Refill holes.

Be sure to treat the interiors of accessible hollow block foundations at the same rate as above. Do not use in structural voids insulated with rigid foam. Be careful to have adequate ventilation when treating areas under structure. For severe infestations or treatment under slabs with heat ducts or radiating heat pipes, in basements and buildings with insufficient crawl space, etc., contact a licensed pest control operator or termite control service.

NOTE: Avoid contamination of public and private water supplies by following these precautions. Do not allow a hose or any type of faucet extension to reach into the termite chemical solution while filling or mixing this material with water. Use of anti-back flow equipment will help ensure that the user of this chemical will not contaminate domestic water supplies. Refer to FHA (Federal Housing Administration), state and local specifications for safe distances of treatment areas from wells. Do not treat structures that have cisterns, wells, French drains, or sump systems. Soil in the vicinity of wells, cisterns or ponds should not be treated. For those situations, contact the local Cooperative Extension Service or consult a commercial pest control service. Do not treat soil that is water saturated or frozen or in any conditions where runoff or movement from the treatment area (site) is likely to occur.

**When to reapply:** Reapplication for subterranean termites should only be made when there has been a disruption of the chemical barrier in the soil due to a construction, excavation, landscaping, etc. Reapplication should be made as a spot treatment to those areas. Annual reapplication of the entire premises must be avoided.

[BACK/SIDE LABEL PANEL]



**ANIMALS AND ANIMAL PREMISE USE  
CATTLE BARNs, SWINE HOUSES, MILK ROOMS, KENNELS, POULTRY  
HOUSES, HORSE BARNs, DAIRIES, MILKING PARLORS, FEED LOTS,  
STABLES, PADDOCKS.**

For initial clean up of severe insect infestation, dilute at a rate of (1) part concentrate in 19 parts water (6.7 ounces per gallon) (0.5%). For normal infestations dilute (1) part concentrate in 100 parts water (1.33 ounces per gallon of water) (1 1/3 tablespoonful per gallon) (0.1%).

**REFERENCE CHART**

PESTS	DILUTIONS (Mix Well)	HOW TO APPLY
<b>PESTS ON FARM PREMISES</b>		
<b>(Barns, Dairies, Loafing Sheds, Milking Parlors, Feedlots, Stables, Paddocks, Poultry and Livestock Housing)</b>		
Horseflies, Stableflies, lesser Houseflies and other manure breeding flies, Fleas, Lice, Mites, Ticks, including Deer tick (carrier of Lyme disease). Aids in control of Cockroaches, Mosquitoes and Spiders.	Dilute 8ozs. in 6.25 gals. water Dilute 1pt. in 12.5 gals. water Dilute 1qt. in 25 gals. water	Spray walls and surfaces thoroughly but do not allow runoff to occur; about 1gal. per 750-1000 sq.ft. Treat no more often than once every 2 weeks. Do not contaminate Animals feed or water by spray. Do not use in milk rooms
<b>PESTS ON DOG PREMISES (Kennels, Dog Houses, Runs and Yards)</b>		
Fleas, Flies, Ticks, Lice and Mange Mites (adult and larvae)	Dilute 8ozs. in 6.25 gals. water Dilute 1pt. in 12.5 gals. water Dilute 1qt. in 25 gals. water	Thoroughly spray infested areas, pet beds and resting quarters. Fresh bedding should be placed in animals quarters following treatment. Most efficient pest control is accomplished when heavy pressure and good contact is achieved. Timing and frequency of applications should be based on pest populations reaching nuisance levels, but accompanying manure removal and sanitation practices should precede sprays. Do not spray feed, food or water. Repeat as needed but not more than once every two weeks.

[BACK/SIDE LABEL PANEL]

**PESTS IN CATTLE BARNs, HORSE BARNs, SWINE HOUSES, KENNELS, MILKING PARLORS, MILK ROOMS, DAIRIES, POULTRY HOUSES, FEEDLOTS STABLES AND PADDOCKS**

Flies, Mosquitoes and Gnats	Dilute 8ozs. in 6.25 gals. water Dilute 1pt. in 12.5 gals. water Dilute 1qt. in 25 gals. water	FOG OR FINE MIST: Apply, directing the spray toward the ceiling and upper corners until the area is filled with mist. Use about 2 ounces per 1000 cubic feet or space. For best results, close doors and windows before spraying and keep closed for 10-15 minutes. Remove animals prior to treatment. ventilate before reoccupying.
-----------------------------	--	--

**PESTS ON LARGE ANIMALS (Dairy or Beef Cattle, Horses, Sheep and Goats)**

Faceflies, Hornflies, Stableflies, Mosquitoes, Lice, Mites, Ticks, including Deer tick (carrier of Lyme disease)	Dilute 8ozs. in 50 gals. water Dilute 1pt. in 100 gals. water Dilute 1qt. in 200 gals. water	Spray to thoroughly cover entire animal. For lice or mites a second treatment is recommended 14-21 days later. Spray lactating dairy cows on after milking is completed.
Hornflies, Faceflies, Stableflies, Ear Ticks	Dilute 2ozs. in 1gal. water Dilute 1pt. in 2.5 gals. water	SPOT TREATMENT - LOW PRESSURE SPRAY Apply 1/2oz. per ear or 2-4oz. on face or 12-16oz. along the backline.

**PESTS ON SWINE AND POULTRY (Do not ship swine for slaughter within 5 days of last treatment)**

Mange Mites	Dilute 8ozs. in 25 gals. water Dilute 1pt. in 50 gals. water Dilute 1qt. in 100 gals. water	Spray, dip or sponge animals. Retreat after 14 days, spraying walls and floor space and bedding to kill later hatching, developing stages.
Blowflies, Flies, Mosquitoes, Hog Lice, Fleas, Ticks, including Deer tick (carrier of Lyme disease)	Dilute 8ozs. in 25 gals. water Dilute 1pt. in 50 gals. water Dilute 1qt. in 100 gals. water	Spray, dip or sponge to apply 1 pint per pig, especially around ears.
Poultry Mites, Northern Fowl Mites & Lice	Dilute 8ozs. in 12.5 gals. water Dilute 1pt. in 25 gals. water Dilute 1qt. in 50 gals. water	Spray at the rate of 1/2oz. per bird, or 1 gal per 100 birds, with a fine mist. Spray roosts, walls and nests or cages.

[BACK/SIDE LABEL PANEL]

**PESTS ON DOGS (DO NOT TREAT PUPPIES LESS THAN TWELVE (12) WEEKS OF AGE)**

Fleas, Ticks, including  
Deer tick (carrier of Lyme  
disease), Lice & Mange

Dilute 8ozs. in 12.5 gals. water  
Dilute 1pt. in 25 gals. water  
Dilute 1qt. in 50 gals. water

Thoroughly wet the animal by dipping, sponging or spraying. Allow animal to dry in a warm place without rinsing or toweling. Do not reapply product for two to three weeks. Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals or animals on medication. **Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.**

**INDOOR USE AREAS**

**HOMES, AND THE NONFOOD/FEED AREAS OF BAKERIES, BEVERAGE PLANTS, CANNERIES, FLOUR MILLS, FOOD PROCESSING PLANTS, GRAIN ELEVATORS, GRAINARIES, HOSPITALS (non-occupied areas), HOTELS, MOTELS, INDUSTRIAL INSTALLATIONS, KENNELS, MEAT PACKING PLANTS, OFFICE BUILDINGS, RAILROAD CARS, RESTAURANTS, SCHOOLS, SHIPSHOLDS, SUPERMARKETS, TRUCK TRAILERS, WAREHOUSES.**

Do not use in food/feed handling establishments, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared food is served such as dining rooms but excluding areas where food may be prepared or held. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food/feed should be covered or removed.

Non-food/feed areas of food/feed manufacturing, processing, and handling establishments are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

**Do not apply to classrooms when in use. Do not apply product in occupied patient rooms or any rooms while occupied by the elderly or infirm.**

[BACK/SIDE LABEL PANEL]

**Not for use in Federally Inspected Meat and Poultry Plants.**

**As a surface spray to control cockroaches, waterbugs, palmetto bugs, ants, silverfish, beetles, mites, weevils:** Use a good sprayer and adjust to deliver a coarse, wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, and behind shelves and drawers. Spray bookcases for silverfish. Spray ant trails, nests and points of entry. If surface application only is to be used, spray floors, walls and other surfaces, applying at a rate of (1) gallon to 750 square feet of surface.

**To control fleas and ticks (adult and larvae),** thoroughly spray infested areas, pets beds, resting quarters, nearby cracks and crevices, along and behind baseboards, molding, window and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Apply with mechanical or compressed air equipment (non-thermal), adjusted to deliver a fine mist. Close doors and windows and shut off ventilating systems.

**For rapid control of house flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, bees, black flies, angoumois grain moths and tobacco moths, SPACE SPRAY -** direct spray at an upward angle, disturbing it uniformly through the entire area at a rate of (1) ounce per 1000 cubic feet of space. Keep area closed for at least 10 minutes. Vacate area after treatment and ventilate and sweep up dead insects before occupying.

**For rapid kill of exposed or accessible stages of other insects named on this insert:** Apply using conventional mechanical or compressed air equipment (non-thermal), following directions for space spraying in the paragraph above. Apply the above dilutions at a rate of (1) ounce per 1000 cubic feet of space.

**STORAGE AND DISPOSAL**

- Prohibitions:** Do not contaminate water, food or feed by storage or disposal.
- Pesticide Storage and Spill Procedures:** Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.
- Pesticide Disposal:** Pesticide spray mixture, or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.
- Container Disposal:** Triple rinse (or equivalent), then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved State and local procedures.
- Containers One Gallon and Smaller:** Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

**[BACK/SIDE LABEL PANEL]**

**NOTE: Buyer assumes all risks of use, storage or handling of this product not in strict accordance with label DIRECTIONS and PRECAUTIONS. [Optional Statement]**

**NOTE: CONTROL SOLUTIONS, INC. WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL AND IS REASONABLY FIT FOR THE PURPOSES STATED ON THE LABEL WHEN USED IN ACCORDANCE WITH THE DIRECTIONS FOR USE. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF CONTROL SOLUTIONS, INC. IN NO CASE SHALL CONTROL SOLUTIONS, INC. BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.**

**EXCEPT AS EXPRESSLY PROVIDED HEREIN, CONTROL SOLUTIONS, INC. MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESSED OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND CONTROL SOLUTIONS, INC. TOTAL LIABILITY, SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.**

**CONTROL SOLUTIONS, INC.  
5903 GENOA-RED BLUFF  
PASADENA, TEXAS 77507-1041  
281-892-2500**

**EPA Registration No. 53883-72  
EPA Establishment No. 53883-TX-002**