

53883-78

8/22/2013

117

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460-0001



OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

8-22-13

Anne Turnbough
Control Solutions, Inc.
5903 Genoa-Red Bluff
Pasadena, TX 77507

Subject: Amended Reregistration Label
Product Name: Permethrin 3.2. EW
EPA Registration Number: 53883-78
EPA Decision Numbers: 413291

Dear Dr. Turnbough:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all of the information submitted with your application to support the reregistration of the above referenced product in connection with the permethrin RED, and has concluded that your submission is acceptable.

NOTE: This product is not being reregistered under sections 3(c)5 and 4(g) of FIFRA at this time.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on non-refillable containers. The code may appear either on the label or durably marked on the container itself, and can be added by non-notification per PRN 98-10.

Please note that the record for this product currently contains the Confidential Statements of Formulation (CSFs) listed below. Any previously dated CSFs are superseded.

- Basic CSF, dated 8/6/02

A copy of your label stamped "Accepted" is enclosed along with copies of the acute toxicity and product chemistry reviews completed for the subject product. Products shipped after 12 months from the date of this amendment or the next printing of the label whichever

occurs first, must bear the new revised label. Your release for shipment of the product bearing the amended label constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e).

If you have any questions about this letter, please contact Dr. Jennifer Urbanski at 703-347-0156 or urbanski.jennifer@epa.gov.

Sincerely,



Richard J. Gebken
Product Manager (10)
Insecticide Branch
Registration Division (7504P)

Enclosures: Label stamped "Accepted"
Acute Toxicity Review, dated 9/27/12
Product Chemistry Review, dated 8/4/11

3/17

Permethrin

3.2 E.W.

[ABN: Martin's Flea Tick & Mange Dip]

For Residential Use Only

[Optional proposed text appears in brackets - the final label may include some or all of the optional text]

- [Kills Termites*[including dry-wood termites] (for localized control only)]
- [For Use around foundations, outside buildings, porches, wood piles, wooden fences and decks, patios]
- [Kills troublesome listed outdoor insects which destroy Lawns, Roses, Flowers, Trees and Shrubs, Vegetables, Fruits, and Nuts]
- [Provides contact and residual control for up to 4 weeks] [on dogs and dog premises]
- [Controls: Flies, Fleas, Lice, Ticks][on dogs and dog premises]
- [KILLS OVER 100 DIFFERENT LISTED INSECTS]
- [Kills Fleas, Flies, Lice, Mange Mites and Ticks, including Deer Tick (carrier of Lyme Disease)] [on dogs and dog premises]
- [Indoor/Outdoor Insecticide]
- [For Use on Dogs and Dog Premises for Control of Fleas and Ticks]
- [Water Based]
- [Makes 8 Gallons] of Dip Solution] [for use on Dogs]] (16 oz size only)
- [Protects your Lawns, Trees & Shrubs, Roses & Flowers, Fruits & Nuts, and Vegetables from: Alfalfa loopers, Ants, Aphids, Armyworms, Bagworms, Beetles, Borers, Brown dog ticks, Budworms, Cabbageworms, Chinch bugs, Cicadas, Citrus black flies, Cluster flies, Codling moths, Corn earworms, Crickets, Cutworms, Earwigs, Elm spanworms, Fall cankerworms, Fire Ants, Fleas, Green fruitworms, Gnats, Grasshoppers, Houseflies, Inchworms, Japanese beetles, Leafhoppers, Leafminers, Leafrollers, Lone Star ticks, Lygus bugs, Mealybugs, Millipedes, Mole crickets, Mosquitoes, Moths, Navel orangeworms, Scales, Sod webworms, Sowbugs (pillbugs), Spider mites, Spring cankerworms, Tent caterpillars, Thrips, Tobacco budworms, Weevils and Whiteflies.

ACTIVE INGREDIENT:

Permethrin 3.20%

OTHER INGREDIENTS: 96.80%

TOTAL: 100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Back/Side Label Panel For Additional Precautionary Statements

Net Contents _____

ACCEPTED
8-22-13
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 53883-78

4/17

[BACK/SIDE LABEL PANEL]

- **READ ENTIRE LABEL BEFORE USE**
- **FOR PET USE - USE ONLY ON DOGS**

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear long-sleeved short and long pants, socks, shoes, and chemical-resistant gloves.

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
<p>NOTE TO PHYSICIAN AND VETERINARIANS: Do not administer or instill milk, cream or other substances containing vegetable or animal fats which will enhance absorption of lipophilic substances.</p> <p>"Have the product container or label with you when calling a poison control center or doctor, or going for treatment. <u>For information on this product, contact the National Pesticide Information Center, 1-800-858-7378, Monday-Friday, 7:30 AM-3:30 PM PST. You may also contact the National Poison Control Center, 1-800-222-1222, day or night, for emergency medical treatment information or SafetyCall® at (866) 897-8050. FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.</u>"</p>	

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish. Do not apply directly to water. Avoid contamination of ornamental fish ponds. Drifts and runoff from treated areas may be hazardous to aquatic organisms in treated areas. Do not contaminate water when cleaning equipment or disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively, visiting the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame. Do not use in undiluted form.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE USE

USE RESTRICTIONS

- Not for use on plants being grown for sale or other commercial seed production, or for research purposes.
- Do not allow adults, children, or pets to enter the treated area until sprays have dried.
- Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.
- All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:
 - Treatment to soil or vegetation around structures;
 - Applications to lawns, turf, and other vegetation;
 - Applications to building foundations, up to a maximum height of 3 feet.
- Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications, only
- Do not apply this product that will contact adults, children, or pets, either directly or through drift.
- Do not water the treated area to the point of run-off. Do not make applications during rain.
- Not for broadcast use on indoor residential surfaces.
- Not for use in outdoor residential misting systems.
- **DO NOT USE ON CATS.** May be toxic or potentially fatal if applied to or ingested by cats. Accidental application to cats and/or grooming a recently treated dog may result in tremors and/or uncoordinated muscle movement. If this occurs, immediate veterinary care should be provided.

Keep people and pets off treated areas until the spray has dried.

SHAKE WELL BEFORE USING

[BACK/SIDE LABEL PANEL]

THREE STEPS TO MIX AND APPLY:

1. For lawns and vegetables, measure the area to be sprayed. Calculate square feet by multiplying length by width. For all other applications, apply as a thorough cover spray.
2. Use a clean sprayer. Carefully measure and mix the amount of concentrate and water as directed.
3. Spray as directed. Thorough coverage is important. Flush sprayer with clean water after each use. Make applications when insects first appear. Repeat applications as directed.

Use Dilution rate as indicated:

1 fl. oz. = 6 tsp. or 2 Tbsp.

8 fl. ozs. = 1/2 pt. or 1 cup

DIRECTIONS FOR APPLYING

HOME LAWNS

Turf	Pests	How to Use
Bentgrass, Bermudagrass, Bluegrass, Dichondra, Fescue, Irish Moss, Merion, St. Augustine Grass	Ants, Armyworms, Brown dog ticks, Chinch bugs, Crickets, Cutworms, Fleas, Grasshoppers Japanese beetle grubs, Mole crickets, Sod webworms, Ticks that may carry Lyme Disease, Leafhoppers	Thoroughly wet down grass a few hours before applying. Home lawns should be no taller than 3 inches at time of application. Slightly more water may be used as long as the amount of product per area is as listed below. For heavy infestations, repeat application after 2 weeks. For ants, crickets and grasshoppers , use 1 1/2 fl. oz. in 3 gals. of water to cover 1,000 sq. ft. For other listed insects , use 4 2/3 fl. ozs. in 10 gals. of water to cover 1,000 sq. ft.

[BACK/SIDE LABEL PANEL]

HOUSE PLANTS

INDOORS: TO PROTECT PLANTS IN RESIDENTIAL GREENHOUSES AND LATHHOUSES AND FOR INDOOR LANDSCAPING:

House Plants	Pests	How to Use
Ageratum, Ardisia, Aster, Baby's Breath, Begonia, Birdsnest Fern, Bleeding Heart, Carnation, Chrysanthemum, Coleus, Crown of Thorns, Cockscomb, Cyclamen, Dracaena, Dumbcane, English Ivy, Fuchsia, Gladiolus, Grape Ivy, Juniper, Lily, Marigold, Orchid, Palm, Pansy, Peperomia, Petunia, Philodendron, Piggy-Back Plant, Poinsettia, Portulaca, Pothos, Prayer Plant, Purple Passion, Rabbits Foot Fern, Rose, Schefflera, Snapdragon, Snake Plant, Statice, Velvet Plant, Verbena, Zinnia	Aphids, Armyworms, Corn earworms, Exposed thrips, Fall cankerworms, Inchworms, Japanese beetles, Leafminers, Leafrollers, Mealybugs, Scales, Solder mites, Spring cankerworms, Whiteflies	Mix ¾ fl. oz. in 1 gal. of water. Contact and residual: Use sufficient spray to obtain full coverage of all foliage. If application is to be made indoors, remove infested plants from living areas prior to application. Use in an area not likely to be occupied by people or pets (particularly fish). After spraying, plants can be returned to original location. Do not use more product than recommended, as this may cause some plant injury. Use with care on ferns, palms and fuchsia. As with all plants listed, especially carnations, chrysanthemums, and roses, there exists a wide variety of species and cultivars, and therefore, a wide variety of susceptibility to injury. To ascertain possible phytotoxicity, treat a few plants as described above, wait 24 hours, and check for possible injury.
Velvet Plants	Fungus Gnats	

Fire Ant Control

Individual Mound and Plant Container Treatment

Mix 4⅔ fl. oz. in 3 gals. of water and apply as a drench using a sprinkler can. Gently apply 1 gallon of the mixture over the surface of each mound. Thoroughly wet mound and surrounding area of 3x4 ft. (12 sq. ft.). For best results apply in cool weather 65⁰-80⁰F. early or late in the day. Treat new mounds as they appear. Pressurized sprays may disturb ants and cause migration which will reduce product effectiveness. **Use the same method as above for the control of Fire Ants that may nest in permanently placed containerized ornamentals plants.**

TREES & SHRUBS (ORNAMENTALS)

Trees & Shrubs	Pests	How to Use
Arizona, Cypress, Azalea, Birch, Non-bearing Citrus, Conifers, Elm, English Ivy, Euonymus, Weeping Fig, Fir, Honeysuckle, Lilac, Mock-orange, Oak, Palm, Philodendron, Pine, Poinsettia, Tulip Poplar, Rhododendron, Taxus	Aphids, Bagworms, Cicadas, Exposed thrips, Fall cankerworms, Inchworms, Leafminers, Leafrollers, Mealybugs, Spider mites, Whiteflies, Gypsy Moth, Tent Caterpillar, Japanese Beetles	Use ¼ fl. oz. in 1 gal. of water. Use sufficient spray to obtain full and uniform coverage. Avoid spraying to the point of runoff. Try to cover underside of leaves and penetrate dense foliage. Spray in the late afternoon or evening, when the temperature ranges from 50 ^o -75 ^o F and when there is little or no wind. Spray at the first sign of insects. Application can be made up to the day of collection.
Conifer	Budworms	
Elm	Elm leaf beetles, Elm spanworms, Fall webworms	
Douglas and Grand Firs	Tussock moths	
Honeysuckle	Fall webworms	
Non-bearing Citrus	Citrus black flies	
Oak	Oakworms	
Pine	Pine beetles, Pine moths, Pine needle miners, Needle scales	

On Tent caterpillars, application should be made when caterpillars are young and tents are first noticed. For best results, apply in the late afternoon or evening, when caterpillars have returned to their tents. Wet tents with spray on all sides. A few caterpillars may be seen escaping from the tent; however, they should die within several hours.

[BACK/SIDE LABEL PANEL]

ROSES & FLOWERS (ORNAMENTALS)

Roses & Flowers (Ornamentals)	Pests	How to Use
<p>Ageratum, Aster, Ardisia, Azalea, Baby's breath, Begonia, Birds-nest fern, Bleeding heart, Carnation, Chrysanthemum, Coleus, Common Ninebark and Snowberry, Crown of thorns, Cockscomb, Cyclamen, Dracaena, Dumbcane, English Ivy, Exacum, Fuchsia, Gladiolus, Grape Ivy, Gold bells, Hypoestes, Ivy, Juniper, Lilac, Marigold, Mock-Orange, Nannyberry, Orchid, Palm, Pansy, Pea shrub, Peperomia, Petunia, Philodendron, Piggy-back Plant, Poinsettia, Portulaca, Pothos, Prayer Plant, Purple Passion, Rabbits foot fern, Rhododendron, Rose, Schefflera, Snapdragon, Snake Plant, Statice Velvet Plant, Verbena, Zinnia</p>	<p>Aphids, Armyworms, Bagworms, Cabbage worms, Cicadas, Cornworms, Cutworms, Exposed thrips, Fall cankerworms, Fall webworms, Fungus gnat, Inchworms, Japanese beetles, Leafminers, Leafrollers, Mealybugs, Scale crawlers, Spring cankerworms, Spider mites, Whiteflies, Rose Chafer, Psyllids, Scale Insects</p>	<p>Use ¾ fl. oz. in 1 gal. of water. Use sufficient spray to obtain full and uniform coverage. Avoid spraying to the point of runoff. Try to cover underside of leaves and penetrate dense foliage. Spray in the later afternoon or evening when the temperature ranges from 50^o-75^oF and when there is little or no wind. Spray at the first sign of insects. Application can be made up to the day of collection.</p>

FRUITS & NUTS TREES

USE DIRECTIONS

Use 1½ fl. oz. in 1 gal. of water. Apply when insects appear. Use sufficient spray to obtain full and uniform coverage. Avoid spraying to the point of runoff. Try to cover underside of leaves and penetrate dense foliage.

Fruits & Nuts Trees	Pests	How to Use
<p>Almonds, Pecans</p>	<p>Navel orangeworms, Peach twig borers</p>	<p>Do not make more than 2 applications during hull split and 5 applications per season. Can be applied up to 7 days prior to harvest.</p>

Apples	Green fruitworms, Aphids, Leafrollers(Oblique-banded, Redbanded), Plum curculio, Rosy apple aphids, Spotted tentiform leafminers, Tarnished plant bugs, White apple leafhoppers, Japanese Beetles	Do not make more than 3 applications. Do not apply after petal fall.
Peaches	Borers(Lesser peachtree, Peach twig), Green fruitworms, Plum curculio, Oriental fruit moths, Tarnished plant bugs	Do not apply within 7 days of harvest. Do not make more than 8 applications per season.
Pears (dormant through delayed dormant)	Pear psylla	Apply during the dormant through delayed dormant growth period only. Do not make more than 2 applications per season.
Pears (summer)	Codling Moths, Green fruitworms, Pear psylla, Aphids	Do not make more than 3 applications per season. Can be applied up to 14 days prior to harvest.

VEGETABLES

Prepare a spray solution at the dilution rates listed for each crop in the chart below. Apply the spray solution at the rate of 1 gallon to cover 440 square feet surface area.

Vegetables	Pests	How to Use
Asparagus (pre-harvest)	Asparagus beetles, Cutworms	Use 2½ fl. ozs. in 2 gals. of water. Apply to ensure uniform coverage. Do not apply more than 4 applications per crop. Can be applied up to 3 days prior to harvest.
Asparagus (post-harvest)	Larval stages of asparagus beetle; Tarnished plant bugs, Lygus bugs, Adult Japanese beetle	Use 2½ fl. ozs. in 1 gal. of water. Treat ferns or bush growth after spear harvest when insects are present.
Celery-see Leafy Vegetables		

<p>Cole Crops, Broccoli, Brussels Sprouts, Cabbage, Cauliflower</p>	<p>Beet armyworms, Cabbage aphids and loopers, Diamondback moths, Imported cabbageworms</p>	<p>Use ¾ fl. oz. in 1 gal. of water. Apply every 5 days or as needed only to maintain insect control. Can be applied up to 1 day prior to harvest. Do not make more than 8 applications on Cauliflower, Brussels Sprouts and Broccoli and 10 applications on Cabbage.</p>
<p>Corn-see Sweet Corn</p>		
<p>Cucumber, Melons, Pumpkins, Squash</p>	<p>Flea beetle, Leafhopper</p>	<p>Use 1½ fl. ozs. in 1 gal. water. Apply as needed, but not more than 8 applications per season. Can be applied up to 3 days prior to harvest.</p>
<p><u>Cucurbits:</u> Balsam pear(bitter melon), Chinese waxgourd(Chinese preserving melon), citron melon, cucumber, gherkin, edible gourds; melons including hybrids such as cantaloupe, casaba, crenshaw, honeydew, honey balls, mango melon, muskmelon, Persian melon, pumpkin, summer & winter squash, watermelon</p>	<p>Aphids, Cabbage looper, Cucumber Beetle (adults), Cutworms, Leafhoppers, Leafminers, Melonworms, Pickleworm, Plant bugs, Squash bugs.</p>	<p>Use 1½ fl. ozs. in 1 gal. of water. Apply to ensure uniform coverage. Can be applied up to the day of harvest.</p>
<p>Eggplant</p>	<p>Colorado potato beetles, Vegetable leafminers</p>	<p>Use 2½ fl. ozs. in 1 gal. of water. Apply to ensure uniform coverage. Do not exceed 16 applications. Can be applied up to 3 days prior to harvest.</p>
<p>Horseradish (Foliar application)</p>	<p>Imported crucifer weevils</p>	<p>Use 2½ fl. ozs. in 1 gal. water. Do not exceed 3 foliar applications. Can be applied up to 22 days prior to harvest.</p>

<p>Head Lettuce</p>	<p>Corn earworms, Beet armyworms, Loopers (Alfalfa, Cabbage), Tobacco budworms</p>	<p>Use 1½ fl. ozs. in 1 gal. water. Apply every 5-10 days only as needed to maintain control. Do not make more than 10 applications per season. Can be applied up to 1 day prior to harvest.</p>
<p><u>Leafy Vegetables:</u> Amaranth, Arrugula, Celery, Celtuce, Chervil, Corn salad, Chrysanthemum(edible-leaved and garland), cress(garden and upland), Dandelion, Dock, Endive, Fennel, Lettuce(head and leaf), Okra, Parsley, Purslane(garden and winter), Rhubarb, Spinach, Swiss chard</p>	<p>Alfalfa looper, Aphids, Beet armyworm, Cabbage looper, Corn earworm, Cutworms, European corn borer, Fall armyworm, Green cloverworm, Southern armyworm, Tobacco budworm, Vegetable leafminer.</p>	<p>Use 1½ fl. ozs. in 1 gal. of water. Apply every 5 to 10 days only as needed to maintain control. Can be applied up to 1 day prior to harvest.</p>
<p>Melons-see Cucurbits</p>		
<p>Potatoes</p>	<p>Beet armyworm, Colorado potato beetle, Potato leafhopper, Potato tuberworm, Cabbage looper, Cutworms, European corn borer, Potato psyllid, Tarnished plant bug</p>	<p>Use 2½ fl. ozs. in 2 gals. of water. Apply as needed, but not more than 12 applications. Can be applied up to 7 days prior to harvest.</p>
<p>Peppers</p>	<p>Vegetable leafminers, Cabbage looper, Flea beetle, Corn earworm, Pepper weevil, European corn borer, Green cloverworms, Leafhoppers</p>	<p>Use 1½ fl. ozs. in 1 gal. of water. Do not make more than 8 applications per season. Can be applied up to 3 days prior to harvest.</p>

Spinach	Beet armyworms, Corn earworms, Cabbage loopers, Cutworms, European corn borers, Green cloverworms, Leafhoppers	Use 2½ fl. ozs. in 2 gals. of water. Apply to ensure uniform coverage. Do not make more than 7 applications per season. Can be applied up to 7 days prior to harvest.
Strawberries, Raspberries, Blueberries	Strawberry weevils, Aphids, Spider mites, Lygus bugs, Flea beetles, Omnivorous leafhopper, Spittlebugs, Japanese beetles, Field crickets, Red Spider, Leafroller, weevils	Use 1½ fl. ozs. in 1 gal. of water. Allow 14 days between last application and harvest. Make no more than 8 foliar applications per growing season.
Sweet Corn	Armyworms(Fall, Southern), Corn earworms, European corn borers	Use 2½ fl. ozs. in 2 gals. of water. Apply every 5 days, but not more than 6 applications per season. Can be applied up to 1 day prior to harvest.
Tomatoes (Do not apply on cherrytomatoes and varieties yielding tomatoes less than one inch in diameter.)	Beet and southern armyworm, Tomato pinworm, Tomato fruitworm, Horn worms, Granulate cutworms, Colorado potato beetle, Cabbage looper	Use 1½ fl. ozs. in 1 gal. of water. Apply to ensure uniform coverage. Can be applied up to day of harvest.

NUISANCE PESTS IN OUTDOOR AREAS

USE DIRECTIONS

Use 5¼ fl. ozs. in 1 quart of water, treating 400 linear feet with a 6-inch spray band.

OUTSIDE SURFACES OF BUILDING

Use Permethrin 3.2 E.W. on Ants, Clover mites, Crickets, Earwigs, Millipedes, Sowbugs(pillbugs) and spiders to help prevent infestation of buildings, treat the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Also, apply as a residual spray to outside surfaces of building, including porches, screens, window frames, eaves, patios, garages, refuse dumps and other areas where these pests congregate or have been seen.

OTHER OUTDOOR AREAS

Use Permethrin 3.2 E.W. on Ants, Crickets, Earwigs, Fleas, Ticks (**including ticks that may carry Lyme Disease**) for treatment of localized infestations of these insects in areas where there are weeds or bushy non-crop areas, spray infested areas thoroughly. **For ants**, thoroughly wet hills and runways. Repeat application as reinfestations occur.

[BACK/SIDE LABEL PANEL]

To reduce annoyance from: **Cluster flies, Flying moths, Gnats, House flies, Mosquitoes.** Spray outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages, around shrubbery, and other places where these insects alight or congregate.

WOOD-INFESTING INSECTS (including termites*): For Beetles (Furniture, Deathwatch, False powderpost, Longhorned, Powderpost), Carpenter Ants (and other wood-infesting ants), Carpenter bees and Termites (including dry-wood termites):

Treat outside surfaces of buildings, porches, wooden decks and patios, wooden fences, window frames, doorways, foundations, eaves, patios, garages, garbage areas and other areas where you may find these pests.

Mix 7 $\frac{3}{4}$ fl. ozs. with 1 gallon of water to treat 100 sq.ft. Spray infested surfaces until thoroughly wet, avoiding dripping and runoff. Apply evenly on wood surfaces. To treat small areas, apply by brushing or spraying the diluted spray. For large areas, use a coarse, low pressure spray. Do not spray into electrical outlets or utilities.

Spray infested areas and areas likely to be infested, such as wood surfaces, voids and channels in damaged wood, spaces between wooden parts of a structure, and between wood and foundations where infestation is likely to occur, also around fenceposts, wood piles (do not use on firewood) and tree stumps.

Suitable protective clothing should be worn in confined or overhead areas to prevent exposure to eyes and skin. The use of eye protection, protective head covering, chemical resistant gloves and footwear, a long-sleeved shirt and long pants or coveralls are recommended. Do not permit contact with treated surfaces until spray has dried. Treated confined areas, such as detached garages, under porches and decks, should be ventilated during and immediately following application before the areas are re-occupied.

For carpenter ants, spray around doors and windows and other places where ants may enter or crawl and hide. Also treat cracks and crevices or openings into wall voids or other areas where these ants may be present. **For carpenter bees**, spray or inject into holes and tunnels already made by the bees with a tank sprayer, and spray areas where they are beginning to build tunnels into the wood. **For Termites* [including dry-wood termites]:** treat as for other wood-infesting insects. This product will kill workers and winged reproductive forms of termites in the treated areas.

*The purpose of such applications of Permethrin 3.2 E.W. for termites is to kill workers or winged reproductive forms which may be present in treated channels at the time of treatment. Such applications are not a substitute for mechanical alteration, soil treatment, or foundation treatment but are merely a supplement. This product is not recommended as sole protection against termites. For active indoor termite infestations, get a professional inspection.

[BACK/SIDE LABEL PANEL]

DOG AND DOG PREMISE USE

REFERENCE CHART

PESTS	DILUTIONS (Mix Well)	HOW TO APPLY
PESTS ON DOG PREMISES (Dog Houses, Runs and Yards)		
Fleas, Flies, Ticks, Lice and Mange Mites (adult and larvae)	Dilute 4 fl. ozs. in 1 gal. water Dilute 1 pt. in 4 gals. water Dilute 1 qt. in 8 gals. water Dilute 1 gal. in 32 gals. water	Thoroughly spray infested areas, pet beds and resting quarters. Fresh bedding should be placed in animals quarters following treatment. Do not spray feed, food or water. Repeat as needed but not more than once every two weeks.
PESTS ON DOGS (DO NOT TREAT PUPPIES LESS THAN TWELVE (12) WEEKS OF AGE)		
Fleas, Ticks, including Deer tick (carrier of Lyme disease), Lice & Mange	Dilute 2 fl. ozs. in 1 gal. water Dilute 1 pt. in 8 gals. water Dilute 1 qt. in 16 gals. water Dilute 1 gal. in 64 gals. water	Thoroughly wet the animal by dipping, sponging or spraying. Allow animal to dry in a warm place without rinsing or toweling. Do not reapply product for two to three weeks. Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals or animals on medication. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage and Spill Procedures: Store upright at room temperature. Avoid exposure to extreme temperatures. Store in a locked area out of reach of children and domestic animals. In case of spillage or leakages, soak up with an absorbent material such as sand, sawdust, earth, Fuller's earth, etc. Dispose of with chemical waste.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spilled bait, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance.

Container Disposal: If empty - Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. If recycling is not available: then dispose of container in a sanitary landfill or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

If partly filled: Call your local solid waste agency for disposal instructions. Unless otherwise instructed, place in trash. Never pour unused product down the drain or on the ground.

[Commercial Language]

For Bulk containers: (Refillable Container) Refill this container with pesticides only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the re-filler. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or re-circulate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

[BACK/SIDE LABEL PANEL]

[NOTE: Buyer assumes all risks of use, storage or handling of this product not in strict accordance with label DIRECTIONS and PRECAUTIONS.]

Control Solutions, Inc. ® Warranty

LIMITATION OF WARRANTY AND LIABILITY

Read the entire direction for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following **CONDITIONS, DISCLAIMER OF WARRANTIES, and LIMITATIONS OF LIABILITY.**

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Control Solutions, Inc. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, Control Solutions, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Control Solutions, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Control Solutions, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or

17/17

handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Control Solutions, Inc election, the replacement of product.

^a

CONTROL SOLUTIONS, INC.
5903 GENOA-RED BLUFF
PASADENA, TEXAS 77507-1041
281-892-2500

EPA Registration No. 53883-78
EPA Establishment No. 53883-TX-002

^a Permethrin 3.2 EW (53883-78) revised per EPA RED comments 8-14-2013