53883-162

6/21/2006

age 124



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

JUN 2 1 2006

Mr. Joe Blake Control Solutions, Inc. 5903 Genoa-Red Bluff Pasadena, TX 77507-1041

Dear Mr. Blake:

Subject: Amendment - Revise per copyright infringement litigation action Control XX-X-X Lawn Food with .083% Bifenthrin Insecticide EPA Registration Number 53883-162 Your submission dated April 20, 2006

The amendment referred to above, submitted in connection with registration under with

registration under FIFRA section 3(c)(7)(a), is acceptable and a stamped copy of the label is

enclosed for your records.

Sincerely,

George LaRocca. Product Manager 13 Insecticide Branch Registration Division (7505P)

Control XX-X-X Lawn Food with .083% Bifenthrin Insecticide

ACCEPTED					
JUN 2 1 2006					
Washer the Federal Incodition, Fungicita, and Rodensistic Acc, as emended, for the pesticide registered under					

This is a Double Action product with Lawn Food and Bifenthrin Insecticide Feeds the Lawn and controls common lawn insects such as: Ants (*including Imported Fire Ants*), Mole Crickets, Sod Webworms, Chinch Bugs and other Pests as listed. For Lawns, in Landscape areas, around Residential, Commercial, Institutional, Public, and Industrial Buildings, Parks, Recreational Lawns, and Athletic Fields.

EPA Reg. No. 53883-162

EPA Est. No. 53883-TX-002

Bifenthrin* CAS [83322-82657]	0.083%
OTHER INGREDIENTS:	99.917%
TOTAL:	100.000%

Cis isomers 97% minimum; trans. isomers 3% maximum.

KEEP OUT OF REACH OF CHILDREN CAUTION

See other panels for Additional Precautionary Statements.

	FIRST AID	
Have the prod	uct container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall® (866) 897-8050 for emergency medical treatment.	
If swallowed	 Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person. 	
If in eyes		
Note to Physic	an: This product is a pyrethroid. If large amounts have been ingested, milk, cream, and other digestible	

fats and oils may increase absorption and so should be avoided.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift and runoff from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply more than 484 lbs. of this product per acre, per year.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial greenhouses, nurseries, commercial seed production or research purposes.

Restrictions concerning use near waters.

Do not apply this product within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes, or natural ponds, estuaries, and commercial fish ponds.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store product in its original container in a cool, dry, locked place out of the reach of children.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

COMMERCIAL CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

HOMEOWNER CONTAINER DISPOSAL: <u>IF EMPTY</u>: Fold bag and discard in trash. <u>IF PARTLY FILLED</u>: Use as directed or call your local waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any drain.

The contents of this 14 lb. bag feeds & treats up to 5,000 sq.ft. The contents of this 28 lb. bag feeds & treats up to 10,000 sq.ft. The contents of this 50 lb. bag feeds & treats up to 15,000 sq.ft.

LAWN AND PERIMETER APPLICATIONS including GOLF COURSE AREAS

Lawn Applications: A Broadcast application is recommended to ensure uniform coverage over the treatment area. Examples of broadcast equipment is listed below.

Perimeter Applications: Control xx-x-x Lawn food with .083% Bifenthrin Insecticide is intended to prevent and reduce insect entry to homes and businesses. When using this product as directed, apply a <u>6 to 10 feet wide band</u> to the lawn areas around and adjacent to the structure. Observe and apply the following application rates.

Use these rates to control the following insects.					
Apply 2.75 lbs. per 1,000 sq.ft. Apply 5.55 lbs. per 1,000 sq.ft.					
Annual Bluegrass	**Ants,				
Weevils (Hyperodes Adult),	**Centipedes,				
Armyworms,	*Fire Ants,				
Billbugs (Adult),	**Fleas,				
Black Turfgrass	Leafhoppers,				
Ataenius (Adult),	**Millipedes,				
Cutworms,	**Mole Crickets,				
Mealybugs,	**Chinch Bugs,				
Sod Webworms	Ticks and Chiggers				

Control XX-X-X Lawn Food with .083% Bifenthrin Insecticide should be applied when insects first appear. Application of this product should be applied evenly and with a properly calibrated drop or rotary spreader over the entire turfgrass area at a rate of 2.77 to 5.55 lbs. per thousand square feet. Best results will be obtained if treated area is thoroughly irrigated with water immediately after application.

Control XX-X-X Lawn Food with .083 Bifenthrin Insecticide will provide optimum control of Imported Fire Ants if combined with a mound treatment. A broadcast application of the area to be treated will control foraging worker ants and newly mated queens. In addition a mound treatment at the same time will aid in the destruction of the existing colony. The highest rate noted on the application table may be required to control Imported Fire Ants as well as Chiggers, Centipedes, Chinch Bugs, Fleas, Millipedes and other foraging ants.

OPTIONAL TEXT WHEN MARKETED IN NEW YORK STATE

In New York State this product may NOT be applied to any grass or turf <u>within 100 feet of a water body</u>; (lake, pond, river, <u>stream, wetlands, or drainage ditch</u>). In New York State do make a single repeat application of this product if there are signs of renewed insect activity, but not sooner than 2 weeks after the first application.

Spreader Setting Suggestions

424

Varied conditions will affect the application rates of spreaders; therefore the settings listed below are approximate. Age and conditions of spreader and walking speed can cause variation in application rate. Calibrate your spreader with each application. Close spreader hopper and adjust the setting using the suggested settings chart below. Fill spreader on a hard surface to eliminate the chance of spillage on the lawn. Close spreader hopper when stopping or turning. For the 5.55 lbs. rate of product per 1,000 sq. ft. make a second application at right angles to the first 2.75 lb. rate or use the spreader setting for that rate as indicated below.

APPLICATION RATES PER 1,000 SQ.FT.				
DROP SPREADERS	2.75 lbs. RATE	5.55 lbs. RATE		
Yard Star 75350	XX	XX		
Republic EZ 7	XX	XX		
Precision FH22	XX	XX		
Scotts Acugreen	XX	XX		
BROADCAST SPREADERS	· · · · · · · · · · · · · · · · · · ·	······································		
Yard Star 75450	XX	XX		
Precision SB50P	XX	XX		
Republic EZGROW 75450	XX	XX		
Scotts Easygreen 745	XX	. XX		
Scotts Speedygreen 740	XX	XX		

GUARANTEED FERTILIZER ANALYSIS Total Nitrogen (N) - - - - - Variable XX% variable XX% Ammonical Nitrogen variable XX% Water Soluble Nitrogen variable XX% Water Insoluble Nitrogen Available Phosphate (P₂O₅) Variable XX%

Available Potash (K₂O)

(.

Plant Nutrients derived from Diammonium Phosphate, Urea formaldehyde, Urea, Sulfate of Ammonia, Sulfur coated Urea, Sulfate of Potash, Magnesium, Statement of Secondary Plant Nutrients.

Variable XX%

NOTICE: Buyer and user assume all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

Control Solutions, Inc. 5903 Genoa-Red Bluff Road Pasadena, TX 77507-1041

Net Weight 14 pounds 28 pounds 50 pounds