



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

November 6th, 2025

Lisa Adamson
Regulatory Manager
Control Solutions, Inc.
5903 Genoa-Red Bluff
Pasadena, TX 77507-1041

Subject: Label Amendment - Registration Review Mitigation for Oryzalin
Product Name: ORYZALIN PRO 4
EPA Registration Number: 53883-168
Case Number: 476072
Application Dates: February 7, 2020

Dear Lisa Adamson:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Oryzalin Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must submit one copy of the final printed labeling before you release the product for

shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Caleb Carr by phone at 202-566-0636, or via email at carr.caleb@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read 'M. K. Muhammad-Perch', with a long, sweeping horizontal line extending to the right.

Maryam K. Muhammad-Perch, Team Lead
Risk Management and Implementation Branch 4
Pesticide Re-Evaluation Division
Office of Pesticide Programs

ENCLOSURE: Stamped label

ACCEPTED
11/06/2025

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 53883-168

ORYZALIN	GROUP	3	HERBICIDE
----------	-------	---	-----------

Oryzalin Pro 4

A selective preemergence surface-applied herbicide for control of annual grasses and many broadleaf weeds in:

- Landscape Ornamentals
- Container Grown Ornamentals
- Field Grown Ornamentals
- Drainage Areas Under Shadehouse Benches
- Ornamental Bulbs
- Ground Covers/Perennials
- Christmas Tree Plantations

- Non-bearing fruit and nut trees and nonbearing vineyards
- Noncropland and Industrial Sites
- Established Wann Season Turf (including Bahiagrass, Bennudagrass, Buffalograss, Centipedegrass, St. Augustinegrass and Zoysiagrass)
- Tall Fescue (warm season areas)

Active Ingredient:

Oryzalin: 3,5-dinitro-N⁴,N⁴-Dipropylsulfanilamide. 41.0%
Other Ingredients: 59.0%
TOTAL: 100.0%

EPA Reg. No. 53883-168

EPA Est No. 53883-TX-002

Net Contents _____

Contains 4 pounds of active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN
CAUTION

Manufactured for:
Control Solutions, Inc.
5903 Genoa Red Bluff
Pasadena, TX 77507

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
CAUTION

Personal Protective Equipment (PPE)
Applicators and other handlers must wear:
- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Mixers and loaders must wear a chemical-resistant apron in addition to other PPE.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Cover or incorporate spills.

Groundwater Advisory: Oryzalin is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

NON-TARGET ORGANISM ADVISORY: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirement in this box only applies to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. Exception: If the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allow workers to enter the treated area if there will be no contact with anything that has been treated.

Workers may enter treated areas without required PPE during the reentry interval following 1/2 to 1 inch of rainfall or irrigation, if they are performing tasks that do not involve contact with the soil subsurface; otherwise, PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Entry Restrictions for Non-WPS Uses: Keep all persons, children and pets out of treated area until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container only. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

MANDATORY SPRAY DRIFT MANAGEMENT

Ground Boom Applications:

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to use a medium or coarser droplet size according to the most current version of the American Society of Agricultural & Biological Engineers Standard (ASAE S572).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

Boom-less Ground Applications:

- Applicators are required to use a medium or coarser droplet size according to the most current version of the American Society of Agricultural & Biological Engineers Standard (ASAE S572) for all applications.
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure – Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce the effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and move

laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

WEED RESISTANCE MANAGEMENT

For resistance management, Oryzalin Pro 4 is a Group 3 herbicide. Any weed population may contain or develop plants naturally resistant to Oryzalin Pro 4 and other Group 3 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of Oryzalin Pro 4 or other Group 3 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact your local Control Solutions Inc. distributor or sales representative.

PRODUCT INFORMATION

Oryzalin Pro 4 herbicide is a preemergence surface-applied product for the control of many annual grasses and broadleaf weeds in ornamental plantings, bulbs, ground covers/perennials,

established warm-season turfgrass, Christmas tree plantations, non-bearing trees and vines, and noncropland and industrial sites.

Oryzalin Pro 4 is orange in color and may cause temporary discoloration of sprayed surfaces. If this discoloration is undesirable, it may be altered by using a commercially available colorant such as Blazon or removed by spraying surface with water or washing with an industrial cleaner immediately after application. Oryzalin Pro 4 may also be applied with mulch colorants, such as Mulch Magic or Nu-Mulch.

Treatment of Plant Species Not Listed on the Label for Oryzalin Pro 4

Users who wish to use Oryzalin Pro 4 on plant species not recommended on this label may determine tie suitability for use by treating a small number of such plants at a recommended rate. Prior to treatment of larger areas, the treated plants should be observed for any sign of herbicidal injury during 30-60 days of normal growing conditions to determine if the treatment is non-injurious to the target plant species. The user assumes responsibility for any plant damage or other liability resulting from use of Oryzalin Pro 4 on plant species not recommended on this label.

Aerial Application: Do not aerially apply this product.

Chemigation: Do not apply this product through any type of irritations system.

Grazing: Do not graze or feed forage from treated areas to livestock.

Precaution: Avoid spray drift to non-target areas when applying Oryzalin Pro 4. Spray drift may result in reduced emergence of non-target plants adjacent to the treated area. Poor weed control may result if directions are not followed. Over-application may result in crop injury or excessive soil residue.

Application

Soil Preparation

Oryzalin Pro 4 controls weeds growing from seed. Oryzalin Pro 4 will not control emerged weeds. Oryzalin Pro 4 does not control established weeds, weeds growing from stolons, rhizomes, or root pieces. Therefore, areas to be treated should be free of emerged weeds. Weed residues, pruning, and trash should be thoroughly mixed into the soil or removed prior to treatment. In field applications, the soil should be in good tilth and free of clods at the time of application.

Ground Application: Apply Oryzalin Pro 4 as a directed spray to the soil surface or over the top of plants. Use only a properly calibrated, low-pressure, herbicide sprayer that will apply the spray uniformly. Use screens no finer than 50 mesh in nozzles and in-line strainers. Apply the appropriate rate of Oryzalin Pro 4 as outlined in "Approved Uses" section of this label. In all cases, use sufficient water volume to obtain uniform coverage and deliver the desired rate of Oryzalin Pro 4 to the treated area. The volume of water used is not critical as long as the desired rate of Oryzalin Pro 4 is delivered uniformly across the area treated. When calibrating, determine the volume of water delivered by the sprayer to a given area (100 Sq ft, 1 acre, etc.). Then mix the desired rate of Oryzalin Pro 4 in the amount of water required to cover the entire area to be treated. As the amount of water used (spray volume) decreases, the importance of accurate calibration and uniform application increases. Check the sprayer daily to ensure proper calibration and uniform application. Maintain continuous agitation from mixing through application. Avoid spray pattern skips and overlaps that may result in incomplete coverage or over-application.

Hand Held or Backpack Sprayer Application: The amount of water used to apply Oryzalin Pro 4 herbicide is not critical, but should be sufficient for uniform coverage of the target area.

Calibrate by determining the volume of water required to treat 1,000 square feet. Use this calibration volume to determine the amount of water and Oryzalin Pro 4 herbicide needed to treat the target area (see the following calibration example). Note: Sprayer calibration (volume of spray needed to treat 1,000 square feet) will vary with each individual operator.

Steps in Calibration

1. Mark an area of 1,000 square feet (i.e. 20 by 50 feet, or 25 by 40).
2. Place the sprayer on a level surface and add water noting the final level of water in the spray tank.
3. Spray the marked area with a sufficient volume of water to provide uniform coverage. Refill the sprayer to the same level as before measuring the amount of water added. The measured water added to the sprayer is the volume needed to cover 1,000 square feet.
4. Determine the application rate (fl oz/1000 sq ft) for Oryzalin Pro 4 from the "Approved Uses" section of this label.
5. To each volume of water used, as measured in step 3, add the amount of Oryzalin Pro 4 as determined in step 4.

Example: If the sprayer used 2 gallons of water to cover 1,000 square feet and the desired application rate of Oryzalin Pro 4 is 3 fluid oz/1,000 square feet, then you would add 3 fluid ounces of Oryzalin Pro 4 to every 2 gallons of water to be used.

Mixing Directions

Shake Well Before Using

Precaution: Do not allow the spray mixture to siphon back into water source.

Oryzalin Pro 4-Alone

Make sure spray tank is clean and use only clean water. Fill spray tank $\frac{1}{2}$ - $\frac{1}{4}$ full. Start agitation and add the required amount of Oryzalin Pro 4. Continue agitation and finish filling the spray tank. Maintain continuous agitation until application is completed.

Oryzalin Pro 4 -Tank Mix Combinations

Prior to mixing, read and carefully follow all label instructions and precautions for each product added to the tank mixture. Vigorous, continuous agitation is required for all tank mixes of Oryzalin Pro 4. Sparger pipe agitators generally provide the best agitation in spray tanks.

Mixing Order: Fill the tank $\frac{1}{4}$ full with clean water. Start agitation and add different formulation types in the order indicated below, allowing time for complete mixing and dispersion after addition of each product. Allow extra mixing and dispersion time for dry flowable products. Add different formulation types in the following order: dry flowables (DF); wettable powders (WP); Oryzalin Pro 4 and other aqueous suspensions (AS); flowables (F); liquids (L); solutions (S); and emulsifiable concentrates (EC).

Continue agitation and finish filling the spray tank with clean water. Maintain agitation until application is completed. If spraying and agitation must be stopped before the spray tank is empty, the materials may settle to the bottom. Settled materials must be completely resuspended before spraying is continued. A sparger agitator is particularly useful for this purpose.

Premixing: When tank mixing, initial mixing and dispersion of certain dry flowable or wettable powder products may be improved by premixing with water (slurrying). Adding the slurried material to the spray tank through a 20 or 35 mesh wetting screen will help assure good initial dispersion.

Equipment Cleaning

If a buildup of material occurs on the walls of the spray tank, it should be removed between fillings by washing with soap and water and rinsing thoroughly. Tanks, lines, screens, and nozzles should be cleaned thoroughly after each use.

Activation and Cultivation

Oryzalin Pro 4 will remain stable on the soil surface up to 21 days following application. In the absence of timely rainfall, irrigation can be used to activate Oryzalin Pro 4. A minimum of one-half (1/2) inch of rain or its equivalent in sprinkler irrigation is necessary to activate Oryzalin Pro 4. If weeds begin to emerge due to lack of rainfall or irrigation, shallow cultivate 1-2 inches deep to destroy existing weeds, or remove them by hand. Shallow cultivation to a depth of 1-2 inches will enhance herbicidal effectiveness. If Oryzalin Pro 4 is not activated by rainfall, irrigation, or cultivation within 21 days of application, or existing weeds have not been removed, erratic weed control may result.

Weeds Controlled by Oryzalin Pro 4

Annual Grasses

Common Name	Scientific Name
Barley, little	<i>Hordeum pusillum</i>
Barnyardgrass (Watergrass)	<i>Echinochloa crus-galli</i>
Bluegrass, annual	<i>Poa annua</i>
Crabgrass, large	<i>Digitaria sanguinalis</i>
Crabgrass, smooth	<i>Digitaria ischaemum</i>
Crowfootgrass	<i>Dactyloctenium aegyptium</i>
Cupgrass, southwestern	<i>Eriochloa gracilis</i>
Foxtail, bristlegass	<i>Setaria magna</i>
Foxtail, giant	<i>Setaria faberi</i>
Foxtail, green (Pigeongrass)	<i>Setaria viridis</i>
Foxtail, robust	<i>Setaria robusta</i>
Foxtail, yellow	<i>Setaria glauca</i>
Goosegrass (Silver crabgrass)	<i>Eleusine indica</i>
Johnsongrass (Seeding only)	<i>Sorghum halepense</i>
Junglerice	<i>Echinochloa colonum</i>
Lovegrass, Mexican	<i>Eragrostis mexicana</i>
Lovegrass, orcutt	<i>Eragrostis orcuttiana</i>
Oat, wild	<i>Avena fatua</i>
Panicum, browntop	<i>Panicum fasciculatum</i>
Panicum, fall (Spreading panicgrass)	<i>Panicum dichotomiflorum</i>
Panicum, Texas (Bufflograss) (Coloradograss)	<i>Panicum texanum</i>
Ryegrass, Italian	<i>Cenchrus incertus</i>
Signalgrass (Brachiaria)	<i>Brachiaria</i> spp.
Sprangletop, red	<i>Leptochloa filiformis</i>
Witchgrass	

	Panicum capillare
--	-------------------

Broadleaf Weeds

Common Name	Scientific Name
Bittercress	Cardamine oligosperma
Carpetweed	Mollugo verticillata
Chickweed, common	Stellaria media
Fiddleneck, coast	Amsinckia intermedia
Filaree, redstem	Erodium cicutarium
Filaree, whitestem	Erodium moschatum
Groundsel, common	Senecio vulgaris
Henbit	Lamium amplexiculare
Knotweed, prostrate	Polygonum aviculare
Lambsquarters	Chenopodium album
Pigweed, prostrate	Amaranthus blitoides
Pigweed, redroot	Amaranthus retroflexus
Pigweed, spring	Amaranthus hybridus
Pigweed, tumble	Amaranthus albus
Puncturevine	Tribulus terrestris
Purslane, common	Portulaca oleracea
Pulsey, Florida (Florida purslane) (Mexican clover) (Pulsey)	Richardia scabra
Rocket, London	Sisymbrium irio
Rockpurslane, desert	Calandrinia ciliata
Shepherdspurse	Capsella bursa-pastoris
Spurge, prostrate	Euphorbia humistrata
Woodsorrel, yellow	Oxalis stricta

Weeds Suppressed by Oryzalin Pro 4

Control of the following weeds may be erratic, ranging from poor to excellent, depending upon soil temperature, time of germination, depth of seed in the soil, and amount and timing of soil moisture:

Common Name	Scientific Name
Horseweed	Conyza canadensis
Ladysthumb	Polygonum persicaria
Lettuce, prickly	Lactuca serriola
Mallow, common	Malva neglecta
Milkweed, climbing	Sarcostemma cynanchoides
Morningglory	Ipomoea spp.
Mustard, black	Brassica nigra
Mustard, wild	Brassica kaber
Nightshade, black	Solanum nigrum
Ragweed, common	Ambrosia artemisiifolia
Smartweed	Polygonum pensylvanicum
Sowthistle, annual	Sonchus oleraceus
Spurge, spotted	Euphorbia maculata
Teaweed (prickly sida)	Sida spinosa

Velvetleaf	Abutilon theophrasti
Wheat, volunteer	Triticum spp.

Approved Uses

Ornamental Plantings

Oryzalin Pro 4 is recommended for use on certain landscape container- and field-grown established ornamental plants including: trees, shrubs, ground covers/perennials, flowers, non-bearing fruit and nut trees, non-bearing vineyards, and in the production of ornamental bulbs (See "Ornamental Bulbs" section for special use directions.)

Broadcast Application Rates

Labeled Use Site	Length of Control	Oryzalin (qt/acre)	Oryzalin Pro 4 (fl oz/1000sq ft)	Minimum Time Between Applications (Months)	Total Amount Allowed Per Year (qt/acre)
Landscape Ornamentals	2-4months	2	1.5	2	8
	3-6 months	3	2.2	4	12
	4-8 months	4	3	4	12
Field-grown and container-grown ornamentals	2-4 months	2	1.5	3	8
	3-6 months	3	2.2	3	9
	4-8 months	4	3	3	12

Tank Mix Combinations

Tank mix combinations of Oryzalin Pro 4 plus Roundup/Glyphosate, and many other labeled herbicides may be used to control undesirable vegetation in ornamental areas. Oryzalin Pro 4 may also be tank mixed with Gallery herbicide and applied preemergence to broaden the spectrum of broadleaf weed control in ornamental areas. Applied as directed, these tank mixes of Oryzalin Pro 4 will provide control of susceptible weed species listed on the respective labels. Refer to tank mix product labels for specific use directions, precautions, and limitations before use.

Oryzalin Pro 4 Plus Roundup/Glyphosate: Tank mix combinations of Oryzalin Pro 4 plus Roundup are recommended to control existing undesirable vegetation. Applied as directed, Oryzalin Pro 4 plus Roundup will provide postemergence control of susceptible weed species listed on the label for Roundup and residual preemergence control of susceptible weed species listed on the label for Oryzalin Pro 4. Refer to the label for Roundup for specific use directions, precautions, and limitations before use.

Precautions: Do not apply sprays containing Roundup over the top of ornamental plants. Extreme care must be exercised to prevent sprays containing Roundup from coming in contact with foliage and stems of turfgrasses, trees, shrubs, or other desirable vegetation since severe damage or death may result. If spraying with Roundup in areas adjacent to desirable plants, use a shield to prevent spray from contacting foliage and stems of desirable plants.

Special Use Precautions:

Apply only to established plants that have been transplanted into their growing location for a sufficient period of time to allow the soil to be firmly settled around the roots from packing and rainfall or irrigation.

Rooted liners should be removed from their original growing containers and placed in new containers at least two weeks prior to treatment or injury may occur.

To avoid possible injury, do not apply Oryzalin Pro 4 to:

- Nursery, forest of Christmas trees: seedling beds, cutting beds, or transplant beds.

- Unrooted liners or cuttings that have been planted in pots for the first time.
- Pots less than four inches wide.
- Ground covers until they are established and well rooted.
- Ornamental planting where there is likelihood of runoff onto lawn areas.
- Areas containing dichondra or cool season turfgrass species.

On container grown ornamentals where weed seed germination continues for extended periods of time, do not make repeat applications of Oryzalin Pro 4 for at least 90 days or crop injury may occur.

Application of Oryzalin Pro 4 over the top of plants with newly forming buds may cause injury. In this situation, a directed spray is recommended.

For soils treated with Oryzalin Pro 4 during the previous season, plant only the ornamental species listed on this label or injury may occur.

Ice Plant: en establishing unrooted ice plant on coarse-textured soils in landscaping plantings, do not exceed the 2 quart per acre rate of Oryzalin Pro 4 or crop injury may occur.

Note: Injury on the following plant species has been observed following applications of Oryzalin Pro 4 and use is not recommended:

Deutzia gracilis (slender deutzia)
Pseudotsuga menziesii (Douglas-fir)
Thuja occidentalis 'Techny' (Techny arborvitae)
Tsuga canadensis (eastern hemlock)
Begonia spp. (begonia)
Coleus hybridus (coleus)

**Oryzalin Pro 4 may be used on the following established plant species:
(Note: limitations on recommended treatment methods.)**

Trees

<u>Scientific Name</u>	<u>Common Name</u>	<u>Recommended Treatment Method</u> F = Field Grown C = Container Grown
Abies balsamea	Fir, balsam	F
Abies concolor	Fir, white	F
Abies fraseri	Fir, fraser	F
Abies grandis	Fir, grand	F
Abies veitchi	Fir, Vietch	F
Abies lasiocarpa	Fir, alpine	F
Abutilon hybridum	Albus-flowering maple Luteus-flowering maple Roseus-flowering maple Tangerine-flowering maple Vesuvius red-flowering maple	F F F F F
Acer ginnala	Flame maple	F
Acer rubrum	Red sunset maple	F
Acer saccharinum	Silver maple	F

<i>Acer</i> spp.	Maple	F
<i>Alsophila australis</i>	Australian tree fern	C, F
<i>Areacastum romanzoffianum</i>	Queen palm	F
<i>Betula nigra</i>	Birch, river	F
<i>Betula papyrifera</i>	Paper birch	F
<i>Betula pendula</i>	Birch, white	F
<i>Bucida buxoides</i>	Black olive	F
<i>Carya</i> spp.	Pecan, ornamental	C, F
<i>Cedrus atlantica</i>	Atlas cedar	C, F
<i>Cedrus deodara</i>	Deodar cedar	C, F
<i>Ceratonia siliqua</i>	Carob	F
<i>Cercidium floridum</i>	Palo Verde, blue	F
<i>Cercis Canadensis</i>	Redbud	C, F
<i>Chamaecyparis lawsoniana</i>	Falsecypress, Lawson	F
<i>Chamaecyparis obtusa</i>	Filicoides-fern spray cypress	F
	Gracilis-slender Hinoki cypress	F
<i>Chamaecyparis pisifera</i>	Sawara-false cypress	F
	Squarrosa-moss cypress	F
<i>Chamaedorea cataractarum</i>	Cat Palm	F
<i>Chamaedorea costaricana</i>	Palm	F
<i>Chamaedorea elegans</i>	parlor palm	F
<i>Citrus</i> spp.	Citrus, ornamental	C, F
<i>Comus florida</i>	Dogwood, flowering	F
<i>Cryptomeria japonica</i>	Cryptomeria, Japanese	C, F
<i>Cupaniopsis anacardioides</i>	Carrot Wood	F
<i>Cupressus arizonica</i> (glabra)	Cypress, Arizona	C, F
<i>Cupressus glabra</i>	Arizona cypress	C, F
<i>Cupressocyparis leylandii</i>	Leyland cypress	C, F
<i>Cupressus sempervirens</i>	Cypress, Italian	C, F
<i>Dicksonia antarctica</i>	Tasmanian tree fern	C, F
<i>Elaeagnus angustifolia</i>	Russian olive	C, F
<i>Eucalyptus camaldulensis</i>	Red gum eucalyptus	F
<i>Eucalyptus cinerea</i>	Eucalyptus, mealy	F
	Silver dollar eucalyptus	F
<i>Eucalyptus nicholii</i>	Eucalyptus, narrow-leaved	F
<i>Eucalyptus sideroxylon</i>	Eucalyptus, red ironbark	F
<i>Ficus benjamina</i>	Ficus	F
<i>Fraxinus</i> spp.	Ash	F
<i>Ginkgo biloba</i>	Ginkgo (Maidenhair tree)	C, F
<i>Gleditsia triacanthos</i>	Honey locust	F
<i>Heteromeles arbutiflora</i>	Toyon	F
<i>Juniperus virginiana</i>	Redcedar, Eastern	F
<i>Koeleria paniculata</i>	Goldenrain tree	F
<i>Liquidambar styraciflua</i>	Sweetgum, American	C, F
<i>Magnolia</i> spp.	Magnolia	F
<i>Malus</i> spp.	Crabapple	F
<i>Morus alba</i>	White mulberry	F
<i>Picea abies</i>	Pendula-weeping Norway Spruce	F
	Repens-spreading Norway Spruce	F
	Spruce, Norway	F
<i>Picea engelmannii</i>	Spruce, Engelmann	F
<i>Picea glauca</i>	Spruce, white	F
	Conica-dwarf Alberta spruce	F
<i>Picea glauca conica</i>	Dwarf Alberta spruce	F

<i>Picea mariana</i>	Spruce, black	F
<i>Picea pungens</i>	Glauca-Colorado blue spruce	F
	Hoopsii-Hoop's blue spruce	F
	Koster-Koster blue spruce	F
	Spruce, Colorado	C, F
<i>Pinus aristata</i>	Bristlecone pine	F
<i>Pinus canariensis</i>	Canary Island pine	F
<i>Pinus contorta</i>	Shore pine, beach pine	F
<i>Pinus eldarica</i>	Eldarica pine	F
<i>Pinus halepensis</i>	Aleppo pine	C, F
<i>Pinus radiata</i>	Monterey pine	F
<i>Pinus</i> spp.	Pine	C, F
<i>Pinus strobus</i>	Eastern white pine	F
<i>Pinus sylvestris</i>	Scotch pine	F
<i>Pinus thunbergiana</i>	Japanese black pine	F
<i>Platanus occidentalis</i>	American sycamore	F
<i>Platanus racemosa</i>	California sycamore	F
<i>Podocarpus</i> spp.	Podocarpus	F
<i>Populus deltoides</i>	Cottonwood	F
	Cottonwood (grown for pulp)	
<i>Prunus caroliniana</i>	Laurelcherry, Carolina	F
<i>Prunus glandulosa</i>	Dwarf flowering almond	C, F
<i>Prunus laurocerasus</i>	Laurelcherry, English	F
<i>Prunus mahaleb</i>	Cherry, Mahaleb	F
<i>Prunus yedoensis</i>	Yoshino flowering cherry	F
<i>Pyrus communis</i>	Pear	F
<i>Quercus palustris</i>	Pin oak	F
<i>Quercus phellos</i>	Willow oak	F
<i>Quercus rubra</i>	Red oak	C, F
<i>Quercus</i> spp.	Oak	C, F
<i>Salix babylonica</i>	Babylon weeping willow	F
	Corkscrew willow	F
<i>Schinus molle</i>	California pepper tree	F
<i>Sequoia sempervirens</i>	Redwood, coast	F
<i>Sequoiadendron giganteum</i>	Giant sequoia	F
<i>Swietenia mahogani</i>	Mahogany	F
<i>Tabebuia caraiba</i>	Yellow Tab	F
<i>Tilia cordata</i>	Linden, little leaf	C, F
<i>Ulmus parvifolia</i>	Chinese elm	F
<i>Umbellularia californica</i>	California laurel	F
<i>Washingtonia robusta</i>	Mexican fan palm	F

Shrubs

<u>Scientific Name</u>	<u>Common Name</u>	<u>Recommended Treatment Method</u> F = Field Grown C = Container Grown
<i>Abelia grandiflora</i>	Glossy abelia	F
<i>Acacia redolens</i>	Acacia, prostrate	F
<i>Agave Americana</i>	Century plan	F
<i>Agave macroculmis</i>	Agave	F
<i>Anisodonteia hypomandarum</i>	Cape mallow	C, F

Arctostaphylos stanfordiana	Manzanita, Stanford	F
Astilbe chinensis	Astilbe/false spirea	C, F
Baccharis pilularis	Coyotebush	F
Berberis thunbergii	Aurea-golden Japanese barberry Crimson pygmy barberry Atropurea-redleaf Japanese barberry Barberry, Japanese	C, F C, F C, F C, F
Bougainvillea spp.	Barbara Karst California gold Scarlet O'Hara Texas Dawn	F F F F
Buxus microphylla	Littleleaf boxwood	F
Buxus microphylla japonica	Boxwood, Japanese	C, F
Buxus sempervirens	Boxwood, common	C, F
Callistemon citrinus	Bottlebrush, lemon	C, F
Cassia artemisioides	Cassia, feathery	F
Ceanothus americanus	Jerseytea, redroot	C, F
Ceanothus spp.	Wild lilac	C, F
Chaenomeles japonica	Flowering quince	C, F
Chamaecyparis obtusa	Kosteri cypress Nana-dwarf Hinoki cypress Torulosa cypress	F F F
Chamaecyparis pisifera	Squarrosa Minima cypress	F
Chamaecyparis pisifera spp.	Filifera-thread cypress	F
Chrysalidocarpus lutescens	Areca palm	F
Cleyera japonica	Cleyera, Japanese	C, F
Coleonema pulchrum	Pink breath of heaven	C, F
Comus alba	Sibirica-Siberian dogwood	F
Comus kousa	Dogwood, kousa	C, F
Comus stolonifera	Flaviramea-yellowtwig dogwood	F
Cotoneaster adpressus	Praecox-early cotoneaster	F
Cotoneaster apiculatus	Cotoneaster, cranberry	C, F
Cotoneaster buxifolius	Cotoneaster, brightbead	F
Cotoneaster congestus	Cotoneaster, Pyrenees	F
Cotoneaster dammeri	Cotoneaster, bearberry	C, F
Cotoneaster himalayan	Himalayan cotoneaster	F
Cotoneaster horizontalis	Cotoneaster, rock	C, F
Cotoneaster lacteus	Cotoneaster, pamey	C, F
Cotoneaster microphyllus	Cotoneaster, rockspray	F
Cotoneaster salicifolia	Willowleaf cotoneaster	C, F
Cytisus praecox	Hollandia-warminster broom	F
Cytisus scoparius	Lena-Scotch broom	F
Dasylium wheeleri	Sotol, desert spoon	F
Deutzia crenata	Nakiana-dwarf deutzia	F
Dodonaea viscosa	Hopseedbush, clammy Hopseed bush	F F
Escallonia exoniensis	Escallonia	C, F
Euonymus alata	Euonymus, winged	F
Euonymus fortunei	Canadale gold euonymus Emerald'n gold euonymus Euonymus, stringybark Wintercreeper	C, F C, F C, F C, F
Euonymus japonica	Euonymus, evergreen Silver king euonymus	C, F F

<i>Euonymus kiatschovica</i>	Spreading euonymus	F
<i>Euonymus vegetus</i>	Bigleaf wintercreeper	C, F
<i>Fatshedera lizei</i>	Fatshedera	C, F
<i>Fatsia japonica</i>	Japanese aralia	C, F
<i>Felicia amelloides</i>	Blue marguerite	C, F
<i>Forsythia intermedia</i>	Forsythia, border	F
<i>Gardenia jasminoides</i>	Gardenia	C, F
<i>Genista pilosa</i>	Woadwaxen	F
<i>Hibiscus rosa-sinesis</i>	Ross Estey-hibiscus	F
	Hibiscus, Chinese	F
<i>Hibiscus syriacus</i>	Rose of Sharon, Red Bird	F
	Rose of Sharon, Red Heart	F
	Rose of Sharon, Woodbridge	F
	Rose of Sharon (Shrubalthea)	F
<i>Ilex aquifolium</i>	Balkans holly	F
	Gold coast holly	F
	Holly, English	F
<i>Ilex aquipemyi</i>	San Jose holly	C, F
<i>Ilex cornuta</i>	Dwarf Burford holly	C, F
	Holly, Chinese	C, F
<i>Ilex crenata</i>	Compacta-dwarf Japanese holly	C, F
	Convexa holly	C, F
	Helleri-Heller's Japanese holly	C, F
	Holly, Japanese	C, F
<i>Ilex glabra</i>	Nordica-inkberry holly	F
<i>Ilex meserveae</i>	Blue boy holly	F
	Blue girl holly	F
	Ebony magic holly	F
<i>Ilex vomitoria</i>	Nana-dwarf yaupon holly	C, F
	Pendula-weeping yaupon holly	C, F
	Yaupon holly	C, F
<i>Juniperus chinensis</i>	Media-old gold juniper	C, F
<i>Juniperus conferta</i>	Emerald sea shore juniper	F
<i>Juniperus horizontalis</i>	Huntington blue juniper	C, F
	Wiltonii-blue carpet juniper	C, F
<i>Juniperus procumbens</i>	Nana-dwarf Japanese garden juniper	C, F
<i>Juniperus prostrata</i>	Prostrata juniper	C, F
<i>Juniperus sabina</i>	Broadmoor juniper	F
	Foemina-Hicks juniper	F
	Tamariscifolia-Tam juniper	F
<i>Juniperus scopulorum</i>	Emerald green juniper	F
<i>Juniperus spp.</i>	Juniper	C, F
<i>Juniperus squamata</i>	Blue juniper	F
	Blue star juniper	F
	Parsonii juniper	F
<i>Justicia brandegeana</i>	Shrimp plant	C, F
<i>Jusiticia spicigera</i>	Honeysuckle, Mexican	F
<i>Kalmia latifolia</i>	Laurel, mountain	F
<i>Lagerstroemia indica</i>	Crape myrtle	C, F
<i>Lavandula angustifolia</i>	English lavender	C, F
<i>Leucothoe axillaris</i>	Leucothoe, coast	F
<i>Leucothoe fontanesiana</i>	Leucothoe, drooping	F
<i>Ligustrum amurense</i>	Privet, amur	C, F
<i>Ligustrum japonicum</i>	Privet, Japanese	C, F

	Yellow tip ligustrum	C, F
Ligustrum lucidum	Privet, glossy	C, F
Ligustrum ovalifolium	California privet	F
Ligustrum texanum	Howardi privet Wax leaf privet	F F
Ligustrum vicaryi	Privet, golden Vicary golden privet	C, F C, F
Livistona chinensis	Chinese fountain palm	F
Lonicera fragrantissima	Winter honeysuckle	F
Lonicera periclymenum	Flowering woodbine Serotina woodbine	F F
Lonicera sempervirens	Trumpet honeysuckle	F
Lorpetalum chinense	(No common name)	C, F
Mahonia aquifolium	Oregon grape	F
Myoporum parvifolium	Myoporum, prostrate	F
Myrtus communis	Myrtle, true	C, F
Nandina domestica	Compacta-dwarf heavenly bamboo Harbour dwarf-heavenly bamboo Heavenly bamboo (Nandina) Nana compacta-heavenly bamboo Nana purpurea-heavenly bamboo Woods dwarf-heavenly bamboo	C, F C, F C, F C, F C, F C, F
Nerium oleander	Hardy red oleander Oleander Ruby lace oleander	C, F C, F C, F
Osmanthus heterophyllus	Osmanthus, holly-leaf	F
Pachysandra terminalis	Japanese spurge	C, F
Philadeiphus spp.	Mockorange	C, F
Phoenix roeloenii	Pigmy date palm	F
Photinia fraseri	Fraser's photinia Photinia	C, F C, F
Pieris japonica	Lily-of-the-valley Snowdrift lily-of-the-valley Temple bells lily-of-the-valley Valley rose lily-of-the-valley Andromeda	F F F F C, F
Pittosporum spp.	Pittosporum	C, F
Pittosporum tobira	Green pittosporum Japanese pittosporum Tibira Wheeler's dwarf pittosporum	F F F F
Platycladus orientalis	Arborvitae, Oriental	C, F
Plumbago ariculata	Blue cape plumbago	F
Podocarpus macrophyllus	Yewpine	C, F
Potentilla fragiformis	Cinquefoil	F
Potentilla fruticosa	Cinquefoil	C, F
Protea neriifolia	Protea	F
Pyracantha coccinea	Firethorn, scarlet	C, F
Pyracantha fortuneana	Lolendei Monrovia pyracantha	C, F
Pyracantha fortuneana	Monon pyracantha Red elf hybrid pyracantha Rutgers hybrid pyracantha Santa Cruz pyracantha Victory pyracantha	C, F C, F C, F C, F C, F

<i>Pyracantha skoidzumi</i>	Firethorn, formosa	C, F
<i>Pyracantha fortuneana</i>	Firethorn	C, F
<i>Raphiolepis indica</i>	Enchantress-Moness raphiolepis Raphiolepis (India hawthorn) Springtime-Monme raphiolepis	F C, F F
<i>Raphiolepis ovata</i>	Roundleaf raphiolepis	F
<i>Rhipsalidopsis gaertneri</i>	Eastercactus	C, F
<i>Rhododendron calendulaceum</i>	Flame azalea	F
<i>Rhododendron campylocarpum</i>	Butterfly rhododendron	F
<i>Rhododendron carolinianum</i> x <i>dauricum</i>	PJM rhododendron	F
<i>Rhododendron catawbiense</i>	Catawba album rhododendron Catawba rhododendron Lord Roberts rhododendron Rocket rhododendron	C, F C, F C, F C, F
<i>Rhododendron forrestii</i> x <i>griersonianum</i>	Elizabeth rhododendron	F
<i>Rhododendron hybrid</i> spp.	America rhododendron English Roseum rhododendron Nova Zembla rhododendron Scintillation rhododendron	F F F F
<i>Rhododendron impeditum</i>	Rhododendron	F
<i>Rhododendron indica</i>	Formosa azalea Waucabusa azalea	C, F C, F
<i>Rhododendron kerume</i>	Coral bells azalea Hino crimson azalea Hino pink azalea Snow azalea	C, F C, F C, F C, F
<i>Rhododendron maximum</i>	Rhodie max (rosebay)	C, F
<i>Rhododendron mucronulatum</i>	Rhododendron	F
<i>Rhododendron satsumi</i>	Gumpo pink azalea Higasa azalea	F F
<i>Rhododendron</i> spp.	Azalea Rhododendron	C, F C, F
<i>Rhododendron</i> spp. hybrids	Carror azalea Girard Roberta azalea Golden flare exbury azalea	C, F F F
<i>Rhus lancea</i>	Sumac, African	C, F
<i>Rosa rugosa</i>	Ramantas rose	F
<i>Rosmarinus officinalis</i>	Rosemary	F
<i>Senecio cineraria</i>	Dusty miller	C, F
<i>Spiraea vanhouttei</i>	Bridal wreath	F
<i>Syringa vulgaris</i>	Lilac, common	C, F
<i>Syzygium paniculata</i>	Brush cherry	C, F
<i>Taxus cuspidate</i>	Yew, Japanese	F
<i>Taxus media</i>	Yew	F
<i>Thuja occidentalis</i>	Arborvitae, American Emerald arborvitae Globosa-globe arborvitae Little giant—dwarf arborvitae Nigra-dark American arborvitae Pyramidalis arborvitae Rheingold arborvitae Woodwardii arborvitae	C, F F F F F F F
<i>Thuja orientalis</i>	Aureus nana-dwarf golden arborvitae	F

	Minima glauca-dwarf arborvitae	F
Thuja plicata	Red Cedar, Western	F
Trachelospermum jasminoides	Star jasmine, Chinese	F
Veitchia merrilli	Christmas palm	F
Viburnum carlesii	Koreanspice viburnum	C, F
Viburnum davidii	David viburnum	F
Viburnum japonicum	Viburnum	F
Viburnum judd (V X Judii)	Viburnum	C, F
Viburnum opulus sterile	Common snowball viburnum	F
Viburnum plicatum tomentosum	Doublefile viburnum	F
Viburnum setigerum	Tea viburnum	F
Viburnum suspensum	Viburnum Sandankwa	F
Viburnum tinus	Viburnum Laurustinus Compactum-spring bouquet viburnum	C, F F
Viburnum tinus compactum	Spring bouquet viburnum	F
Viburnum trilobum Compactum	Dwarf cranberry bush	F
Viburnum x pragense	Viburnum	F
Weigela florida	Bristol ruby weigela Java red weigela Minuet weigela Weigela, oldfashioned	F F F F
Xylosma congestum	Xylosma	F
Yucca elata	Yucca soaptree	C, F
Yucca recurvifolia	Yucca pendulous	F

Groundcovers/Perennials

<u>Scientific Name</u>	<u>Common Name</u>	<u>Recommended Treatment Method</u> F = Field Grown C = Container Grown
Agapanthus africanus	Lily-of-the-Nile	C, F
Ajuga spp.	Carpet bugle	F
Arctotheca calendula	Cape weed	F
Asparagus retrofractus	(No common name)	C, F
Asparagus varieegata	Tree fern	C, F
Aster novae-angliae	New England aster	C, F
Aster novi-belgii	New York aster	C, F
Athyrium nipponimcum	Japanese painter fern	C, F
Brassica oleracea	Wild cabbage	C, F
Callistepheus chinensis	China Aster	C, F
Campanula elatines	Bellflower	C, F
Carpobrotus edulis	Ice plant, largeleaf (see label)	F
Clytostoma callistegiodes	Trumpet vine, violet	C, F
Cortaderia selloana	Pampas grass	F
Cuphea hyssopifolia	False Mexican heather	C, F
Delosperma alba	White iceplant	F
Dietes vegeta	Fortnight lily	C, F

<i>Digitalis mertonensis</i>	Foxglove	C, F
<i>Doronicum cordatum</i>	Leopard's bane	C, F
<i>Drosanthemum floribundum</i>	Trailing rosea iceplant	F
<i>Erianthus ravennae</i>	Hardy pampus grass	C, F
<i>Festuca ovina glauca</i>	Blue fescue	F
<i>Gaillardia grandiflora</i>	Blanket flower	C, F
<i>Gazania rigens leucolaena</i>	Gazania, trailing	C, F
<i>Gazania</i> spp.	Gazania	F
<i>Hedera canariensis</i>	Ivy, Algerian	F
<i>Hedera helix</i>	Ivy, English	F
<i>Heliotropium fragrans</i>	Common heliotrope	C, F
<i>Hemerocallis</i> spp.	Daylily	C, F
<i>Hosta lancifolia</i>	Albo-marginata hosta	C, F
<i>Hosta</i> spp.	Lily, plantain	C, F
<i>Heuchera micrantha</i>	Coral bells	C, F
<i>Hypericum</i> spp.	St. Johnswort	C, F
<i>Iberis sempervirens</i>	Evergreen candytuft	C, F
<i>Lampranthus spectabilis</i>	Trailing iceplant	F
<i>Leptospermum scaparium</i>	New Zealand teatree/Manuka	C, F
<i>Limonium perezii</i>	Statice/Sea lavender	C, F
<i>Liriope gigantea</i>	White lily turf	F
<i>Liriope muscari</i>	Lilac beauty lily turf	C, F
	Majestic lily turf	C, F
	Monroe white lily turf	C, F
	Silvery sunproof lily turf	C, F
	Variegated liriope lily turf	C, F
	Big blue lily turf	C, F
<i>Lobelia erinus</i>	Edging lobelia	C, F
<i>Lonicera japonica</i>	Honeysuckle, Japanese	F
<i>Mesembryanthemum crystallinum</i>	Ice plant (see label)	F
<i>Ophiopogon japonicus</i>	Mondo grass	F
<i>Osteospermum fruitocum</i>	Daisy, trailing African	F
<i>Pachysandra terminalis</i>	Japanese spurge	F
<i>Pennisetum setaceum</i>	Fountaingrass	C, F
<i>Polystichum polyblepharum</i>	Tassel fern	C, F
<i>Sedum brevifolium</i>	Stonecrop	C, F
<i>Sedum kamtschaticum</i>	Stonecrop	C, F
<i>Sedum spurium</i>	Stonecrop, two row	C, F
<i>Tulbaghia violacea</i>	Society garlic	C, F
<i>Verbena rigida</i>	Veined verbena	C, F
<i>Veronica</i> spp.	Speedwell	C, F
<i>Vinca major</i>	Periwinkle, bigleaf	F
<i>Vinca minor</i>	Periwinkle, dwarf	F

Flowers

<u>Scientific Name</u>	<u>Common Name</u>	<u>Recommended Treatment Method</u> F = Field Grown C = Container Grown
<i>Actillea</i> app.	Yarrow	C, F
<i>Antirrhinum majus</i>	Snapdragon	F

Caladium bicolor	Caladium, fancyleaf	F
Chrysanthemum spp.	Chrysanthemum	C, F
Coreopsis lanceolata	Coreopsis	F
Coreopsis verticulata	Threadleaf coreopsis	C, F
Dianthus barbatus	Sweet William	F
Dianthus gratianopolitanus	Cheddar pink	C, F
Dicentra spectabilis	Bleeding heart	C, F
Dimorphotheca spp.	Marigold, cape	F
Echinacea purpurea	Coneflower, purple	C, F
Evolvulus nuttallianus	Blue daze	C, F
Geum quellyon	Geum	F
Gladiolus hortulanus	Gladiolus	F
Gypsophila paniculata	Baby's breath	F
Impatiens wallerana	Impatiens (Busy lizzy)	F
Iris spp.	Iris, bearded	F
Liatris spicata	Blazing star	C, F
Pelargonium hortorum	Geranium	F
Petunia spp.	Petunia	C, F
Portulaca grandiflora	Moss, rose	F
Ranunculus asiaticus	Ranunculus, Persian	F
Rosa spp.	Rose	F
Rudbeckia fulgida	Blackeyed susan	C, F
Rudbeckia hirta	Daisy, gloriosa (black-eyed Susan)	F
Salvia spp.	Salvia (Sage)	F
Stokesia laevis	Aster, stokes	F
Strelitza reginae	Bird of paradise	F
Tagetes spp.	Marigold	F
Viola wittrockiana	Pansy	F
Zinnea elegans	Zinnia, common	F

Non-bearing * Trees and Vines

	<u>Recommended Treatment Method</u> F = Field Grown C = Container Grown
Almond	F
Apple	F
Apricot	F
Avocado	F
Blackberry	F
Blueberry	F
Boysenberry	F
Cherry, sour	F
Cherry, sweet	F
Currant	F
Dewberry	F
Elderberry	F
Fig	F
Filbert	F
Gooseberry	F
Grape, American	F
Grape, European	F

Grapefruit	F
Kiwi	F
Kumquat	C, F
Lemon	F
Loganberry	F
Macadamia nut	F
Nectarine	F
Olive	F
Orange	C, F
Peach	F
Pear	F
Pecan	C, F
Pistachio	F
Plum	F
Pomegranate	F
Prime	F
Raspberry	F
Walnut, black	F
Walnut, English	F

*Non-bearing plants are defined as those that will not bear fruit for at least one year after treatment.

Ornamental Bulbs

Oryzalin Pro 4 may be applied for control of susceptible annual weeds in ornamental bulbs, e.g. bulbous iris, daffodil (narcissus), hyacinth, and tulip. Apply Oryzalin Pro 4 to the soil surface 2-4 weeks after planting, but prior to the emergence of annual weeds. For fall planted bulbs, apply Oryzalin Pro 4 again in late winter or early spring to weed-free soil surfaces.

Broadcast Application Rates

Time of Application	Soil Texture	Oryzalin Pro 4 (qt/acre)	Oryzalin Pro 4 (fl oz/1000 sq ft)	Minimum Time Between Applications (months)	Total Amount Allowed Per Year (qt/acre)
Fall	Coarse	0.75	0.5	3	1.5
Fall	Medium and Fine	1.5	1.0	3	2.25
Feb-March	All Soil Textures	0.75	0.5	3	2.25

Special Use Precautions

Do not apply to tulip plants that have emerged to a height greater than 3/4 inch.

Do not apply to gladioli corins prior to emergence or less than one (1) inch in diameter.

Shadehouse Areas

Oryzalin Pro 4 may be applied to drainage areas under benches in open shadehouse-type structures where the natural flow of air is unimpeded. Do not apply in enclosed greenhouses or in enclosed shadehouse type structures. Do not apply within 3 weeks prior to enclosure of greenhouse or poly-type structures.

Christmas Tree Plantations

Oryzalin Pro 4 alone

Apply Oryzalin Pro 4 as a directed spray to the soil surface or as an overtop spray to established plantings of field grown Christmas tree species, including fir (*Abies* spp.), Pine (*Pinus* spp.), and spruce (*Picea* spp.). Follow all instructions provided in the "General Information" section of this label.

Broadcast Application Rates

Length of Control	Oryzalin Pro 4 (qt/acre)	Oryzalin Pro 4 (fl oz/1000 sq ft)	Minimum Time Between Applications (months)	Total Amount Allowed Per Year (qt/acre)
2-4 months	2	1.5	2	8
4-8 months	4	3	2	8

Tank Mix Combinations

Tank mix combinations of Oryzalin Pro 4 plus other labeled herbicides may be used as directed sprays or overtop sprays in established Christmas tree plantings. When applied according to use directions, these tank mixes will provide control of susceptible weed species listed on the respective product labels. Refer to tank mix product labels for specific use directions, precautions, and limitations before use.

Oryzalin Pro 4 Plus Roundup/Glyphosate: Apply tank mix combinations of Oryzalin Pro 4 plus Roundup only as directed sprays in Christmas tree plantings. When applied according to use directions, Oryzalin Pro 4 plus Roundup will provide postemergence control of susceptible weed species listed on the label for Roundup and residual preemergence control of susceptible weed species listed on the label for Oryzalin Pro 4. Refer to the label for Roundup for specific use directions, precautions, and limitations before use.

Special Use Precautions

Do not apply to Douglas-fir (*Pseudotsuga menziesii*). Do not apply to seedbeds or seedling transplant beds. Apply only to established plants that have been transplanted into their final growing location for a sufficient period of time to allow the soil to be firmly settled around the roots from packing and rainfall or irrigation.

Noncropland Areas and Industrial Sites

Noncropland Areas—Tank Mix Combinations

Tank mix combinations of Oryzalin Pro 4 plus Roundup/Glyphosate and many other labeled herbicides may be used to control undesirable vegetation in noncropland areas such as roadsides, rights-of-way, etc. When applied according to use directions, these tank mixes will provide control of susceptible weed species listed on the respective product labels. Refer to tank mix product labels for specific use directions, precautions, and limitations before use.

Broadcast Application Rates

Length of Control	Oryzalin Pro 4 (qt/acre)	Oryzalin Pro 4 (fl oz/1000 sq ft)	Minimum Time Between Applications (months)	Total Amount Allowed Per Year (qt/acre)
2-4 months	2	1.5	2	6
4-8 months	4	3	4	12
8-12 months	6	4.5	8	12

Industrial Sites—Tank Mix Combinations

Tank mix combinations of Oryzalin Pro 4 plus Roundup, Spike* herbicide, and many other labeled herbicides may be used as overtop sprays to control existing vegetation on industrial sites such as utility substations, highway guard rails, sign posts, and delineators. When applied according to use directions, these tank mixes will provide control of susceptible weed species listed on the respective product labels. Refer to tank mix product labels for specific use directions, precautions, and limitations before use.

Warm Season Turfgrasses

Oryzalin Pro 4 may be applied as a preemergence treatment for control of annual grasses and certain broadleaf weeds in established warm season turf including bahiagrass, bermudagrass, buffalograss, centipedegrass, St. Augustinegrass, zoysiagrass, and established tall fescue growing in warm season areas. Established turf is defined as a dense turf having a well-anchored root system and healthy vigorous top growth. Use Oryzalin Pro 4 only as a part of a total turf management program that includes good fertilization practices.

Oryzalin Pro 4 may be tank mixed with Gallery* herbicide (California registration pending) and applied preemergence to broaden the spectrum of broadleaf weed control in warm season turf. Refer to the label for Gallery for specific use directions, precautions, and limitations before use.

Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of Oryzalin Pro 4.

Oryzalin Pro 4 will not control emerged weeds. Successful preemergence control of weeds listed on this label requires that Oryzalin Pro 4 be applied prior to weed germination and be activated by at least one-half (1/2) inch of rainfall or irrigation within 21 days of application.

Oryzalin Pro 4 may injure turf that is not well-established or is stressed or weakened due to unfavorable winter climatic conditions, drought, nematodes, or other factors which damage or weaken turf root systems. Apply Oryzalin Pro 4 only to healthy, well-established turf that has a well-anchored root system.

Do not apply Oryzalin Pro 4 in the spring or early summer to tall fescue turfgrass reseeded the previous fall. In such cases, apply Balan* 2.5G granular herbicide at 60-80 pounds per acre in early summer (Round 1) and Oryzalin Pro 4 at 1.5 quarts per acre approximately eight weeks later (Round 2). Do not apply Oryzalin Pro 4 at the single application rate (2 quarts per acre) to established tall fescue; in such cases, apply 1.5 quarts per acre of Oryzalin Pro 4 in an initial application, followed by a second application of 1.5 quarts per acre 8-10 weeks later.

In Bermudagrass areas that have been overseeded with winter grasses, a spring application of Oryzalin Pro 4 will thin the overseeded grasses.

Annual Grasses Controlled by Oryzalin Pro 4 Summer Annuals:

<u>Common Name</u>	<u>Scientific Name</u>
Barnyardgrass (watergrass)	Echinochloa crus-galli
Crabgrass, large	Digitaria sanguinalis
Crabgrass, smooth	Digitaria ischaemum
Crabgrass	Digitaria spp.
Crowfootgrass	Dactyloctenium aegyptium
Foxtail, bristleglass	Setaria magna
Foxtail, giant	Setaria faberi
Foxtail, green (pigeongrass)	Setaria viridis
Foxtail, robust	Setaria robusta

Foxtail, yellow	Setaria glauca
Goosegrass (silver crabgrass)	Eleusine indica
Johnsongrass (seedling only)	Sorghum halepense
Ryegrass, Italian	Lolium multiflorum
Sandbur, field	Cenchrus incertus

Winter Annuals:

<u>Common Name</u>	<u>Scientific Name</u>
Bluegrass, annual	Poa annua

Annual Broadleaf Weeds Controlled by Oryzalin Pro 4 Summer Annuals:

<u>Common Name</u>	<u>Scientific Name</u>
Carpetweed	Mollugo verticillata
Knotweed, prostrate	Polygonum aviculare
Purslane, common	Portulaca oleracea

Winter Annuals:

<u>Common Name</u>	<u>Scientific Name</u>
Chickweed, common	Stellaria media
Henbit	Lamium amplexicaule

Broadleaf Weeds Suppressed by Oryzalin Pro 4

<u>Common Name</u>	<u>Scientific Name</u>
Groundsel, common	Senecio vulgaris
Spurge, prostrate	Euphorbia humistrata
Woodsorrel, yellow	Oxalis stricta

Application Rates, Frequency, and Timing of Application

Oryzalin Pro 4 can be applied in the spring for summer annual grass and broadleaf weed control, and in the fall for annual bluegrass (*Poa annua*) and winter annual broadleaf weed control.

Broadcast Application Rates (Warm Season Turfgrasses)

Use Area	Oryzalin Pro 4 (qt/acre)	Oryzalin Pro 4 (fl oz/1000 sq ft)	Minimum Time Between Applications (months)	Total Amount Allowed Per Year (qt/acre)
All, except Florida	1.5 to 2	1 to 1.5	3	6
Florida	1.5	1	3	4.5

1. Summer Annual Grasses and Broadleaf Weeds

Single Application Program: Apply 2 quarts per acre of Oryzalin Pro 4 in late winter or early spring, prior to the onset of conditions favorable for annual weed germination.

Split Application Program: As an alternative to a single application program, Oryzalin Pro 4 may be applied in a split application. This program is desirable when the initial application is made well in advance of weed germination and where weed control is desired for a longer period of time. Apply 1.5 quarts per acre of Oryzalin Pro 4 in an initial application, followed by a second application of 1.5 quarts per acre 8-10 weeks later.

The second treatment of the split application may follow application of a different preemergence grass herbicide in place of the initial application of Oryzalin Pro 4.

2. Annual Bluegrass (*Poa annua*) and Winter Annual Broadleaf Weeds

In areas of heavy annual bluegrass infestation, its elimination will result in temporary thinning of turfgrass cover. Proper fertilization, irrigation, and soil-incorporated reseeding should be employed to speed the restoration of desirable turfgrass cover in areas previously occupied by annual bluegrass (see section on reseeding).

Apply Oryzalin Pro 4 as a preemergence treatment in late summer or early fall, prior to the expected germination period for annual bluegrass and winter annual broadleaf weeds. If annual bluegrass infestation is severe and its elimination will result in thinning of turfgrass cover, apply Oryzalin Pro 4 at 1.5 quarts per acre. If thinning of turfgrasses cover is not a potential problem, Oryzalin Pro 4 may be applied at 2 quarts per acre.

Weed Control in Florida

In Florida, apply 1.5 quarts per acre of Oryzalin Pro 4 three times per year, or every 90 — 100 days in the fall, early spring, and early summer. Do not apply more than 1.5 quarts per acre of Oryzalin Pro 4 in any single application.

Application Equipment

Apply Oryzalin Pro 4 evenly over the turfgrass area. Avoid spray pattern skips and overlaps that may result in incomplete coverage or over-application. For best results, use application equipment designed to uniformly broadcast liquid herbicides. Calibrate application equipment prior to use, according to manufacturer's directions. Check equipment frequently to make sure it is working properly and distributing spray uniformly.

Reseeding

Herbicides that control annual weeds may also affect establishment of desirable turfgrass seedlings. Reseeding should be delayed for at least 90-120 days following application of Oryzalin Pro 4. When reseeding, it is essential that proper cultural practices such as soil cultivation and seedbed preparation, irrigation, and fertilization be followed. For satisfactory reseeding results following use of Oryzalin Pro 4 the seeding rate should be increased and equipment designed to place seed in full contact with soil (such as the Rogers Aero Seeder) should be employed.

Special Use Precautions

To avoid possible injury, do not apply Oryzalin Pro 4 to:

- Cool season turfgrass species
- Golf course putting greens and tees or lawns containing dichondra or cool season turfgrass species.
- Newly sprigged or sodded areas of bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass until these turfgrasses are well-established and have well-anchored root systems.
- Newly hydromulched areas of bermudagrass until such areas are well-established.
- Bermudagrass variety "Sun Turf" when tank mixed with atrazine.

IMPORTANT INFORMATION. READ BEFORE USING PRODUCT.

IMPORTANT: Read the information below before using this product. If the terms are not

acceptable, you should return the unopened product container immediately for a complete refund.

WARRANTY STATEMENT

Control Solutions, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Control Solutions, Inc. To the extent allowed by law, Control Solutions, Inc. shall in no event be liable for consequential, special, or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. In addition to the foregoing, no purchaser of this product (other than an end user) shall be entitled to any reimbursement for any loss suffered as a result of any suspension or cancellation of the registration for this product by the U.S. Environmental Protection Agency. Except, as expressly provided herein, Control Solutions, Inc. makes no warranties, guarantees, or representations of any kind, either expressed or implied, or by usage of trade, statutory or otherwise, with regard to the product sold, including, but not limited to merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage. The exclusive remedy of any buyer or user of this product for any and all losses, injuries, or damage resulting from or in any way arising from the use, handling, or application of this product, whether in contract, warranty, tort, negligence, strict liability, or otherwise, shall be damages not exceeding the purchase price paid for this product or, at Control Solutions, Inc. election, the replacement of this product.