



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (H7505C)  
401 "M" St., S.W.  
Washington, D.C. 20460

EPA Reg. Number:  
53883-172

Date of Issuance:  
JUN 8 2005

NOTICE OF PESTICIDE:

Registration  
 Reregistration

(under FIFRA, as amended)

Term of Issuance:  
Conditional

Name of Pesticide Product:  
Prodiamine 0.37 Plus

Name and Address of Registrant (include ZIP Code):

Control Solution  
5903 Genoa-Red Bluff  
Pasadena, TX 77507

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

- 1. Add the phrase "EPA Registration No.53883-172" to the label before you release the product for shipment.

COMMENTS CONTINUED ON PAGE 2 OF THIS NOTICE OF REGISTRATION

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product under the enclosed stamped copy of the label constitutes acceptance of these conditions.

Enclosure

Joanne I. Miller  
Product Manager (23)  
Herbicide Branch  
Registration Division (7505C)

Signature of Approving Official:

*Joanne I. Miller*

Date:

JUN 8 2005

- 2. Submit the following data required for the registration of this pesticide product within 1 year from the date of this Notice of Registration:

<u>EPA Guideline Data Number</u>	<u>Guideline Descriptor</u>
830.6317	Storage Stability Study
830.6320	Corrosion Characteristics Study

- 3. On the front panel of the labeling and on page 3, delete the claim "selective" as a descriptor of this pesticide product.
- 4. On the front panel, delete the descriptor "UGLY"; "& MORE". and either revise the "Active Ingredient" chemical name for prodiamine to the Chemical Abstract name or show the name of the active ingredient as described in PR Notice 97-5(IV-D).
- 5. On page 3 add a map showing the approximate crabgrass germination dates in all areas where this product is to be marketed. In the "USE PRECAUTIONS" section, sixth line correct the statement that reads "Do not apply....." to read: "Do not apply to newly set sod until...".
- 6. On page 4, under "When To Overseed After Application" add a map which accompanies the chart "Months Before Overseeding". The map must clearly show the North, Transition and South Regions of the United States, where this product is to be marketed.
- 7. Submit one (1) copy of the final printed labeling before you release this product for shipment.
- 8. Submit and/or cite all data required for the registration of this product when the Agency requires all registrants of similar products to submit data; and submit acceptable responses required for reregistration of this product under FIFRA, section 4. A Stamped copy of the label is enclosed for your records.

---

Enclosure

**PRODIAMINE 0.37 PLUS**  
**FOR SELECTIVE PRE-EMERGENCE CONTROL OF GRASS AND**  
**BROADLEAF WEEDS IN LAWNS AND ORNAMENTAL TURF**  
(excluding golf course putting greens)  
**LANDSCAPE ORNAMENTALS, ESTABLISHED PERENNIALS AND**  
**WILDFLOWER PLANTINGS**  
**STOPS MANY UGLY WEED GRASSES BEFORE THEY GROW**  
**PREVENTS CRABGRASS, ANNUAL BLUEGRASS, FOXTAIL & MORE**

**ACTIVE INGREDIENT:**

Prodiamine: [N <sup>3</sup> N <sup>3</sup> -Di-n-propyl-2,4-dinitro-6-(trifluoromethyl)-m-phenylenediamine]	.0.37%
<b>INERT INGREDIENTS:</b>	<u>99.63%</u>
TOTAL:	100.00%

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

See back panel for further precautions and First Aid information

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION**

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals.

EPA REG NO. 53883-XXX  
EPA EST. NO.

Control Solutions, Inc.  
5903 Genoa-Red Bluff Rd.  
Pasadena, TX 77507-1041  
281-892-2500

**ACCEPTED**  
**with COMMENTS**  
**In EPA Letter Dated:**

JUN 8 2005  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

53883-172

NET CONTENT: \_\_\_\_\_

<b>FIRST AID</b>	
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>● Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>● Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>● Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>● Take off contaminated clothing.</li> <li>● Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>● Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>● Call a poison control center or physician immediately for treatment advice.</li> <li>● Have person sip a glass of water if able to swallow.</li> <li>● Do not induce vomiting unless told to do so by the poison control center or physician.</li> <li>● Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>● Move person to fresh air.</li> <li>● If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>● Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.</p>	

#### **ENVIRONMENTAL HAZARDS**

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

#### **DIRECTION FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product is a selective pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in established lawns and ornamental turf (excluding golf course putting greens), landscape ornamentals and established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seeds, germination and root development. When used according to directions, it will control; barnyardgrass; bluegrass; annual (*Poa annua*); carpetweed; chickweed, common; chickweed, mouseear (from seed); crabgrass (large, smooth); crowfootgrass; cupgrass, woolly; foxtails, annual; goosegrass<sup>1</sup>; henbit; itchgrass; johnsongrass (from seed); junglerice; knotweed; kochia; lambsquarter, common; lovegrass; panicum (Texas, fall, browntop); pigweed; purslane, common; pusley, Florida; rescuegrass<sup>3</sup>, shepherd's purse<sup>2</sup>, signalgrass, broadleaf; speedwell, Persian; sprangletop; spurge, prostrate; witchgrass; woodsorrel, yellow (from seed).

<sup>1</sup>In many areas a single application of 4 to 9.3 pounds/1,000 square feet (175 to 405 pounds/acre) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 4 to 6.2 pounds/1,000 square feet (175 to 270 pounds/acre), followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see "Minimum and Maximum Application Rate by Turfgrass Species" table in the Rates of Application section of this label).

<sup>2</sup>For this weed, apply in late summer, fall or winter prior to germination.

<sup>3</sup>Suppression only.

**WHEN TO APPLY TO ESTABLISHED TURF:** This product must be applied prior to germination of weed seeds, as it will not control established weeds.

**[MAP OF USA]**

**THIS MAP SHOWS  
APPROXIMATE CRABGRASS  
GERMINATION DATES**

**APPLICATION DIRECTIONS:** Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days application.

**USE PRECAUTIONS:** The following precautions apply to the use of this product in turf grasses and lawns. Application of this product may thin emerged annual bluegrass and newly overseeded grasses. Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil). Do not apply newly set sod until the following year. Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural practices (such as core aeration) may result in reduced weed control. Do not apply to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species. Do not apply through any type of irrigation system. Do not apply aerially. Do not graze or feed livestock forage cut from treated areas.

**RATES OF APPLICATION:** This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not exceed the maximum yearly rate listed:

<b>MINIMUM AND MAXIMUM APPLICATION RATES (by Turf Grass Species)</b>		
<b>Turf Species</b>	<b>Minimum Rate</b>	<b>Maximum Annual Rate<sup>1</sup></b>
Bahiagrass, Bermudagrass <sup>2</sup> Centipedegrass, Zoysiagrass, Seashore Paspalum, St. Augustinegrass Tall Fescue (including turf- Type)	4 lbs/1,000 sq. ft. (175 lbs/acre)	9.3 lbs/1,000 sq. ft. per year (405 lbs/acre per year)
Buffalograss Kentucky Bluegrass Perennial Ryegrass	4 lbs/1,000 sq. ft. (175 lbs/acre)	6.2 lbs/1,000 sq. ft. per year (270 lbs/acre per year)
Creeping Red Fescue	4 lbs/1,000 sq. ft. (175 lbs/acre)	4.6 lbs/1,000 sq. ft. per year (202 lbs/acre per year)
Creeping Bentgrass	4 lbs/1,000 sq. ft. (175 lbs/acre)	4.0 lbs/1,000 sq. ft. per year (175 lbs/acre per year)

<sup>1</sup>These are the maximum rates per calendar year.

<sup>2</sup>May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 3.3 pounds/1,000 square feet (145 pounds/acre). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control.

**WHEN TO OVERSEED AFTER APPLICATION:** This product will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/reseeding results.

<b>USE RATE</b>	<b>MONTHS BEFORE OVERSEEDING</b>		
	<b>North</b>	<b>Transition</b>	<b>South</b>
3 pounds/1,000 sq. ft. (135 pounds/acre)	4	4	4
4 pounds/1,000 sq. ft. (175 pounds/acre)	5	4	4
4.6 pounds/1,000 sq. ft. (202 pounds/acre)	6	5	5
4.9 pounds/1,000 sq. ft. (215 pounds/acre)	-	6	6
6.2 pounds/1,000 sq. ft. (270 pounds/acre)	-	7	7
7 pounds/1,000 sq. ft. (308 pounds/acre)	-	-	9
8 pounds/1,000 sq. ft. (350 pounds/acre)	-	-	10
9.3 pounds/1,000 sq. ft. (405 pounds/acre)	-	-	12

**LANDSCAPE ORNAMENTALS:** (including established perennials and wildflower plantings): This product may be applied for residual pre-emergence weed control in ornamentals. Apply at the rate of 4 to 9.3 pounds/1,000 square feet (175 to 405 pounds/acre) in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 9.3 pounds/1,000 square feet (405 pounds/acre) per year.

SPREADER SETTINGS to apply at the listed pounds per 1,000 square feet						
	3 lbs.	4 lbs.	4.6 lbs.	6.2 lbs.	8 lbs.	9.3 lbs.
Spreader A	X	X	X	X	X	X
(May contain additional spreader brands/models, spreader settings and spreader use instructions as appropriate for the product)						

**APPLICATION TIMING AND INFORMATION:** This product may be applied to newly transplanted and established ornamentals a broadcast, over the top or directed application. Around new transplants, delay application to allow soil to settle and water thoroughly before applying. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union. This product is a pre-emergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when applied to soil free of clods, weeds and debris such as leaves. Prior to application, control existing vegetation by hand weeding, cultivation or the use of an appropriate post-emergence herbicide.

**TOLERANT ORNAMENTAL SPECIES:** This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

**SCIENTIFIC NAME**

- Abies spp.
- Abelia grandiflora
- Acer palmatum
- A. Platanoides
- Achillea spp.
- Actinidia chinensis
- Agapanthus africanus
- Agapanthus orientalis
- Akebia quintata
- Allium cernuum
- Anemone hybrida
- Aquilegia spp.
- Arctostaphylos densiflora
- Arctotheca calendula
- Artemisia spp.
- Aster spp.
- Aster X Frikartii
- Athrium Filiz-femina
- Aucuba japonica
- Begonia spp.
- Berberis gladwynensis
- B. Julianae
- B. mentorensis
- B. Thunbergii
- B. Verruculosa
- Bergenia cordifolia
- Boltonia asteroides
- Buddleia davidii
- Buxus microphylla
- Callistemon viminalis

**COMMON NAME**

- Fir species\*\* (Balsam, Frasier, Noble, etc.)
- Abelia: Sherwood
- Japanese Maple
- Norway Maple
- Yarrow: King Edward
- Kiwi\*
- Lily-of-the-Nile (African Lily)
- Five-leaf or Chocolate Vine
- Lady's Leek, Nodding Onion
- Japanese Anemone
- Aquilegia: Red and Gold
- Vine Hill Manzanita
- Cape Weed
- Wormwood; Silver Mound, Castle
- Aster: Bonnie Blue, Purple Dome
- Lady Fern; Fern Lady
- Japanese Aucuba
- Fibrous Begonia: Hardy Grandis
- Barberry
- Wintergreen Barberry
- Mentor Barberry
- Japanese Barberry
- Warty Barberry
- Snowbank
- Butterfly-Bush (Dwarf Blue); Royal Red
- Japanese Boxwood
- Weeping Bottlebrush

Calluna vulgaris	Scotch Heather
Campanula carpatica	Tussock Bellflower, (White Clips)
Campis X Tagliabuana	Trumpet Flower, Madame Galen
Carpobrotus edulis	Hottentot Fig (Ice Plant)
Cassia artemisoides	Feathery Cassia
Ceanothus rigidus	Wild Lilac
Ceratostigma plumbagonoides	
Chamaecyparis pisifera	False Cypress
Cleyera japonica	Cleyera
Citrus spp.	Citrus species*
Coreopsis spp.	Coreopsis (Calliopsis) Early Sunrise, Moonbeam
Cornus florida	Flowering Dogwood
C. Stolonifera	American Dogwood
Cortaderia seloana	Pampas Grass
Cotoneaster apiculatus	Cranberry Cotoneaster
C. buxifolius	Cotoneaster
C. dammeri	Bearberry Cotoneaster
C. Microphyllus	Rockspray Cotoneaster
Cretaegus spp.	Hawthorne
Crocsmia spp.	Lucifer
Cupressus sempervirens	Italian Cypress
Delosperma alba	White Trailing Ice Plant
Delosperma spp.	Cooperi Pink
Delphinium spp.	Larkspur: Blue Elf
Dianthus deltoides	Dianthus: Maiden Pinks "Zing"
D. gratianopolitanus	Cheddar Pink
Dodonea viscosa	Hop Bush
Echinacea purpurea	Coneflower, Purple; Magnus
Elaeagnus pungens	Silverberry
Euonymus fortunei	Wintercreeper
E. japonica	Evergreen Euonymus
E. kiautschovica	Spreading Euonymus
Fatsia japonica	Japanese Aralia
Forsythia intermedia	Border Forsythia
Forsythia suspensa	Weeping Forsythia
F. viridissima	Greenstem Forsythia
Gaillardia spp.	Gaillardia, Blanket Flower: "Goblin"
Gardenia jasminoides	Gardenia, Cape-Jasmine
Gaura spp.	
Gentiana dahurica	Gentian
Geranium cinereum	Cranesbill
Gladiolus spp.	Gladiolus species**
Gypsophilia repens	Baby's Breath
Hedera helix	English Ivy
Hellanthemum spp.	Sunrose
Hemerocallis spp.	Daylily: Aztec Gold, Stella de Oro, Tender Love
Hibiscus	Rose of Sharon**
Hibiscus spp.	Mallow: Disco Belle White
Hibiscus rosa-sinensis	Chinese Hibiscus
Hosta plantaginea	Hosta, Plantain Lily (Fragrant)
H. Sieboldiana	Hosta, "Seersucker"
Houttuynia cordata var. variegata	
Hydrangea macrophylla	Bigleaf Hydrangea
Ilex Bernyi	Holly
Ilex cornuta	Chinese Holly
I. crenata	Japanese Holly



<i>I. opaca</i>	American Holly
<i>I. vomitoria</i>	Yaupon Holly, Schillings
<i>Inula ensifolia</i>	
<i>Iris ensata</i>	Sword-Leaved Iris; Jodlesong
<i>Iris siberica</i>	Siberian Iris; Cabernet
<i>Iris</i> spp.	<i>Iris</i> species**
<i>Jasminium nudiflorum</i>	Winter Jasmine
<i>Juniperus chinensis</i>	Chinese Juniper; Nick's Compact, Parsonnii
<i>J. conferta</i>	Shore Juniper; Blue Pacific
<i>J. davurica</i>	Parsoni
<i>J. horizontalis</i>	Creeping Juniper
<i>Juglans</i> spp.	Walnut*
<i>Justicia brandegeana</i>	Shrimp Plant
<i>Lagerstromia indica</i>	Crape Myrtle
<i>Lagerstromia indica</i> X <i>Fauriei</i>	Crape Myrtle; Tuscarora
<i>Lantana montevidensis</i>	Weeping Lantana
Lavender spp.	Lavender; Munstead
<i>Leonopodium alpinum</i>	Edelweiss
<i>Ligustrum amurense</i>	Amur Privet
<i>Ligustrum sinense</i>	Chinese Privet; Variegata
<i>L. japonicum</i>	Japanese Privet
<i>L. lucidum</i>	Glossy Privet (Waxleaf)
<i>Lilium</i> spp.	Lily; Jazz
<i>Liriope muscari</i>	Big Blue Lily, Silver Mound, Evergreen Giant
<i>Liriope muscari</i> var. <i>variegata</i>	<i>Liriope</i> , Variegated
<i>Liriope spicata</i>	<i>Liriope</i> , Creeping
<i>Lobelia Cardinalis</i>	Cardinal Flower, Indian Pink
<i>Lonicera japonica</i>	Japanese Honeysuckle
<i>L. tatarica</i>	Tatarian Honeysuckle
<i>Lorapetalum chinense</i>	Burgundy
<i>Lythrum</i> spp.	Loosestrife; Modern Pink
<i>Magnolia</i> spp.	<i>Magnolia</i> species**
<i>Maleophora luteola</i>	Ice Plant
<i>Malus</i> spp.	Crabapple*
<i>Miscanthus sinensis</i>	Yaku Jima, Silberfeder
<i>Nandina domestica</i>	Heavenly Bamboo
<i>Narcissus</i> spp.	<i>Narcissus</i> species**
<i>Nerium oleander</i>	Oleander
<i>Olea europaea</i>	Olive*
<i>Ophiopogon japonicus</i>	Mondo Grass
<i>Osmanthus heterophyllus</i>	<i>Osmanthus</i> (False Holly); Gulf Tide
<i>Osteospermum fruticosum</i>	Trailing African Daisy
<i>Oxydendrum arboretum</i>	Sourwood
<i>Paeonia suffruticosa</i>	Tree Peony
<i>Pennisetum setaceum</i>	Fountain Grass (Dwarf)
<i>Perovskia atriplicifolia</i>	
<i>Persea americana</i>	Avocado*
<i>Photinia Fraseri</i>	Fraser's Photinia (Redtip)
<i>Physotegia Virginiana</i>	Dragonhead, False; Vivid
<i>Picea</i> spp.	Spruce species** (Colorado Blue, Norway, etc.)
<i>Pieris japonica</i>	Lily-of-the-Valley Bush
<i>Pinus brutia</i>	Calabrian Pine
<i>P. canariensis</i>	Canary Island Pine
<i>P. Elliottii</i>	Slash Pine
<i>P. halepensis</i>	Aleppo Pine
<i>P. nigra</i>	Austrian Pine

*P. palustris*  
*P. radiata*  
*P. strobus*  
*P. sylvestris*  
*P. taeda*  
*P. Thunbergiana*  
*P. Virginiana*  
 Pistachio spp.  
*Pittosporum rhombifolium*  
*P. tobira*  
*Podocarpus macrophyllus*  
*Prunus laurocerasus*  
*Prunus* spp.

*Pseudotsuga menziesii*  
*Pyracantha coccinea*  
*P. fortuneana*  
*P. koidzumii*  
*Pyrus* spp.  
*Quercus Shumardii*  
*Quercus* spp.  
*Raphiolepis indica*  
*Raphiolepis umbellata*  
*Rhododendron*  
 (including *Azalea*)

*Rosa banksiae*  
*Rosmarinus officinalis*  
*Rudbeckia* spp.  
*Santolina virens*  
*Saxifraga* spp.  
*Scabiosa* spp.  
*Sedum album*  
*Sedum caudicicola*  
*Sedum dasyphyllum*  
*Sedum spurium*  
*Spiraea bumalda*  
*Syzygium paniculatum*  
*Taxus cuspidata*  
*T. media*  
*Teucrium* spp.  
*Thalictrum dipterocarpum*  
*Thuja occidentalis*  
*Trachelospermum asiaticum*  
*Tsuga canadensis*  
*Tulipa* spp.  
*Veronica* spp.  
*Viburnum japonicum*  
*V. odoratissimum*  
*V. plicatum*  
*V. rigidum*  
*V. suspensum*  
*V. tinus*  
*V. trilobium*  
*V. wrightii*

Longleaf Pine  
 Monterey Pine  
 Eastern White Pine  
 Scotch Pine  
 Loblolly Pine  
 Japanese Black Pine  
 Virginia Pine  
 Pistachio\*  
 Queensland Pittosporum  
 Japanese Pittosporum  
 Japanese Yew  
 English Laurel  
 Almond, Apricot, Nectarine, Peach, Plum and Prune\*  
 Douglas Fir\*\*  
 Firethorn (Scarlet)  
 Firethorn  
 Firethorn  
 Bradford Pear spp.  
 Oak, Shumard's Red  
 Oak species  
 Indian Hawthorne  
 Yedda Hawthorne  
 'Coral Bells,' 'Formosa,' 'Hino-crimson,' 'PJM'  
 'Delaware Valley White', 'Roseum Elegans,'  
 'Flame Creeper', 'Girard Crimson', 'Wakeiebisu'  
 'White Gumbo', 'George L. Tabor'  
 Lady Bank's Rose  
 Rosemary\*  
 Black-Eyed Susan; Goldsturm  
 Santolina  
 Saxifrage; Purple Dome  
 Pincushion Flower  
 Stonecrop  
 Stonecrop; Lidakense  
 Stonecrop  
 Stonecrop; Dragon's Blood  
 Spirea; Anthony Waterer  
 Australian Brushcherry  
 Japanese Yew  
 Yew  
 Germander  
 Meadow Rue  
 American Arborvitae  
 Star Jasmine  
 Canada Hemlock  
 Tulip species  
 Veronica, Speedwell; Sunny Border  
 Japanese Viburnum  
 Sweet Viburnum  
 Japanese Snowball  
 Canary Island Viburnum  
 Arrowwood Viburnum  
 Laurustinus  
 Cranberry Bush  
 Leatherleaf Viburnum

Vinca major  
Vinca Minor  
Vitis spp.  
Weigela florida  
Yucca aloifolia  
Y. filametosa

Greater Periwinkle, Vinca  
Common Periwinkle, Vinca  
Grape\*  
Old Fashioned Weigela  
Spanish Bayonet, Yucca  
Adam's Needle, Yucca

\*Do not use on food producing trees, vines or plants.

\*\*Not for use on container grown plants.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in original container in a locked storage area out of reach of children and domestic animals.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, say out of smoke

#### CONDITIONS OF SALE AND WARRANTY

The DIRECTIONS FOR USE of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop, injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of Control Solutions, Inc. or the Seller. All such risks shall be assumed by the Buyer.

Control Solutions, Inc. warrants that this product conforms to the chemical description on the label, and is reasonably fit for the purposes set forth in the DIRECTIONS FOR USE, when it is used in accordance with such directions, subject to the inherent risks mentioned above.

CONTROL SOLUTIONS, INC. NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESSED OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. THIS WARRANTY EXTENDS TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNING, OR CAUTIONS. BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. CONTROL SOLUTIONS, INC. and the seller offer this product, and the Buyer and User accept it, subject to the foregoing CONDITIONS OF SALE AND WARRANTY.