

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY Washington, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

February 22, 2013

Lisa Adamson Control Solutions, Inc. 5903 Genoa-Red Bluff Pasadena, TX 77507-1041

Subject:

Notification per PR Notice 2007-4 (container handling)

Prodiamine 0.37 Plus EPA Reg. No. 53883-172

Application Dated February 1, 2013

Dear Ms. Adamson:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 for the subject product. The Registration Division has conducted a review of this request and finds that the action falls within the scope of PRN 2007-4. The label submitted with the application has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on <u>non-refillable</u> containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please contact Mindy Ondish at (703)605-0723 or at ondish.mindy@epa.gov.

Sincerely,

Kathryn V. Montague Product Manager 23 Herbicide Branch

Registration Division (7505P)

Mindy andh, for

lease read instructions on reverse before completing form.							Form A	proved. ON		
OFDA United States				3		Regis	stration	OPP Ide	ntifier h	Number
Environmental Protection Ager				icy		Amer	ndment			
1200 Pennsylvania Avenue, N.W.					X	Other	r			
Washington, DC 20460-0001				,						·
		pplication								-,.
Company/Product Numb	53883-172		2.	EPA Prod		_{mager} ontague		Proposed C	assinc	ation
4. Company/ Product (Nam			PN	VI #	yn m	ontague	×	None		Restricted
	diamine 0.37 Plu	S		23				11001110100		
5. Name and Address of Ap	opticant (<i>Include ZIP C</i>	ode)								RA Section
Control Solutions, Inc.	•			3(c)(3)(b)(i) my product is similar or identical in composition and						
5903 Genoa-Red Bluff	.44		lac	labeling to: NOTIFICATION						
Pasadena, TX 77507-10	141		EF	PA Reg. N	No.			FFD A		346
☐ Che	eck if this is a new add	race ·						FEB 2	2 21	JI3
	ick ii lilla ia a new add			roduct Na	me					<u></u>
			Section -	 _						
Amendment - Explain				J Final pri	nted la	bels in res	ponse to Ag	ency letter da	ated _	
Resubmission in respo	nse to Agency letter d	ated	_ 🗆	J *Me Too	* Appli	cation.				
Notification - Explain b	elow.			Other -	Explair	n below.	· · ···			
Explanation: Use add										
Notification of label ch										
and the requirements										
changes have been m a violation of 18 U.S.C										
label is not consistent										
may be in violation of I										
FIFRA.										
		;	Section -	<u> </u>						
Material This Product V Child Resistant Packaging	Vill Be Packaged In: Unit Packaging		Water Solu	uhia Packa	ging		2 Type of	f Container		
Yes*	Yes		Yes Yes	ubic i auna	91119		☐ Meta			
							x Plast	tic		
X No *Certification must	X No If "Yes"	No. Per	X No		No	Per	☐ Glas			
be submitted	Unit Packaging Wt.	Container	Packaging	y Wt.		ntainer	x Pape	er r (Specify): _		
Location of Net Contents	Information	4. Size(s) Reta	il Containes		<u> </u>	1 = 1		abel Direction		
3. Location of Net Contents	пиоппации	4. Size(s) Rela	a Container			1		acei Directivi	15	
X Label	X Container		1 pt to 50	lb		lâ	On Label			
A Label							On Labeling	Accompany	ing Pro	oduct
6. Manner in Which Label is	Affixed to Product	ΧL	ithograph			stenciled				
		ΧP	aper glued			Other				
		1	Section -	- IV						
1. Contact Point (Comp	lete items directly be		ation of indiv	vidual to b	e cont					lication).
Name		Title				Tele		nclude Area		
Lisa Adam	ison	Ren	ulatorv ∆e	sistant			2	81-892-2	JUU	
Lisa Adamson Regulatory Assistant 6. Date Application										
certify that the statements I have made on this form and all attachments thereto are true Received										
accurate and complete. I acknowledge that any knowingly false or misleading statement may be										
punishable by fine or it	mprisonment or bo							_		
2. Signature	1 .	3. Ti	lle							
/ Sura C	Typa Adamson					Regulatory Assistant				
4. Typed Name 5. Date				izeguiatory Assistant						
				2-22	2 <mark>-20</mark> 13					
	Adamson									i
EPA Form 8570-1 (Rev. 8-9	4) Previous editions a	re obsolete.								



February 1, 2013

US Environmental Protection Agency Document Processing Desk (NOTIF) Office of Pesticide Programs (7504P) Room S4900, 4th Floor One Potomac Yard 2777 S. Crystal Drive Arlington, VA 22202

Attention: Ms. Kathryn Montague (PM #23)

Re: Notification per PR Notice 2007-4 Prodiamine .37 Plus (EPA Reg No. 53883-172)

Dear Ms. Montague:

On behalf of Control Solutions, Inc. please find enclosed an Application for Pesticide Notification to Re-Add Storage and Disposal language per PR Notice 2007-4. It was previously accepted by the EPA 02-21-2009 but was not included in the EPA accepted label dated 11-15-2012.

The following documents are enclosed to process this Notification:

- I. Application for Pesticide Notification (EPA Form 8570-1)
- II. One (1) copy of the proposed label with the additions highlighted in yellow.
- III. copies of proposed label clean

Should you have any questions, or wish to reach me, please feel free to contact me at 281-892-2522.

Sincerely,

Lisa Adamson

Regulatory Assistant

Lisa Adamsa

PRODIAMINE 0.37 PLUS

[with Lawn Fertilizer]

NOTIFICATION

FEB 2 2 2013

Optional Marketing Language:

[FOR PRE-EMERGENCE CONTROL OF GRASS AND BROADLEAF WEEDS IN LAWNS AND ORNAMENTAL TURF(excluding golf course putting greens)]

[LANDSCAPE ORNAMENTALS, ESTABLISHED PERENNIALS AND WILDFLOWER PLANTINGS]

[STOPS MANY WEED GRASSES BEFORE THEY GROW PREVENTS CRABGRASS, ANNUAL BLUEGRASS, FOXTAIL]

[Prevents Crabgrass in Lawns]
[Phosphorus-Free][Helping our water sources]
[With Lawn Fertilizer]
[Controls crabgrass and broadleaf weeds before they germinate]
[Feeds for 3 Months guaranteed!]
[Prevent crabgrass and fertilize your lawn in one application]
[Up to 5 month weed and grass control]
[Covers 5,000 sq. ft.] for 15 lb. bag

ACTIVE INGREDIENT:

Prodiamine: (CAS # 29091-21-2)	0.37%
INERT INGREDIENTS:	
TOTAL:	100 00%

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for further precautions and First Aid information

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals.

EPA REG NO. 53883-172

EPA EST. NOFEGGG

Control Solutions, Inc. 5903 Genoa-Red Bluff Rd. Pasadena, TX 77507-1041 281-892-2500

1	NET	CON	١T	Е١	V٦	:	•	

5/13

FIRST AID						
IF IN EYES:	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. 					
 Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. 						
	 Call a poison control center or doctor for treatment advice. 					
IF ON SKIN OR CLOTHING:	Take off contaminated clothing.					
	 Rinse skin immediately with plenty of water for 15-20 minutes. 					
	Call a poison control center or doctor for treatment advice.					
IF SWALLOWED:	 Call a poison control center or physician immediately for treatment advice. 					
	 Have person sip a glass of water if able to swallow. 					
	 Do not induce vomiting unless told to do so by the poison control center or physician. 					
.	 Do not give anything by mouth to an unconscious person. 					
IF INHALED:	Move person to fresh air.					
	 If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. 					
	Call a poison control center or doctor for further treatment advice. HOT LINE NUMBER					

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall® International (866) 897-8050 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

DIRECTION FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product is a pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in established lawns and ornamental turf (excluding golf course putting greens),


landscape ornamentals and established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seeds, germination and root development. When used according to directions, it will control; barnyardgrass; bluegrass; annual (*Poa annua*); carpetweed; chickweed, common; chickweed, mouseear (from seed); crabgrass (large, smooth); crowfootgrass; cupgrass, woolly; foxtails, annual; goosegrass¹; henbit; itchgrass; johnsongrass (from seed); junglerice; knotweed; kochia; lambsquarter, common; lovegrass; panicum (Texas, fall, browntop); pigweed; purslane, common; pusley, Florida; rescuegrass³, shepherd's purse², signalgrass, broadleaf; speedwell, Persian; sprangletop; spurge, prostrate; witchgrass; woodsorrel, yellow (from seed).

¹In many areas a single application of 3.2 to 9.3 pounds/1,000 square feet (144 to 405 pounds/acre) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 3.2 to 6.2 pounds/1,000 square feet (144 to 270 pounds/acre), followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see "Minimum and Maximum Application Rate by Turfgrass Species" table in the Rates of Application section of this label).

²For this weed, apply in late summer, fall or winter prior to germination.

WHEN TO APPLY TO ESTABLISHED TURF: This product must be applied prior to germination of weed seeds, as it will not control established weeds.

THIS MAP SHOWS APPROXIMATE CRABGRASS GERMINATION DATES



APPLICATION DIRECTIONS: Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days application.

use precautions: The following precautions apply to the use of this product in turf grasses, and lawns. Application of this product may thin emerged annual bluegrass and newly overseeded grasses. Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil).

Do not apply newly set sod until the following year. Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural

³Suppression only.

practices (such as core aeration) may result in reduced weed control. Do not apply to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species. Do not apply through any type of irrigation system. Do not apply aerially. Do not graze or feed livestock forage cut from treated areas.

RATES OF APPLICATION: This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not exceed the maximum yearly rate listed:

MINIMUM AND MAXIMUM APPLICATION RATES (by Turf Grass Species)							
Turf Species	Minimum Rate	Maximum Annual Rate ¹					
Bahiagrass, Bermudagrass ²	2.7 4 lbs/1,000 sq. ft.	9.3 lbs/1,000 sq. ft. per year					
Centipedegrass, Zoysiagrass,	(118 lbs/acre)	(405 lbs/acre per year)					
Seashore Paspalum,							
St. Augustinegrass							
Tall Fescue (including turf-		·					
Туре)							
Buffalograss	2.7 4 lbs/1,000 sq. ft.	6.2 lbs/1,000 sq. ft. per year					
Kentucky Bluegrass	(118 lbs/acre)	(270 lbs/acre per year)					
Perennial Ryegrass		·					
Creeping Red Fescue	2.7 4 lbs/1,000 sq. ft.	4.6 lbs/1,000 sq. ft. per year					
	(118 lbs/acre)	(202 lbs/acre per year)					
Creeping Bentgrass	2.7 4 lbs/1,000 sq. ft.	4.0 lbs/1,000 sq. ft. per year					
	(118 lbs/acre)	(175 lbs/acre per year)					

¹These are the maximum rates per calendar year.

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control.

WHEN TO OVERSEED AFTER APPLICATION: This product will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/reseeding results. (See map above.)

USE RATE	MONTHS BEFORE OVERSEEDING.				
	North	Transition	South '…'		
3 pounds/1,000 sq. ft. (135 pounds/acre)	4	4	. 4 .		
4 pounds/1,000 sq. ft. (175 pounds/acre)	5	4	4		
4.6 pounds/1,000 sq. ft. (202 pounds/acre)	6	5 6666	<u>5</u> 5		
4.9 pounds/1.000 sq. ft. (215 pounds/acre)	-	6 ((6، ﴿ ذُوْرِ		
6.2 pounds/1,000 sq. ft. (270 pounds/acre)	-	7	. 7		
7 pounds/1,000 sq. ft. (308 pounds/acre)	-		9 ,		
8 pounds/1,000 sq. ft. (350 pounds/acre)	-	-	16,000		
9.3 pounds/1,000 sq. ft. (405 pounds/acre)	-	_	12, ,		

LANDSCAPE ORNAMENTALS: (including established perennials and wildflower plantings): This product may be applied for residual pre-emergence weed control in ornamentals. Apply at

²May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 3.3 pounds/1,000 square feet (145 pounds/acre). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.

the rate of 2.7 to 9.3 pounds/1,000 square feet (118 to 405 pounds/acre) in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 9.3 pounds/1,000 square feet (405 pounds/acre) per year.

SPREADER SETTINGS to apply at the listed pounds per 1,000 square feet								
	3 lbs.	4 lbs.	4.6 lbs.	6.2 lbs.	8 lbs.	9.3 lbs.		
Spreader A	Χ .	X	X	X	X	X		
(May contain additional spreader brands/models, spreader settings and spreader use instructions								
as appropriate for	the product)			_				

APPLICATION TIMING AND INFORMATION: This product may be applied to newly transplanted and established ornamentals as a broadcast, over the top or directed application. Around new transplants, delay application to allow soil to settle and water thoroughly before applying. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union. This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when applied to soil free of clods, weeds and debris such as leaves. Prior to application, control existing vegetation by hand weeding, cultivation or the use of an appropriate post-emergence herbicide.

TOLERANT ORNAMENTAL SPECIES: This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

SCIENTIFIC NAME

Abies spp.

Abelia grandflora

Acer palmatum

A. Platanoides

Achillea spp.

Actinidia chinensis

Agapanthus africanus

Agapanthus orientalis

Akebia quintata

Allium cemuum

Anemone hybrida

Aquilegia spp.

Arctostaphylos densiflora

Arctotheca calendula

Artemesia spp.

Aster spp.

Aster X Frikartii

Athrium Filiz-femina

Aucuba japonica

Begonia spp.

Berberis gladwynensis

B. Julianae

B. mentorensis

B. Thunbergii

B. Verruculosa

COMMON NAME

Fir species** (Balsam, Frasier, Noble, etc.)

Abelia: Sherwood

Japanese Maple

Norway Maple

Yarrow: King Edward

Kiwi*

Lily-of-the-Nile (African Lily)

Five-leaf or Chocolate Vine

Lady's Leek, Nodding Onion

Japanese Anenome

Aquilegia: Red and Gold

Vine Hill Manzanita

Cape Weed

Wormwood; Silver Mound, Castle

Aster: Bonnie Blue, Purple Dome.ccc

Lady Fern; Fern Lady Japanese Aucuba

Fibrous Begonia: Hardy Grandis

Barberry

Wintergreen Barberry

Mentor Barberry

Japanese Barberry

Warty Barberry

Bergenia cordifolia
Boltonia asteroides
Buddleia davidii
Buxus microphylla
Callistemon viminalis
Calluna vulgaris
Campanula carpatica
Campis X Tagliabuana
Carpobrotus edulis
Cassia artemisoides
Ceanothus rigidus

Ceratostigma plumbagonoides

Chamaecyparis pisifera

Cleyera japonica
Citrus spp.
Coreopsis spp.
Cornus florida
C. Stolonifera
Cortaderia selloana
Cotoneaster apiculatus

C. buxifolius
C. dammeri
C. Microphyllus
Cretaegus spp.
Crocosmia spp.

Cupressus sempervirens

Delosperma alba
Delosperma spp.
Delphinium spp.
Dianthus deltoides
D. gratianopolitanus
Dodonea viscosa
Echinacea purpurea
Elaeagnus pungens
Euonymus fortunei

E. japonica
E. kiautschovica
Fatsia japonica
Forsythia intermedia
Forsythia suspensa
F. viridissima

Gardenia jasminoides

Gaura spp.

Gaillardia spp.

Gentiana dahurica Geranium cinereum Gladiolus spp. Gypsophilia repens Hedera helix

Hellanthemum spp. Hemerocallis spp.

Hibiscus Hibiscus spp.

Hibiscus rosa-sinensis Hosta plantaginea H. Sieboldiana Snowbank

Butterfly-Bush (Dwarf Blue); Royal Red

Japanese Boxwood Weeping Bottlebrush Scotch Heather

Tussock Bellflower, (White Clips) Trumpet Flower, Madame Galen

Hottentot Fig (Ice Plant) Feathery Cassia

Wild Lilac

False Cypress

Cleyera

Citrus species*

Coreopsis (Calliopsis) Early Sunrise, Moonbeam

Flowering Dogwood American Dogwood Pampas Grass

Cranberry Cotoneaster

Cotoneaster

Bearberry Cotoneaster Rockspray Cotoneaster

Hawthorne Lucifer

Italian Cypress

White Trailing Ice Plant

Cooperi Pink Larkspur: Blue Elf

Dianthus: Maiden Pinks "Zing"

Cheddar Pink Hop Bush

Coneflower, Purple; Magnus

Silverberry Wintercreeper Evergreen Euonymus

Spreading Euonymus
Japanese Aralia
Border Forsythia
Weeping Forsythia
Greenstem Forsythia

Gaillardia, Blanket Flower: "Goblin"

Gardenia, Cape-Jasmine

Gentian Cranesbill

Gladiolus species*

Baby's Breath English Ivy Sunrose

Daylily: Aztec Gold, Stella de Oro, Tender Love

Rose of Sharon**

Mallow: Disco Belle White

Chinese Hibiscus

Hosta, Plantain Lily (Fragrant)

Hosta, "Seersucker"

Houttuynia cordata var. variegata

Hydrangea macrophylla

Ilex Bernyi Ilex cornuta I. crenata I. opaca I. vomitoria Inula ensifolia Iris ensata Iris siberica Iris spp.

Jasminium nudiflorum Juniperus chinensis

J. conferta J. davurica J. horizontalis Juglans spp.

Justicia brandegeana Lagerstromia indica

Lagerstromia indica X Fauriei Lantana montevidensis

Lavender spp.

Leonopodium alpinum Ligustrum amurense Ligustrum sinense L. japonicum L. lucidum Lilium spp. Liriope muscari

Liriope muscari var. variegata

Liriope spicata Lobelia Cardinalis Lonicera japonica

L. tatarica

Lorapetalum chinense

Lythrum spp. Magnolia spp. Maleophora luteola

Malus spp.

Miscanthus sinensis Nandina domestica Narcissus spp. Nerium oleander Olea europaea

Ophiopogon japonicus
Omanthus heterphyllus
Osteospermum fruticosum
Oxydendrum arboretum
Paeonia suffruticosa
Pennisetum setaceumi
Perovskia atriplicifolia
Persea americana
Photinia Fraseri
Physotegia Virginiana

Picea spp. Pieris japonica Bigleaf Hydrangea

Holly

Chinese Holly Japanese Holly American Holly

Yaupon Holly, Schillings

Sword-Leaved Iris; Jodlesong

Siberian Iris; Cabernet

Iris species** Winter Jasmine

Chinese Juniper; Nick's Compact, Parsonnii

Shore Juniper; Blue Pacific

Parsoni

Creeping Juniper

Walnut* Shrimp Plant Crape Myrtle

Crape Myrtle; Tuscarora Weeping Lantana Lavender; Munstead

Edelweiss Amur Privet

Chinese Privet; Variegata

Japanese Privet

Glossy Privet (Waxleaf)

Lily; Jazz

Big Blue Lily, Silver Mound, Evergreen Giant

Liriope, Variegated Liriope, Creeping

Cardinal Flower, Indian Pink Japanese Honeysuckle Tatarian Honeysuckle

Burgundy

Loosestrife; Modern Pink Magnolia species**

Ice Plant Crabapple*

Yaku Jima, Silberfeder Heavenly Bamboo Narcissus species**

Oleander Olive* Mondo Grass

O U (F.)

Osmanthus (False Holly); Gulf Tide

Trailing African Daisy Sourwood

Tree Peony

Fountain Grass (Dwarf)

Avocado*

Frasier's Photinia (Redtip) Dragonhead, False; Vivid

Spruce species** (Colorado Blue, Norway, etc.)

Lily-of-the-Valley Bush

Pinus brutia
P. canariensis
P. Elliottii
P. halepensis
P. nigra
P. palustrus
P. radiata
P. strobus
P. sylvestris
P, taeda
P. Thunbergiana
P. Virginiana
Pistachio spp.

Pittosporum rhombifolium

P. tobira

Podocarpus macrophyllus Prunus laurocerasus

Prunus spp.

Pseudotsuga menziesii Pyracantha coccinea

P. fortuneana P. koidzumii Pyrus spp.

Quercus Shumardii Quercus spp. Raphiolepsis indica Raphiolepsis umbellata

Rhododendron (including Azalea)

Rosa banksiae Rosmarinus officinalis

Rudbeckia spp.
Santolina virens
Saxifraga spp.
Scabiosa spp.
Sedum album
Sedum cauticola
Sedum dasyphyllum
Sedum spurium
Spiraea bumalda
Syzygium paniculatum

Taxus cuspidata T. media

Teucrium spp.

Thalictrum diptercarpum

Thuja occidentalis

Trachelospermum asiatum

Tsuga canadensis

Tulipa spp.
Veronica spp.
Viburnum japonicum
V. odoratissimum
V. plicatum

Calabrian Pine Canary Island Pine

Slash Pine Aleppo Pine Austrian Pine Longleaf Pine Monterey Pine Eastern White Pine Scotch Pine

Loblolly Pine

Japanese Black Pine

Virginia Pine Pistachio*

Queensland Pittosporum Japanese Pittosporum

Japanese Yew English Laurel

Almond, Apricot, Nectarine, Peach, Plum and

Prune*
Douglas Fir**
Firethorn (Scarlet)

Firethorn Firethorn

Bradford Pear spp. Oak, Shumard's Red

Oak species Indian Hawthorne Yedda Hawthorne

'Coral Bells,' 'Formosa,' 'Hino-crimson,' 'PJM' 'Delaware Valley White', 'Roseum Elegans,' 'Flame Creeper', 'Girard Crimson', 'Wakeiebisu

'White Gumbo', 'George L. Tabor'

Lady Bank's Rose

Rosemary*

Black-Eyed Susan; Goldsturm

Santolina

Saxifrage; Purple Dome Pincushion Flower

Stonecrop

Stonecrop; Lidakense

Stonecrop

Stonecrop; Dragon's Blood Spirea; Anthony Waterer Australian Brushcherry

Japanese Yew

Yew Germander Meadow Rue

American Arborvitae Star Jasmine

Canada Hemlock
Tulip species

Veronica, Speedwell; Sunny Border

Japanese Viburnum Sweet Viburnum Japanese Snowball

02-01-2013 Label notification to add Storage and Disposal language per PR Notice 2007-4

V. rigidum V. suspensum V. tinus

V. trilobium
V. wrightii
Vinca major
Vinca Minor

Vitis spp. Weigela florida Yucca aloifolia

Y. filametosa

Canary Island Viburnum Arrowood Viburnum

Laurustinus Cranberry Bush Leatherleaf Viburnum Greater Periwinkle, Vinca Common Periwinkle, Vinca

Grape*

Old Fashioned Weigela Spanish Bayonet, Yucca Adam's Needle, Yucca

*Do not use on food producing trees, vines or plants.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a dry place. Store in locked area out of reach of children and dometic animals.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spilled bait, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance.

CONTAINER Disposal: Non-refillable container. Do not reuse or refill this container. Empty container through use according to label instructions. Do not rinse unless required for recycling. Offer for recycling, if available. If recycling Is not available; then dispose of container in a sanitary landfill or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONDITIONS OF SALE AND WARRANTY

The DIRECTIONS FOR USE of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop, injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of Control Solutions, Inc. or the Seller. All such risks shall be assumed by the Buyer.

Control Solutions, Inc. warrants that this product conforms to the chemical description on the label, and is reasonably fit for the purposes set forth in the DIRECTIONS FOR USE, when it is used in accordance with such directions, subject to the inherent risks mentioned above.

CONTROL SOLUTIONS, INC. NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESSED OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. THIS WARRANTY EXTENDS TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNING, OR, CAUTIONS, BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER

^{**}Not for use on container grown plants.

02-01-2013 Label notification to add Storage and Disposal language per PR Notice 2007-4

OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. CONTROL SOLUTIONS, INC. and the seller offer this product, and the Buyer and User accept it, subject to the foregoing CONDITIONS OF SALE AND WARRANTY.