

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

July 20, 2015

Lisa Adamson Regulatory Group Leader Control Solutions Inc. 5903 Genoa Red Bluff Pasadena, TX 77507-1041

Subject: Notification per PRN 98-10 – Addition of State Specific Language Product Name: Prodiamine 0.37 Plus EPA Registration Number: 53883-172 Application Date: June 11, 2015 Decision Number: 506198

Dear Ms. Adamson:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please contact Terri Stowe by phone at (703) 305-6117, or via email at <u>stowe.terri@epa.gov</u>.

Sincerely,

aytryn V. W Tontagues

Kathryn V. Montague Product Manager 23 Herbicide Branch Registration Division (7505P) Office of Pesticide Programs

Enclosure

NOTIFICATION

53883-172

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

07/20/2015

6-11-2015 Notification to EPA - Add state restrictive language

PRODIAMINE 0.37 PLUS

[with Lawn Fertilizer]

Optional Marketing Language:

[FOR PRE-EMERGENCE CONTROL OF GRASS AND BROADLEAF WEEDS IN LAWNS AND ORNAMENTAL TURF(excluding golf course putting greens)]

[LANDSCAPE ORNAMENTALS, ESTABLISHED PERENNIALS AND WILDFLOWER PLANTINGS]

[STOPS MANY WEED GRASSES BEFORE THEY GROW PREVENTS CRABGRASS, ANNUAL BLUEGRASS, FOXTAIL]

[Prevents Crabgrass in Lawns] [Phosphorus-Free][Helping our water sources] [With Lawn Fertilizer] [Controls crabgrass and broadleaf weeds before they germinate] [Feeds for 3 Months guaranteed!] [Prevent crabgrass and fertilize your lawn in one application] [Up to 5 month weed and grass control] [Covers 5,000 sq. ft.] for 15 lb. bag [Covers 15,000 sq ft] for 45 lb. bag [Covers 17,500 Square Feet] for 50 lb bag Prevents up to 200 Weeds Instantly activated by rain or irrigation Apply in Early Spring & Late Fall Plus Lawn Food Quick Spring Green Up after Winter Stops Crabgrass before it starts Fast Acting Formula Prevents Craborass All Season Long Inhibits weed seeds germination and root development Feeds up to 4 months guaranteed Phosphorus Free Up to 4 Month Feeding Controlled-Release Fertilizer Driven By Duration CR Engineered for Performance For a beautiful [Product name] lawn

ANNUAL FEEDING PROGRAM: IMPORTANCE OF FEED: Regularly feeding your lawn provides the necessary nutrients it needs to help fight weeds, heat, drought and traffic. EARLY SPRING FEEDING: After the first time you mow, feed to help your lawn greenup quickly after the harsh winter months. LATE SPRING FEEDING: Before the hot summer months, feed to prepare and strengthen for stresses of heat, drought and traffic. EARLY FALL FEEDING: After your lawn comes out of dormancy, feed for a quick recovery from summer stress.

LATE FALL FEEDING: Once the leaves have fallen, feed to build a strong root system and prepare your lawn for winter.

ACTIVE INGREDIENT:

Prodiamine: (CAS # 29091-21-2)
OTHER INGREDIENTS:
TOTAL:

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for further precautions and First Aid information

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals.

EPA REG NO. 53883-172

EPA EST. NO.

Control Solutions, Inc. 5903 Genoa-Red Bluff Rd. Pasadena, TX 77507-1041 281-892-2500

NET CONTENT:____

FIRST AID				
IF IN EYES:	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. 			
	 Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. 			
	Call a poison control center or doctor for treatment advice.			
IF ON SKIN OR CLOTHING:	Take off contaminated clothing.			
	Rinse skin immediately with plenty of water for 15-20 minutes.			
	• Call a poison control center or doctor for treatment advice.			
IF SWALLOWED:	 Call a poison control center or physician immediately for treatment advice. 			
	Have person sip a glass of water if able to swallow.			
	 Do not induce vomiting unless told to do so by the poison control center or physician. 			
	Do not give anything by mouth to an unconscious person.			
IF INHALED:	Move person to fresh air.			

	 If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
	 Call a poison control center or doctor for further treatment advice.
	HOT LINE NUMBER
going for treatment. You	ner or label with you when calling a poison control center or doctor, or may also contact SafetyCall [®] International (866) 897-8050 for
emergency medical treat	ment information.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

DIRECTION FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

*This [XXXX] Ib bag covers [XXXX] sq ft of lawn area. A lawn typically will utilize 1 to 4 lbs of nitrogen per 1,000 sq ft per growing season. When used as directed, this product will supplement your regular fertilizer program by providing [XXXX] Ib of nitrogen per [XXXX] sq ft. Please consult your local extension office for questions regarding a regular fertilizer program.
* This statement is only for the use and sale in the state of Florida

*Apply only to actively growing turf. Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands in the driveway, sidewalk, or street, back onto your lawn. Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices. Check with your county or city government to determine if there are local regulations for fertilizer use.

* This statement is only for the use and sale in the state of Kentucky

This product is a pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in established lawns and ornamental turf (excluding golf course putting greens), landscape ornamentals and established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seeds, germination and root development. When used according to directions, it will control; barnyardgrass; bluegrass; annual (*Poa annua*); carpetweed; chickweed, common; chickweed, mouseear (from seed); crabgrass (large, smooth); crowfootgrass; cupgrass, woolly; foxtails, annual; goosegrass¹; henbit; itchgrass; johnsongrass (from seed); junglerice; knotweed; kochia; lambsquarter, common; lovegrass; panicum (Texas, fall, browntop); pigweed; purslane, common; pusley, Florida; rescuegrass³, shepherd's purse², signalgrass, broadleaf; speedwell, Persian; sprangletop; spurge, prostrate; witchgrass; woodsorrel, yellow (from seed).

¹In many areas a single application of 4.1 to 9.3 pounds/1,000 square feet (175to 405 pounds/acre) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 4.1 to 6.2 pounds/1,000 square feet (175 to 270 pounds/acre), followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see"Minimum and Maximum Application Rate by Turfgrass Species" table in the Rates of Application section of this label).

²For this weed, apply in late summer, fall or winter prior to germination.

³Suppression only.

WHEN TO APPLY TO ESTABLISHED TURF: This product must be applied prior to germination of weed seeds, as it will not control established weeds.



THIS MAP SHOWS APPROXIMATE CRABGRASS GERMINATION DATES

APPLICATION DIRECTIONS: Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days of application.

USE PRECAUTIONS: The following precautions apply to the use of this product in turf grasses and lawns. Application of this product may thin emerged annual bluegrass and newly overseeded grasses. Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil).

Do not apply newly set sod until the following year. Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural practices (such as core aeration) may result in reduced weed control. Do not apply to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species. Do not apply through any type of irrigation system. Do not apply aerially. Do not graze or feed livestock forage cut from treated areas.

RATES OF APPLICATION: This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not exceed the maximum yearly rate listed:

MINIMUM AND MAXIMUM APPLICATION RATES (by Turf Grass Species)					
Turf Species	Minimum Rate	Maximum Annual Rate ¹			
Bahiagrass, Bermudagrass ² Centipedegrass, Zoysiagrass, Seashore Paspalum, St. Augustinegrass Tall Fescue (including turf- Type)	2.7 lbs/1,000 sq. ft. (118 lbs/acre)	9.3 lbs/1,000 sq. ft. per year (405 lbs/acre per year)			
Buffalograss	2.7 lbs/1,000 sq. ft.	6.2 lbs/1,000 sq. ft. per year			
Kentucky Bluegrass	(118 lbs/acre)	(270 lbs/acre per year)			
Perennial Ryegrass					
Creeping Red Fescue	2.7 lbs/1,000 sq. ft.	4.6 lbs/1,000 sq. ft. per year			
	(118 lbs/acre)	(202 lbs/acre per year)			
Creeping Bentgrass	2.7 lbs/1,000 sq. ft.	4.0 lbs/1,000 sq. ft. per year			
	(118 lbs/acre)	(175 lbs/acre per year)			

¹These are the maximum rates per calendar year.

²May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 3.3 pounds/1,000 square feet (145 pounds/acre). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control.

WHEN TO OVERSEED AFTER APPLICATION: This product will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/reseeding results. (See map above.)

USE RATE	MONTHS BEFORE OVERSEEDING		
	North	Transition	South
3 pounds/1,000 sq. ft. (135 pounds/acre)	4	4	4
4 pounds/1,000 sq. ft. (175 pounds/acre)	5	4	4
4.6 pounds/1,000 sq. ft. (202 pounds/acre)	6	5	5
4.9 pounds/1.000 sq. ft. (215 pounds/acre)	-	6	6
6.2 pounds/1,000 sq. ft. (270 pounds/acre)	-	7	7
7 pounds/1,000 sq. ft. (308 pounds/acre)	-	-	9
8 pounds/1,000 sq. ft. (350 pounds/acre)	-	-	10
9.3 pounds/1,000 sq. ft. (405 pounds/acre)	-	-	12

LANDSCAPE ORNAMENTALS: (including established perennials and wildflower plantings): This product may be applied for residual pre-emergence weed control in ornamentals. Apply at the rate of 2.7 to 9.3 pounds/1,000 square feet (118 to 405 pounds/acre) in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 9.3 pounds/1,000 square feet (405 pounds/acre) per year.

SPREADER SETTINGS to apply at the listed pounds per 1,000 square feet						
	3 lbs.	4 lbs.	4.6 lbs.	6.2 lbs.	8 lbs.	9.3 lbs.
Spreader A	Х	Х	Х	Х	Х	Х
(May contain additional spreader brands/models, spreader settings and spreader use instructions as appropriate for the product)						

APPLICATION TIMING AND INFORMATION: This product may be applied to newly transplanted and established ornamentals as a broadcast, over the top or directed application. Around new transplants, delay application to allow soil to settle and water thoroughly before applying. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union. This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when applied to soil free of clods, weeds and debris such as leaves. Prior to application, control existing vegetation by hand weeding, cultivation or the use of an appropriate post-emergence herbicide.

TOLERANT ORNAMENTAL SPECIES: This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

SCIENTIFIC NAME

Abies spp. Abelia grandflora Acer palmatum A. Platanoides Achillea spp. Actinidia chinensis Agapanthus africanus Agapanthus orientalis Akebia quintata Allium cemuum Anemone hybrida Aquilegia spp. Arctostaphylos densiflora Arctotheca calendula Artemesia spp. Aster spp. Aster X Frikartii Athrium Filiz-femina Aucuba japonica Begonia spp. Berberis gladwynensis B. Julianae B. mentorensis B. Thunbergii B. Verruculosa Bergenia cordifolia Boltonia asteroides Buddleia davidii Buxus microphylla Callistemon viminalis Calluna vulgaris Campanula carpatica Campis X Tagliabuana Carpobrotus edulis Cassia artemisoides

COMMON NAME

Fir species** (Balsam, Frasier, Noble, etc.) Abelia: Sherwood Japanese Maple Norway Maple Yarrow: King Edward Kiwi* Lily-of-the-Nile (African Lily) Five-leaf or Chocolate Vine Lady's Leek, Nodding Onion Japanese Anenome Aquilegia: Red and Gold Vine Hill Manzanita Cape Weed Wormwood; Silver Mound, Castle Aster: Bonnie Blue, Purple Dome Lady Fern; Fern Lady Japanese Aucuba Fibrous Begonia: Hardy Grandis Barberry Wintergreen Barberry Mentor Barberry Japanese Barberry Warty Barberry Snowbank

Butterfly-Bush (Dwarf Blue); Royal Red Japanese Boxwood Weeping Bottlebrush Scotch Heather Tussock Bellflower, (White Clips) Trumpet Flower, Madame Galen Hottentot Fig (Ice Plant) Feathery Cassia

Ceanothus rigidus Ceratostigma plumbagonoides Chamaecyparis pisifera Cleyera japonica Citrus spp. Coreopsis spp. Cornus florida C. Stolonifera Cortaderia selloana Cotoneaster apiculatus C. buxifolius C. dammeri C. Microphyllus Cretaegus spp. Crocosmia spp. Cupressus sempervirens Delosperma alba Delosperma spp. Delphinium spp. Dianthus deltoides D. gratianopolitanus Dodonea viscosa Echinacea purpurea Elaeagnus pungens Euonymus fortunei E. japonica E. kiautschovica Fatsia japonica Forsythia intermedia Forsythia suspensa F. viridissima Gaillardia spp. Gardenia jasminoides Gaura spp. Gentiana dahurica Geranium cinereum Gladiolus spp. Gypsophilia repens Hedera helix Hellanthemum spp. Hemerocallis spp. Hibiscus Hibiscus spp. Hibiscus rosa-sinensis Hosta plantaginea H. Sieboldiana Houttuynia cordata var. variegata Hydrangea macrophylla Ilex Bernyi llex cornuta I. crenata I. opaca I. vomitoria Inula ensifolia Iris ensata Iris siberica

Wild Lilac

False Cypress Cleyera Citrus species* Coreopsis (Calliopsis) Early Sunrise, Moonbeam Flowering Dogwood American Dogwood Pampas Grass Cranberry Cotoneaster Cotoneaster **Bearberry Cotoneaster** Rockspray Cotoneaster Hawthorne Lucifer Italian Cypress White Trailing Ice Plant Cooperi Pink Larkspur: Blue Elf Dianthus: Maiden Pinks "Zing" Cheddar Pink Hop Bush Coneflower, Purple; Magnus Silverberrv Wintercreeper Evergreen Euonymus Spreading Euonymus Japanese Aralia Border Forsythia Weeping Forsythia Greenstem Forsythia Gaillardia, Blanket Flower: "Goblin" Gardenia, Cape-Jasmine Gentian Cranesbill Gladiolus species** Baby's Breath English Ivy Sunrose Daylily: Aztec Gold, Stella de Oro, Tender Love Rose of Sharon** Mallow: Disco Belle White Chinese Hibiscus Hosta, Plantain Lily (Fragrant) Hosta, "Seersucker" **Bigleaf Hydrangea** Holly Chinese Holly Japanese Holly American Holly Yaupon Holly, Schillings Sword-Leaved Iris; Jodlesong Siberian Iris; Cabernet

Iris spp. Jasminium nudiflorum Juniperus chinensis J. conferta J. davurica J. horizontalis Juglans spp. Justicia brandegeana Lagerstromia indica Lagerstromia indica X Fauriei Lantana montevidensis Lavender spp. Leonopodium alpinum Ligustrum amurense Ligustrum sinense L. iaponicum L. lucidum Lilium spp. Liriope muscari Liriope muscari var. variegata Liriope spicata Lobelia Cardinalis Lonicera japonica L. tatarica Lorapetalum chinense Lythrum spp. Magnolia spp. Maleophora luteola Malus spp. Miscanthus sinensis Nandina domestica Narcissus spp. Nerium oleander Olea europaea Ophiopogon japonicus Omanthus heterphyllus Osteospermum fruticosum Oxydendrum arboretum Paeonia suffruticosa Pennisetum setaceumi Perovskia atriplicifolia Persea americana Photinia Fraseri Physotegia Virginiana Picea spp. Pieris japonica Pinus brutia P. canariensis P. Elliottii P. halepensis P. nigra P. palustrus P. radiata P. strobus P. sylvestris

P, taeda

Iris species** Winter Jasmine Chinese Juniper; Nick's Compact, Parsonnii Shore Juniper; Blue Pacific Parsoni **Creeping Juniper** Walnut* Shrimp Plant Crape Myrtle Crape Myrtle; Tuscarora Weeping Lantana Lavender; Munstead Edelweiss Amur Privet Chinese Privet; Variegata Japanese Privet Glossy Privet (Waxleaf) Lily; Jazz Big Blue Lily, Silver Mound, Evergreen Giant Liriope, Variegated Liriope, Creeping Cardinal Flower, Indian Pink Japanese Honeysuckle Tatarian Honevsuckle Burgundy Loosestrife; Modern Pink Magnolia species** Ice Plant Crabapple* Yaku Jima, Silberfeder Heavenly Bamboo Narcissus species** Oleander Olive* Mondo Grass Osmanthus (False Holly); Gulf Tide Trailing African Daisy Sourwood Tree Peony Fountain Grass (Dwarf) Avocado* Frasier's Photinia (Redtip) Dragonhead, False; Vivid Spruce species** (Colorado Blue, Norway, etc.) Lily-of-the-Valley Bush Calabrian Pine Canary Island Pine Slash Pine Aleppo Pine

Austrian Pine Longleaf Pine Monterey Pine Eastern White Pine Scotch Pine Loblolly Pine

P. Thunbergiana P. Virginiana Pistachio spp. Pittosporum rhombifolium P. tobira Podocarpus macrophyllus Prunus laurocerasus Prunus spp. Pseudotsuga menziesii Pyracantha coccinea P. fortuneana P. koidzumii Pyrus spp. Quercus Shumardii Quercus spp. Raphiolepsis indica Raphiolepsis umbellata Rhododendron (including Azalea) Rosa banksiae Rosmarinus officinalis Rudbeckia spp. Santolina virens Saxifraga spp. Scabiosa spp. Sedum album Sedum cauticola Sedum dasyphyllum Sedum spurium Spiraea bumalda Syzygium paniculatum Taxus cuspidata T. media Teucrium spp. Thalictrum diptercarpum Thuja occidentalis Trachelospermum asiatum Tsuga canadensis Tulipa spp. Veronica spp. Viburnum japonicum V. odoratissimum V. plicatum V. rigidum V. suspensum V. tinus V. trilobium V. wrightii Vinca major Vinca Minor Vitis spp. Weigela florida Yucca aloifolia

Japanese Black Pine Virginia Pine Pistachio* Queensland Pittosporum Japanese Pittosporum Japanese Yew English Laurel Almond, Apricot, Nectarine, Peach, Plum and Prune* Douglas Fir** Firethorn (Scarlet) Firethorn Firethorn Bradford Pear spp. Oak, Shumard's Red Oak species Indian Hawthorne Yedda Hawthorne 'Coral Bells,' ' Formosa,' ' Hino-crimson,' ' PJM' 'Delaware Valley White', ' Roseum Elegans,' 'Flame Creeper', ' Girard Crimson', 'Wakeiebisu 'White Gumbo', 'George L. Tabor' Lady Bank's Rose Rosemarv* Black-Eyed Susan; Goldsturm Santolina Saxifrage; Purple Dome **Pincushion Flower** Stonecrop Stonecrop; Lidakense Stonecrop Stonecrop; Dragon's Blood Spirea; Anthony Waterer Australian Brushcherry Japanese Yew Yew Germander Meadow Rue American Arborvitae Star Jasmine Canada Hemlock Tulip species Veronica, Speedwell; Sunny Border Japanese Viburnum Sweet Viburnum Japanese Snowball Canary Island Viburnum Arrowood Viburnum Laurustinus Cranberry Bush Leatherleaf Viburnum Greater Periwinkle, Vinca Common Periwinkle, Vinca Grape* Old Fashioned Weigela Spanish Bayonet, Yucca

Y. filametosa

Adam's Needle, Yucca *Do not use on food producing trees, vines or plants.

**Not for use on container grown plants.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a dry place. Store in locked area out of reach of children and domestic animals.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spilled bait, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance.

CONTAINER Disposal: Non-refillable container. Do not reuse or refill this container. Empty container through use according to label instructions. Do not rinse unless required for recycling. Offer for recycling, if available. If recycling Is not available; then dispose of container in a sanitary landfill or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONDITIONS OF SALE AND WARRANTY IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Control Solutions Inc. (CSI). All such risks shall be assumed by the user or buyer. DISCLAIMER OF WARRANTIES:

To the extent consistent with applicable law, CSI makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of CSI is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, CSI disclaims any liability what so ever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at CSI's election, the replacement of product. The DIRECTIONS FOR USE of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop, injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of Control Solutions, Inc. or the Seller. All such risks shall be assumed by the Buyer.

Control Solutions, Inc. warrants that this product conforms to the chemical description on the label, and is reasonably fit for the purposes set forth in the DIRECTIONS FOR USE, when it is used in accordance with such directions, subject to the inherent risks mentioned above.

Formatted: Centered

CONTROL SOLUTIONS, INC. NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESSED OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. THIS WARRANTY EXTENDS TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNING, OR CAUTIONS. BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. CONTROL SOLUTIONS. INC. and the seller offer this product, and the Buyer and User accept it, subject to the foregoing CONDITIONS OF SALE AND WARRANTY.

Optional State Specific Language

[This product may stain driveways, walks, patios, etc.]

[Wash or sweep off thoroughly, immediately after application]

[Do Not Apply Near Water, Storm Drains or Drainage Ditches.]

[Do Not Apply if heavy rain is expected]

[Apply this product only to your lawn or garden, and sweep any product that lands on the driveway, sidewalk or street, back onto your lawn or garden]

[Help do your part in protecting our rivers and lakes by following label directives. Do not apply near water storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk or street back onto your lawn. Do not apply to frozen ground.]

{Label may also include state specific language regarding Nitrogen application}