

53883-173

7/1/2014

1/13



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
Washington, D.C. 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

JUL -1 2014

Lisa Adamson
Control Solutions, Inc.
5903 Genoa-Red Bluff
Pasadena, TX 77507-1041

Subject: Notification per PR Notice 98-10 – Addition of marketing language
Product Name: Prodiamine 0.20 Plus
EPA Registration Number: 53883-173
Application Dated: April 29, 2014
Decision Number: 490820

Dear Ms. Adamson:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the subject product. Our review finds that the action falls within the scope of PRN 98-10. The label submitted with the application has been date-stamped "Notification" and placed in our records.

If you have any questions, please contact Sarah Meadows at (703)347-0505 or at meadows.sarah@epa.gov.

Sincerely,

Kathryn V. Montague *for*
Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

3/13



**CONTROL
SOLUTIONS**
incorporated

May 28, 2014

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
US EPA
One Potomac Yard
2777 Crystal Drive, Room S-4900
Arlington, VA 22202

Re: Prodiamine 0.20 Plus EPA Reg # 53883-173
Addition of Optional Marketing Language Allowed by PRN 98-10

To Whom It May Concern:

Control Solutions Inc. (CSI) is notifying the Agency of its intention to add optional marketing language to the above reference product.

We certify the following:

This notification is consistent with the provisions of PR Notices 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

In support of this submission, the following documents are attached:

- Application for Pesticide Registration (EPA Form 8570-1)
- One Copy of annotated labeling highlighting the added text in blue (page 10)
- Three Copies of the final printed labeling

Should you have any questions about this application, please contact me at 800-242-5562 or email ladamson@controlsolutionsinc.com

Sincerely,

Lisa Adamson
Regulatory Specialist

4/13

PRODIAMINE 0.20 PLUS
FOR PRE-EMERGENCE CONTROL OF GRASS AND BROADLEAF WEEDS
IN LAWNS AND ORNAMENTAL TURF
 (excluding golf course putting greens)
LANDSCAPE ORNAMENTALS, ESTABLISHED PERENNIALS AND
WILDFLOWER PLANTINGS
STOPS MANY WEED GRASSES BEFORE THEY GROW
PREVENTS CRABGRASS, ANNUAL BLUEGRASS, FOXTAIL

ACTIVE INGREDIENT:

Prodiamine (CAS # 29091-21-2)0.20%

INERT INGREDIENTS: 99.80%

TOTAL: 100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION

See back panel for further precautions and First Aid information

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals.

NOTIFICATION

JUL -1 2014

EPA REG NO. 53883-173
EPA EST. NO

Control Solutions, Inc.
5903 Genoa-Red Bluff Rd.
Pasadena, TX 77507-1041
281-892-2500

NET CONTENT: _____

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> ● Hold eye open and rinse slowly and gently with water for 15-20 minutes. ● Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. ● Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> ● Take off contaminated clothing. ● Rinse skin immediately with plenty of water for 15-20 minutes. ● Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> ● Call a poison control center or physician immediately for treatment advice. ● Have person sip a glass of water if able to swallow. ● Do not induce vomiting unless told to do so by the poison control center or physician. ● Do not give anything by mouth to an unconscious person.
IF INHALED:	<ul style="list-style-type: none"> ● Move person to fresh air. ● If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, <u>if possible</u>. ● Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall® International (866) 897-8050 for emergency medical treatment information.</p>	

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

DIRECTION FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product is a pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in established lawns and ornamental turf (excluding golf course putting greens), landscape ornamentals and established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seeds, germination and root development. When used according to directions, it will control; barnyardgrass; bluegrass; annual (*Poa annua*); carpetweed; chickweed, common; chickweed, mouseear (from seed); crabgrass (large, smooth); crowfootgrass; cupgrass, woolly; foxtails, annual; goosegrass¹; henbit; itchgrass; johnsongrass (from seed); junglerice; knotweed; kochia; lambsquarter, common; lovegrass; panicum (Texas, fall, browntop); pigweed; purslane, common; pusley, Florida; rescuegrass³, shepherd's purse², signalgrass, broadleaf; speedwell, Persian; sprangletop; spurge, prostrate; witchgrass; woodsorrel, yellow (from seed).


¹In many areas a single application of 7.6 to 17.2 pounds/1,000 square feet (330 to 750 pounds/acre) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 7.6 to 11.5 pounds/1,000 square feet (330 to 500 pounds/acre), followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see "Minimum and Maximum Application Rate by Turfgrass Species" table in the Rates of Application section of this label).

²For this weed, apply in late summer, fall or winter prior to germination.

³Suppression only.

WHEN TO APPLY TO ESTABLISHED TURF: This product must be applied prior to germination of weed seeds, as it will not control established weeds.

THIS MAP SHOWS APPROXIMATE CRABGRASS GERMINATION DATES



APPLICATION DIRECTIONS: Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days application.

USE PRECAUTIONS: The following precautions apply to the use of this product in turf grasses and lawns. Application of this product may thin emerged annual bluegrass and newly overseeded grasses. Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil). Do not apply to newly set sod until the following year. Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural

practices (such as core aeration) may result in reduced weed control. Do not apply to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species. Do not apply through any type of irrigation system. Do not apply aerially. Do not graze or feed livestock forage cut from treated areas.

RATES OF APPLICATION: This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not exceed the maximum yearly rate listed:

MINIMUM AND MAXIMUM APPLICATION RATES (by Turf Grass Species)		
Turf Species	Minimum Rate	Maximum Annual Rate¹
Bahiagrass, Bermudagrass ² Centipedegrass, Zoysiagrass, Seashore Paspalum, St. Augustinegrass Tall Fescue (including turf-Type)	4 lbs/1,000 sq. ft. (174 lbs/acre)	17.2 lbs/1,000 sq. ft. per year (750 lbs/acre per year)
Buffalograss Kentucky Bluegrass Perennial Ryegrass	4 lbs/1,000 sq. ft. (174 lbs/acre)	11.5 lbs/1,000 sq. ft. per year (500 lbs/acre per year)
Creeping Red Fescue	4 lbs/1,000 sq. ft. (174 lbs/acre)	8.5 lbs/1,000 sq. ft. per year (370 lbs/acre per year)
Creeping Bentgrass	4 lbs/1,000 sq. ft. (174 lbs/acre)	7.6 lbs/1,000 sq. ft. per year (330 lbs/acre per year)

¹These are the maximum rates per calendar year.

²May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 6 pounds/1,000 square feet (260 pounds/acre). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control.

WHEN TO OVERSEED AFTER APPLICATION: This product will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/reseeding results. (See map above)

USE RATE	MONTHS BEFORE OVERSEEDING		
	North	Transition	South
4 pounds/1,000 sq. ft. (174 pounds/acre)	4	4	4
7.6 pounds/1,000 sq. ft. (330 pounds/acre)	5	4	4
8.5 pounds/1,000 sq. ft. (370 pounds/acre)	6	5	5
9.2 pounds/1,000 sq. ft. (400 pounds/acre)	-	6	6
11.5 pounds/1,000 sq. ft. (500 pounds/acre)	-	7	7
13 pounds/1,000 sq. ft. (570 pounds/acre)	-	-	9
14.9 pounds/1,000 sq. ft. (650 pounds/acre)	-	-	10
17.2 pounds/1,000 sq. ft. (750 pounds/acre)	-	-	12

LANDSCAPE ORNAMENTALS: (including established perennials and wildflower plantings): This product may be applied for residual pre-emergence weed control in ornamentals. Apply at the rate of 7.6 to 17.2 pounds/1,000 square feet (330 to 750 pounds/acre) in fall and/or spring.

Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 17.2 pounds/1,000 square feet (750 pounds/acre) per year.

SPREADER SETTINGS to apply at the listed pounds per 1,000 square feet						
	4 lbs.	6 lbs.	7.6 lbs.	8.5 lbs.	11.5 lbs.	17.2 lbs.
Spreader A	X	X	X	X	X	X
(May contain additional spreader brands/models, spreader settings and spreader use instructions as appropriate for the product)						

APPLICATION TIMING AND INFORMATION: This product may be applied to newly transplanted and established ornamentals a broadcast, over the top or directed application. Around new transplants, delay application to allow soil to settle and water thoroughly before applying. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union. This product is a pre-emergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when applied to soil free of clods, weeds and debris such as leaves. Prior to application, control existing vegetation by hand weeding, cultivation or the use of an appropriate post-emergence herbicide.

TOLERANT ORNAMENTAL SPECIES: This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

SCIENTIFIC NAME

Abies spp.
Abelia grandiflora
Acer palmatum
A. Platanoides
Achillea spp.
Actindia chinensis
Agapanthus africanus
Agapanthus orientalis
Akebiaq quintata
Allium cemuum
Anemone hybrida
Aquilegia spp.
Arctostaphylos densiflora
Arctotheca calendula
Artemesia spp.
Aster spp.
Aster X Frikartii
Athrium Filiz-femina
Aucuba japonica
Begonia spp.
Berberis gladwynensis
B. Julianae
B. mentorensis
B. Thunbergijl
B. Verruculosa

COMMON NAME

Fir species** (Balsam, Frasier, Noble, etc.)
Abelia: Sherwood
Japanese Maple
Norway Maple
Yarrow: King Edward
Kiwi*
Lily-of-the-Nile (African Lily)

Five-leaf or Chocolate Vine
Lady's Leek, Nodding Onion
Japanese Anemone
Aquilegia: Red and Gold
Vine Hill Manzanita
Cape Weed
Wormwood; Silver Mound, Castle
Aster: Bonnie Blue, Purple Dome

Lady Fern; Fern Lady
Japanese Aucuba
Fibrous Begonia: Hardy Grandis
Barberry
Wintergreen Barberry
Mentor Barberry
Japanese Barberry
Warty Barberry

Bergenia cordifolia
 Boltonia asteroides
 Buddleia davidii
 Buxus microphylla
 Callistemon viminalis
 Calluna vulgaris
 Campanula carpatica
 Campis X Tagliabuana
 Carpobrotus edulis
 Cassia artemisoides
 Ceanothus rigidus
 Ceratostigma plumbagonoides
 Chamaecyparis pisifera
 Cleyera japonica
 Citrus spp.
 Coreopsis spp.
 Cornus florida
 C. Stolonifera
 Cortaderia selloana
 Cotoneaster apiculatus
 C. buxifolius
 C. dammeri
 C. Microphyllus
 Cretaegus spp.
 Crocosmia spp.
 Cupressus sempervirens
 Delosperma alba
 Delosperma spp.
 Delphinium spp.
 Dianthus deltoides
 D. gratianopolitanus
 Dodonea viscosa
 Echinacea purpurea
 Elaeagnus pungens
 Euonymus fortunei
 E. japonica
 E. kiautschovica
 Fatsia japonica
 Forsythia intermedia
 Forsythia suspensa
 F. viridissima
 Gaillardia spp.
 Gardenia jasminoides
 Gaura spp.
 Gentiana dahurica
 Geranium cinereum
 Gladiolus spp.
 Gypsophila repens
 Hedera helix
 Hellanthenum spp.
 Hemerocallis spp.
 Hibiscus
 Hibiscus spp.
 Hibiscus rosa-sinensis
 Hosta plantaginea
 H. Sieboldiana

Snowbank
 Butterfly-Bush (Dwarf Blue); Royal Red
 Japanese Boxwood
 Weeping Bottlebrush
 Scotch Heather
 Tussock Bellflower, (White Clips)
 Trumpet Flower, Madame Galen
 Hottentot Fig (Ice Plant)
 Feathery Cassia
 Wild Lilac

 False Cypress
 Cleyera
 Citrus species*
 Coreopsis (Calliopsis) Early Sunrise, Moonbeam
 Flowering Dogwood
 American Dogwood
 Pampas Grass
 Cranberry Cotoneaster
 Cotoneaster
 Bearberry Cotoneaster
 Rockspray Cotoneaster
 Hawthorne
 Lucifer
 Italian Cypress
 White Trailing Ice Plant
 Cooperi Pink
 Larkspur: Blue Elf
 Dianthus: Maiden Pinks "Zing"
 Cheddar Pink
 Hop Bush
 Coneflower, Purple; Magnus
 Silverberry
 Wintercreeper
 Evergreen Euonymus
 Spreading Euonymus
 Japanese Aralia
 Border Forsythia
 Weeping Forsythia
 Greenstem Forsythia
 Gaillardia, Blanket Flower: "Goblin"
 Gardenia, Cape-Jasmine

 Gentian
 Cranesbill
 Gladiolus species**
 Baby's Breath
 English Ivy
 Sunrose
 Daylily: Aztec Gold, Stella de Oro, Tender Love
 Rose of Sharon**
 Mallow: Disco Belle White
 Chinese Hibiscus
 Hosta, Plantain Lily (Fragrant)
 Hosta, "Seersucker"

Houttuynia cordata var. variegata
 Hydrangea macrophylla
 Ilex Bernyi
 Ilex cornuta
 I. crenata
 I. opaca
 I. vomitoria
 Inula ensifolia
 Iris ensata
 Iris siberica
 Iris spp.
 Jasminium nudiflorum
 Juniperus chinensis
 J. conferta
 J. davurica
 J. horizontalis
 Juglans spp.
 Justicia brandegeana
 Lagerstromia indica
 Lagerstromia indica X Fauriei
 Lantana montevidensis
 Lavender spp.
 Leonopodium alpinum
 Ligustrum amurense
 Ligustrum sinense
 L. japonicum
 L. lucidum
 Liliium spp.
 Liriope muscari
 Liriope muscari var. variegata
 Liriope spicata
 Lobelia Cardinalis
 Lonicera japonica
 L. tatarica
 Lorapetalum chinense
 Lythrum spp.
 Magnolia spp.
 Maleophora luteola
 Malus spp.
 Miscanthus sinensis
 Nandina domestica
 Narcissus spp.
 Nerium oleander
 Olea europaea
 Ophiopogon japonicus
 Osmanthus heterophyllus
 Osteospermum fruticosum
 Oxydendrum arboreum
 Sourwood
 Paeonia suffruticosa
 Pennisetum setaceum
 Perovskia atriplicifolia
 Persea americana
 Photinia Fraseri
 Physotegia Virginiana
 Picea spp.

Bigleaf Hydrangea
 Holly
 Chinese Holly
 Japanese Holly
 American Holly
 Yaupon Holly, Schillings

 Sword-Leaved Iris; Jodlesong
 Siberian Iris; Cabernet
 Iris species**
 Winter Jasmine
 Chinese Juniper; Nick's Compact, Parsonii
 Shore Juniper; Blue Pacific
 Parsoni
 Creeping Juniper
 Walnut*
 Shrimp Plant
 Crape Myrtle
 Crape Myrtle; Tuscarora
 Weeping Lantana
 Lavender; Munstead
 Edelweiss
 Amur Privet
 Chinese Privet; Variegata
 Japanese Privet
 Glossy Privet (Waxleaf)
 Lily; Jazz
 Big Blue Lily, Silver Mound, Evergreen Giant
 Liriope, Variegated
 Liriope, Creeping
 Cardinal Flower, Indian Pink
 Japanese Honeysuckle
 Tatarian Honeysuckle
 Burgundy
 Loosestrife; Modern Pink
 Magnolia species**
 Ice Plant
 Crabapple*
 Yaku Jima, Silberfeder
 Heavenly Bamboo
 Narcissus species**
 Oleander
 Olive*
 Mondo Grass
 Osmanthus (False Holly); Gulf Tide
 Trailing African Daisy

 Tree Peony
 Fountain Grass (Dwarf)

 Avocado*
 Frasier's Photinia (Redtip)
 Dragonhead, False; Vivid
 Spruce species** (Colorado Blue, Norway, etc.)

Pieris japonica
Pinus brutia
P. canariensis
P. Elliottii
P. halepensis
P. nigra
P. palustris
P. radiata
P. strobus
P. sylvestris
P. taeda
P. Thunbergiana
P. Virginiana
 Pistachio spp.
Pittosporum rhombifolium
P. tobira
Podocarpus macrophyllus
Prunus laurocerasus
Prunus spp.

Pseudotsuga menziesii
Pyracantha coccinea
P. fortuneana
P. koidzumii
Pyrus spp.
Quercus Shumardii
Quercus spp.
Raphiolepis indica
Raphiolepis umbellata
Rhododendron
 (including *Azalea*)

Rosa banksiae
Rosmarinus officinalis
Rudbeckia spp.
Santolina virens
Saxifraga spp.
Scabiosa spp.
Sedum album
Sedum caucicola
Sedum dasyphyllum
Sedum spurium
Spiraea bumalda
Syzygium paniculatum
Taxus cuspidata
T. media
Teucrium spp.
Thalictrum diptercarpum
Thuja occidentalis
Trachelospermum asiaticum
Tsuga canadensis
Tulipa spp.
Veronica spp.
Viburnum japonicum
V. odoratissimum

Lily-of-the-Valley Bush
 Calabrian Pine
 Canary Island Pine
 Slash Pine
 Aleppo Pine
 Austrian Pine
 Longleaf Pine
 Monterey Pine
 Eastern White Pine
 Scotch Pine
 Loblolly Pine
 Japanese Black Pine
 Virginia Pine
 Pistachio*
 Queensland Pittosporum
 Japanese Pittosporum
 Japanese Yew
 English Laurel
 Almond, Apricot, Nectarine, Peach, Plum and Prune*
 Douglas Fir**
 Firethorn (Scarlet)
 Firethorn
 Firethorn
 Bradford Pear spp.
 Oak, Shumard's Red
 Oak species
 Indian Hawthorne
 Yedda Hawthorne
 'Coral Bells,' 'Formosa,' 'Hino-crimson,' 'PJM'
 'Delaware Valley White', 'Roseum Elegans,'
 'Flame Creeper', 'Girard Crimson', 'Wakeiebisu'
 'White Gumbo', 'George L. Tabor'
 Lady Bank's Rose
 Rosemary*
 Black-Eyed Susan; Goldsturm
 Santolina
 Saxifrage; Purple Dome
 Pincushion Flower
 Stonecrop
 Stonecrop; Lidakense
 Stonecrop
 Stonecrop; Dragon's Blood
 Spirea; Anthony Waterer
 Australian Brushcherry
 Japanese Yew
 Yew
 Germander
 Meadow Rue
 American Arborvitae
 Star Jasmine
 Canada Hemlock
 Tulip species
 Veronica, Speedwell; Sunny Border
 Japanese Viburnum
 Sweet Viburnum

V. plicatum
 V. rigidum
 V. suspensum
 V. tinus
 V. trilobium
 V. wrightii
 Vinca major
 Vinca Minor
 Vitis spp.
 Weigela florida
 Yucca aloifolia
 Y. filametosa

Japanese Snowball
 Canary Island Viburnum
 Arrowwood Viburnum
 Laurustinus
 Cranberry Bush
 Leatherleaf Viburnum
 Greater Periwinkle, Vinca
 Common Periwinkle, Vinca
 Grape*
 Old Fashioned Weigela
 Spanish Bayonet, Yucca
 Adam's Needle, Yucca

*Do not use on food producing trees, vines or plants.

**Not for use on container grown plants.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container in a dry place. Store in locked area out of reach of children and domestic animals.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spilled bait, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance.

Container Disposal: Non-refillable container. Do not reuse or refill this container. Empty container through use according to label instructions. Do not rinse unless required for recycling. Offer for recycling, if available. If recycling is not available; then dispose of container in a sanitary landfill or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONDITIONS OF SALE AND WARRANTY

The DIRECTIONS FOR USE of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop, injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of Control Solutions, Inc. or the Seller. All such risks shall be assumed by the Buyer.

Control Solutions, Inc. warrants that this product conforms to the chemical description on the label, and is reasonably fit for the purposes set forth in the DIRECTIONS FOR USE, when it is used in accordance with such directions, subject to the inherent risks mentioned above.

CONTROL SOLUTIONS, INC. NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESSED OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. THIS WARRANTY EXTENDS TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNING, OR CAUTIONS. BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN

CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. CONTROL SOLUTIONS, INC. and the seller offer this product, and the Buyer and User accept it, subject to the foregoing CONDITIONS OF SALE AND WARRANTY.

Optional State Specific Language

[Help do your part in protecting our rivers and lakes by following label directives. Do not apply near water storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back onto your lawn. Do not apply to frozen ground.]

{Label may also include state specific language regarding Nitrogen application}