



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

June 28, 2024

Haley Wade
Lead Regulatory Manager
Control Solutions, Inc.
5903 Genoa-Red Bluff
Pasadena, TX 77507-1041

Subject: Label Amendment – Separated label into consumer and commercial sublabels, miscellaneous label updates, updated First Aid Box, MOA and HRM text added, added marketing claims

Product Name: PRODIAMINE 0.20 PLUS

EPA Registration Number: 53883-173

Application Date: July 1, 2020

Case Number: 483914

Dear Haley Wade:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website

is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Curtis Hildebrandt by phone at (202) 566-2770, or via email at hildebrandt.curtis@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read 'Kable Bo Davis', enclosed within a hand-drawn oval.

Kable Bo Davis
Senior Regulatory Specialist
Office of Pesticide Programs
Registration Division, Immediate Office

Enclosure

[Text in brackets and italics are notes to the label editor.]
 [Text in brackets alone are optional marketing statements or alternative statements.]

PRODIAMINE	GROUP	3	HERBICIDE
-------------------	--------------	----------	------------------

Sub label A: Commercial label

PRODIAMINE 0.20 PLUS

[FOR PRE-EMERGENCE CONTROL OF GRASS AND BROADLEAF WEEDS IN LAWNS AND
 ORNAMENTAL TURF

(excluding golf course putting greens)]

[LANDSCAPE ORNAMENTALS, ESTABLISHED PERENNIALS AND WILDFLOWER
 PLANTINGS]

ACTIVE INGREDIENT:

Prodiamine (CAS # 29091-21-2) 0.20%

OTHER INGREDIENTS: 99.80%

TOTAL: 100.00%



ADAMA


Consumer &
Professional
Solutions

Manufactured for:
 Control Solutions, Inc.
 5903 Genoa Red Bluff
 Pasadena, TX 77507

EPA Reg. No: 53883-173

EPA Est. No: _____

Net Contents: _____



KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for further precautions and First Aid information

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse. Wear long-sleeved shirt, long pants, and waterproof gloves.

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> ● Hold eye open and rinse slowly and gently with water for 15-20 minutes. ● Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. ● Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> ● Take off contaminated clothing. ● Rinse skin immediately with plenty of water for 15-20 minutes. ● Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center (800-222-1222) or doctor, or going for treatment. You may also contact SafetyCall® International (866) 897-8050 for emergency medical treatment information.	

[Text in brackets and italics are notes to the label editor.]

[Text in brackets alone are optional marketing statements or alternative statements.]

PRODIAMINE	GROUP	3	HERBICIDE
-------------------	--------------	----------	------------------

NON-TARGET ORGANISM ADVISORY STATEMENT: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize drift.

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

DIRECTION FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

[Not for use on plants being grown for sale or other commercial use, for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.] *[When label omits all applications other than turf, this section will be omitted.]*

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

WEED RESISTANCE MANAGEMENT

For resistance management, Prodiamine 0.20 Plus is a Group 3 herbicide. Any weed population may contain or develop plants naturally resistant to Prodiamine 0.20 Plus and other Group 3 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of Prodiamine 0.20 Plus or other Group 3 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide used and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g. higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing

[Text in brackets and italics are notes to the label editor.]

[Text in brackets alone are optional marketing statements or alternative statements.]

PRODIAMINE	GROUP	3	HERBICIDE
-------------------	--------------	----------	------------------

- or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
 - Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
 - Report any incidence of non-performance of this product against a particular weed species to your local retailer or your local extension specialist. If resistance is suspected, treat weed escapes with a herbicide having a different mechanism of action and/or use non-chemical means to remove escapes, as practical, with the goal of preventing further seed production.

BEST MANAGEMENT PRACTICES

Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is recommended. A diversified weed management program may include the use of multiple herbicides with different modes of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices. Research has demonstrated that using the labeled rate and directions for use is important to delay the selection for resistant weeds. Scouting after herbicide application is important because it can facilitate the early identification of weed shifts and/or weed resistance and thus provide direction on future weed management practices. One of the best ways to contain resistant weed populations is to implement measures to avoid allowing weeds to reproduce by seed or to proliferate vegetatively. Cleaning equipment between sites and avoiding movement of plant material between sites will greatly aid in reducing the spread of resistant weed seed.

This product is a pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in established lawns and ornamental turf (excluding golf course putting greens), landscape ornamentals and established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seeds, germination and root development. When used according to directions, it will control; barnyardgrass; bluegrass; annual (*Poa annua*); carpetweed; chickweed, common; chickweed, mouseear (from seed); crabgrass (large, smooth); crowfootgrass; cupgrass, woolly; foxtails, annual; goosegrass¹; henbit; itchgrass; johnsongrass (from seed); junglerice; knotweed; kochia; lambsquarter, common; lovegrass; panicum (Texas, fall, browntop); pigweed; purslane, common; pusley, Florida; rescuegrass³, shepherd's purse², signalgrass, broadleaf; speedwell, Persian; sprangletop; spurge, prostrate; witchgrass; woodsorrel, yellow (from seed).

¹In many areas [where not prohibited by fertilizer use restrictions**] a single application of 7.6 to 17.2 pounds/1,000 square feet (330 to 750 pounds/acre) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 7.6 to 11.5 pounds/1,000 square feet (330 to 500 pounds/acre), followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see "Minimum and Maximum Application Rate by Turfgrass Species" table in the Rates of Application section of this label).

²For this weed, apply in late summer, fall or winter prior to germination.

³Suppression only.

[**Consult your State Cooperative Extension Service for current information on annual application limitations for lawn fertilizers in your region.]

[Text in brackets and italics are notes to the label editor.]
 [Text in brackets alone are optional marketing statements or alternative statements.]

PRODIAMINE	GROUP	3	HERBICIDE
-------------------	--------------	----------	------------------

WHEN TO APPLY TO ESTABLISHED TURF: This product must be applied prior to germination of weed seeds, as it will not control established weeds.

THIS MAP SHOWS APPROXIMATE CRABGRASS GERMINATION DATES



APPLICATION DIRECTIONS: Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days application.

USE RESTRICTIONS:

- Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer.
- Do not apply to newly set sod until the following year.
- Do not apply to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.
- Do not apply through any type of irrigation system.
- Do not apply aurally.
- Do not use on food producing trees, vines or plants.

USE PRECAUTIONS:

The following precautions apply to the use of this product in turf grasses and lawns. Application of this product may thin emerged annual bluegrass and newly overseeded grasses.

- Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil).
- Application to turf stressed by drought, low fertility, or pest damage may result in turf injury.
- Disturbing the herbicide barrier with cultural practices (such as core aeration) may result in reduced weed control.

RATES OF APPLICATION: This product is labeled for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not exceed the maximum yearly rate listed:

MINIMUM AND MAXIMUM APPLICATION RATES (by Turf Grass Species)		
Turf Species	Minimum Rate	Maximum Annual Rate¹
Bahiagrass ^[3] , Bermudagrass ² , Centipedegrass, Zoysiagrass, Seashore Paspalum,	4 lbs/1,000 sq. ft. (0.008 lbs/ai) (174 lbs/acre)	17.2 lbs/1,000 sq. ft. per year (0.0344 lbs/ai) (750 lbs/acre per year)

[Text in brackets and italics are notes to the label editor.]

[Text in brackets alone are optional marketing statements or alternative statements.]

PRODIAMINE	GROUP	3	HERBICIDE
-------------------	--------------	----------	------------------

St. Augustinegrass Tall Fescue (including turf-Type)		
Buffalograss Kentucky Bluegrass Perennial Ryegrass	4 lbs/1,000 sq. ft. (0.008 lbs/ai) (174 lbs/acre)	11.5 lbs/1,000 sq. ft. per year (0.023 lbs/ai) (500 lbs/acre per year)
Creeping Red Fescue	4 lbs/1,000 sq. ft. (0.008 lbs/ai) (174 lbs/acre)	8.5 lbs/1,000 sq. ft. per year (0.017 lbs/ai) (370 lbs/acre per year)
Creeping Bentgrass	4 lbs/1,000 sq. ft. (0.008 lbs/ai) (174 lbs/acre)	7.6 lbs/1,000 sq. ft. per year (0.0152 lbs/ai) (330 lbs/acre per year)

¹These are the maximum rates per calendar year.

²May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 6 pounds/1,000 square feet (260 pounds/acre). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.

³[In Florida: Do not exceed 2 applications per year to Bahiagrass and Centipedegrass in the north. Do not exceed 3 applications per year to Centipedegrass in the central and south regions. Consult your State Cooperative Extension Service for current information on annual application limitations for lawn fertilizers in your region.]

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates labeled for each turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control.

WHEN TO OVERSEED AFTER APPLICATION: This product will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/reseeding results. (See map above.)

USE RATE	MONTHS BEFORE OVERSEEDING		
	North	Transition	South
4 pounds/1,000 sq. ft. (174 pounds/acre)	4	4	4
7.6 pounds/1,000 sq. ft. (330 pounds/acre)	5	4	4
8.5 pounds/1,000 sq. ft. (370 pounds/acre)	6	5	5
9.2 pounds/1,000 sq. ft. (400 pounds/acre)	-	6	6
11.5 pounds/1,000 sq. ft. (500 pounds/acre)	-	7	7
13 pounds/1,000 sq. ft. (570 pounds/acre)	-	-	9
14.9 pounds/1,000 sq. ft. (650 pounds/acre)	-	-	10
17.2 pounds/1,000 sq. ft. (750 pounds/acre)	-	-	12

LANDSCAPE ORNAMENTALS: (including established perennials and wildflower plantings): This product may be applied for residual pre-emergence weed control in ornamentals. Apply at the rate of 7.6 to 17.2 pounds/1,000 square feet (330 to 750 pounds/acre) (0.0152 – 0.0344 lbs/ai) in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 17.2 pounds/1,000 square feet (750 pounds/acre) (0.0344 lbs/ai) per year.

[Spreader Settings Chart is Optional]

SPREADER SETTINGS to apply at the listed pounds per 1,000 square feet
--

[Text in brackets and italics are notes to the label editor.]

[Text in brackets alone are optional marketing statements or alternative statements.]

PRODIAMINE	GROUP	3	HERBICIDE
-------------------	--------------	----------	------------------

	4 lbs.	6 lbs.	7.6 lbs.	8.5 lbs.	11.5 lbs.	17.2 lbs.
Spreader A	X	X	X	X	X	X

[Note to reviewer: May contain additional spreader brands/models, spreader settings and spreader use instructions as appropriate for the product]

APPLICATION TIMING AND INFORMATION: This product may be applied to newly transplanted and established ornamentals as a broadcast, over the top, or directed application. Around new transplants, delay application to allow soil to settle and water thoroughly before applying. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union. This product is a pre-emergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when applied to soil free of clods, weeds and debris such as leaves. Prior to application, control existing vegetation by hand weeding, cultivation or the use of an appropriate post-emergence herbicide.

TOLERANT ORNAMENTAL SPECIES: This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

SCIENTIFIC NAME	COMMON NAME
<i>Abies spp.</i>	Fir species** (Balsam, Frasier, Noble, etc.)
<i>Abelia grandiflora</i>	Abelia: Sherwood
<i>Acer palmatum</i>	Japanese Maple
<i>A. Platanoides</i>	Norway Maple
<i>Achillea spp.</i>	Yarrow: King Edward
<i>Actindia chinensis</i>	Kiwi*
<i>Agapanthus africanus</i>	Lily-of-the-Nile (African Lily)
<i>Agapanthus orientalis</i>	
<i>Akebia quintata</i>	Five-leaf or Chocolate Vine
<i>Allium cemuum</i>	Lady's Leek, Nodding Onion
<i>Anemone hybrida</i>	Japanese Anemone
<i>Aquilegia spp.</i>	Aquilegia: Red and Gold
<i>Arctostaphylos densiflora</i>	Vine Hill Manzanita
<i>Arctotheca calendula</i>	Cape Weed
<i>Artemisia spp.</i>	Wormwood; Silver Mound, Castle
<i>Aster spp.</i>	Aster: Bonnie Blue, Purple Dome
<i>Aster X Frikartii</i>	
<i>Athrium Filiz-femina</i>	Lady Fern; Fern Lady
<i>Aucuba japonica</i>	Japanese Aucuba
<i>Begonia spp.</i>	Fibrous Begonia: Hardy Grandis
<i>Berberis gladwynensis</i>	Barberry
<i>B. Julianae</i>	Wintergreen Barberry
<i>B. mentorensis</i>	Mentor Barberry
<i>B. Thunbergii</i>	Japanese Barberry
<i>B. Verruculosa</i>	Warty Barberry
<i>Bergenia cordifolia</i>	
<i>Boltonia asteroides</i>	Snowbank
<i>Buddleia davidii</i>	Butterfly-Bush (Dwarf Blue); Royal Red

[Text in brackets and italics are notes to the label editor.]

[Text in brackets alone are optional marketing statements or alternative statements.]

PRODIAMINE	GROUP	3	HERBICIDE
------------	-------	---	-----------

<i>Buxus microphylla</i>	Japanese Boxwood
<i>Callistemon viminalis</i>	Weeping Bottlebrush
<i>Calluna vulgaris</i>	Scotch Heather
<i>Campanula carpatica</i>	Tussock Bellflower, (White Clips)
<i>Campis X Tagliabuana</i>	Trumpet Flower, Madame Galen
<i>Carpobrotus edulis</i>	Hottentot Fig (Ice Plant)
<i>Cassia artemisoides</i>	Feathery Cassia
<i>Ceanothus rigidus</i>	Wild Lilac
<i>Ceratostigma plumbagonoides</i>	
<i>Chamaecyparis pisifera</i>	False Cypress
<i>Cleyera japonica</i>	Cleyera
<i>Citrus spp.</i>	Citrus species*
<i>Coreopsis spp.</i>	Coreopsis (Calliopsis) Early Sunrise, Moonbeam
<i>Cornus florida</i>	Flowering Dogwood
<i>C. Stolonifera</i>	American Dogwood
<i>Cortaderia selloana</i>	Pampas Grass
<i>Cotoneaster apiculatus</i>	Cranberry Cotoneaster
<i>C. buxifolius</i>	Cotoneaster
<i>C. dammeri</i>	Bearberry Cotoneaster
<i>C. Microphyllus</i>	Rockspray Cotoneaster
<i>Cretaeus spp.</i>	Hawthorne
<i>Crocosmia spp.</i>	Lucifer
<i>Cupressus sempervirens</i>	Italian Cypress
<i>Delosperma alba</i>	White Trailing Ice Plant
<i>Delosperma spp.</i>	Cooperi Pink
<i>Delphinium spp.</i>	Larkspur: Blue Elf
<i>Dianthus deltoides</i>	Dianthus: Maiden Pinks "Zing"
<i>D. gratianopolitanus</i>	Cheddar Pink
<i>Dodonea viscosa</i>	Hop Bush
<i>Echinacea purpurea</i>	Coneflower, Purple; Magnus
<i>Elaeagnus pungens</i>	Silverberry
<i>Euonymus fortunei</i>	Wintercreeper
<i>E. japonica</i>	Evergreen Euonymus
<i>E. kiautschovica</i>	Spreading Euonymus
<i>Fatsia japonica</i>	Japanese Aralia
<i>Forsythia intermedia</i>	Border Forsythia
<i>Forsythia suspensa</i>	Weeping Forsythia
<i>F. viridissima</i>	Greenstem Forsythia
<i>Gaillardia spp.</i>	Gaillardia, Blanket Flower: "Goblin"
<i>Gardenia jasminoides</i>	Gardenia, Cape-Jasmine
<i>Gaura spp.</i>	
<i>Gentiana dahurica</i>	Gentian
<i>Geranium cinereum</i>	Cranesbill
<i>Gladiolus spp.</i>	Gladiolus species**
<i>Gypsophilia repens</i>	Baby's Breath
<i>Hedera helix</i>	English Ivy
<i>Hellanthemum spp.</i>	Sunrose
<i>Hemerocallis spp.</i>	Daylily: Aztec Gold, Stella de Oro, Tender Love
<i>Hibiscus</i>	Rose of Sharon**
<i>Hibiscus spp.</i>	Mallow: Disco Belle White

[Text in brackets and italics are notes to the label editor.]

[Text in brackets alone are optional marketing statements or alternative statements.]

PRODIAMINE	GROUP	3	HERBICIDE
------------	-------	---	-----------

<i>Hibiscus rosa-sinensis</i>	Chinese Hibiscus
<i>Hosta plantaginea</i>	Hosta, Plantain Lily (Fragrant)
<i>H. Sieboldiana</i>	Hosta, "Seersucker"
<i>Houttuynia cordata</i> var. <i>variegata</i>	
<i>Hydrangea macrophylla</i>	Bigleaf Hydrangea
<i>Ilex Bernyi</i>	Holly
<i>Ilex cornuta</i>	Chinese Holly
<i>I. crenata</i>	Japanese Holly
<i>I. opaca</i>	American Holly
<i>I. vomitoria</i>	Yaupon Holly, Schillings
<i>Inula ensifolia</i>	
<i>Iris ensata</i>	Sword-Leaved Iris; Jodlesong
<i>Iris siberica</i>	Siberian Iris; Cabernet
<i>Iris</i> spp.	Iris species**
<i>Jasminum nudiflorum</i>	Winter Jasmine
<i>Juniperus chinensis</i>	Chinese Juniper; Nick's Compact, Parsonii
<i>J. conferta</i>	Shore Juniper; Blue Pacific
<i>J. davurica</i>	Parsoni
<i>J. horizontalis</i>	Creeping Juniper
<i>Juglans</i> spp.	Walnut*
<i>Justicia brandegeana</i>	Shrimp Plant
<i>Lagerstromia indica</i>	Crape Myrtle
<i>Lagerstromia indica</i> X <i>Fauriei</i>	Crape Myrtle; Tuscarora
<i>Lantana montevidensis</i>	Weeping Lantana
<i>Lavender</i> spp.	Lavender; Munstead
<i>Leonopodium alpinum</i>	Edelweiss
<i>Ligustrum amurense</i>	Amur Privet
<i>Ligustrum sinense</i>	Chinese Privet; Variegata
<i>L. japonicum</i>	Japanese Privet
<i>L. lucidum</i>	Glossy Privet (Waxleaf)
<i>Lilium</i> spp.	Lily; Jazz
<i>Liriope muscari</i>	Big Blue Lily, Silver Mound, Evergreen Giant
<i>Liriope muscari</i> var. <i>variegata</i>	Liriope, Variegated
<i>Liriope spicata</i>	Liriope, Creeping
<i>Lobelia Cardinalis</i>	Cardinal Flower, Indian Pink
<i>Lonicera japonica</i>	Japanese Honeysuckle
<i>L. tatarica</i>	Tatarian Honeysuckle
<i>Lorapetalum chinense</i>	Burgundy
<i>Lythrum</i> spp.	Loosestrife; Modern Pink
<i>Magnolia</i> spp.	Magnolia species**
<i>Maleophora luteola</i>	Ice Plant
<i>Malus</i> spp.	Crabapple*
<i>Miscanthus sinensis</i>	Yaku Jima, Silberfeder
<i>Nandina domestica</i>	Heavenly Bamboo
<i>Narcissus</i> spp.	Narcissus species**
<i>Nerium oleander</i>	Oleander
<i>Olea europaea</i>	Olive*
<i>Ophiopogon japonicus</i>	Mondo Grass
<i>Omanthus heterophyllus</i>	Osmanthus (False Holly); Gulf Tide
<i>Osteospermum fruticosum</i>	Trailing African Daisy
<i>Oxydendrum arboreum</i>	Sourwood
<i>Paeonia suffruticosa</i>	Tree Peony

[Text in brackets and italics are notes to the label editor.]

[Text in brackets alone are optional marketing statements or alternative statements.]

PRODIAMINE	GROUP	3	HERBICIDE
------------	-------	---	-----------

<i>Pennisetum setaceum</i>	Fountain Grass (Dwarf)
<i>Perovskia atriplicifolia</i>	
<i>Persea americana</i>	Avocado*
<i>Photinia Fraseri</i>	Fraser's Photinia (Redtip)
<i>Physotegia Virginiana</i>	Dragonhead, False; Vivid
<i>Picea spp.</i>	Spruce species** (Colorado Blue, Norway, etc.)
<i>Pieris japonica</i>	Lily-of-the-Valley Bush
<i>Pinus brutia</i>	Calabrian Pine
<i>P. canariensis</i>	Canary Island Pine
<i>P. Elliottii</i>	Slash Pine
<i>P. halepensis</i>	Aleppo Pine
<i>P. nigra</i>	Austrian Pine
<i>P. palustris</i>	Longleaf Pine
<i>P. radiata</i>	Monterey Pine
<i>P. strobus</i>	Eastern White Pine
<i>P. sylvestris</i>	Scotch Pine
<i>P. taeda</i>	Loblolly Pine
<i>P. Thunbergiana</i>	Japanese Black Pine
<i>P. Virginiana</i>	Virginia Pine
<i>Pistachio spp.</i>	Pistachio*
<i>Pittosporum rhombifolium</i>	Queensland Pittosporum
<i>P. tobira</i>	Japanese Pittosporum
<i>Podocarpus macrophyllus</i>	Japanese Yew
<i>Prunus laurocerasus</i>	English Laurel
<i>Prunus spp.</i>	Almond, Apricot, Nectarine, Peach, Plum and Prune*
<i>Pseudotsuga menziesii</i>	Douglas Fir**
<i>Pyracantha coccinea</i>	Firethorn (Scarlet)
<i>P. fortuneana</i>	Firethorn
<i>P. koidzumii</i>	Firethorn
<i>Pyrus spp.</i>	Bradford Pear spp.
<i>Quercus Shumardii</i>	Oak, Shumard's Red
<i>Quercus spp.</i>	Oak species
<i>Raphiolepis indica</i>	Indian Hawthorne
<i>Raphiolepis umbellata</i>	Yedda Hawthorne
<i>Rhododendron</i> (including Azalea)	'Coral Bells,' 'Formosa,' 'Hino-crimson,' 'PJM' 'Delaware Valley White,' 'Roseum Elegans,' 'Flame Creeper,' 'Girard Crimson', 'Wakeiebisu 'White Gumbo', 'George L. Tabor'
<i>Rosa banksiae</i>	Lady Bank's Rose
<i>Rosmarinus officinalis</i>	Rosemary*
<i>Rudbeckia spp.</i>	Black-Eyed Susan; Goldsturm
<i>Santolina virens</i>	Santolina
<i>Saxifraga spp.</i>	Saxifrage; Purple Dome
<i>Scabiosa spp.</i>	Pincushion Flower
<i>Sedum album</i>	Stonecrop
<i>Sedum caudicola</i>	Stonecrop; Lidakense
<i>Sedum dasyphyllum</i>	Stonecrop
<i>Sedum spurium</i>	Stonecrop; Dragon's Blood
<i>Spiraea bumalda</i>	Spiraea; Anthony Waterer
<i>Syzygium paniculatum</i>	Australian Brushcherry

[Text in brackets and italics are notes to the label editor.]

[Text in brackets alone are optional marketing statements or alternative statements.]

PRODIAMINE	GROUP	3	HERBICIDE
-------------------	--------------	----------	------------------

<i>Taxus cuspidata</i>	Japanese Yew
<i>T. media</i>	Yew
<i>Teucrium spp.</i>	Germander
<i>Thalictrum diptercarpum</i>	Meadow Rue
<i>Thuja occidentalis</i>	American Arborvitae
<i>Trachelospermum asiaticum</i>	Star Jasmine
<i>Tsuga canadensis</i>	Canada Hemlock
<i>Tulipa spp.</i>	Tulip species
<i>Veronica spp.</i>	Veronica, Speedwell; Sunny Border
<i>Viburnum japonicum</i>	Japanese Viburnum
<i>V. odoratissimum</i>	Sweet Viburnum
<i>V. plicatum</i>	Japanese Snowball
<i>V. rigidum</i>	Canary Island Viburnum
<i>V. suspensum</i>	Arrowwood Viburnum
<i>V. tinus</i>	Laurustinus
<i>V. trilobium</i>	Cranberry Bush
<i>V. wrightii</i>	Leatherleaf Viburnum
<i>Vinca major</i>	Greater Periwinkle, Vinca
<i>Vinca Minor</i>	Common Periwinkle, Vinca
<i>Vitis spp.</i>	Grape*
<i>Weigela florida</i>	Old Fashioned Weigela
<i>Yucca aloifolia</i>	Spanish Bayonet, Yucca
<i>Y. filametosa</i>	Adam's Needle, Yucca

*Do not use on food producing trees, vines or plants.

**Not for use on container grown plants.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a dry place. Store in locked area out of reach of children and domestic animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Non-refillable container. Do not reuse or refill this container. Empty container through use according to label instructions. Offer for recycling, if available, or dispose of empty bag in a sanitary landfill or by other procedures allowed by state and local authorities.

CONDITIONS OF SALE AND WARRANTY

The DIRECTIONS FOR USE of this product are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Injury/ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of Control Solutions, Inc. or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

Control Solutions, Inc. warrants that this product conforms to the chemical description on the label, and is reasonably fit for the purposes set forth in the DIRECTIONS FOR USE, when it is used in accordance with such directions, subject to the inherent risks mentioned above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, CONTROL SOLUTIONS, INC. NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESSED OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF

[Text in brackets and italics are notes to the label editor.]

[Text in brackets alone are optional marketing statements or alternative statements.]

PRODIAMINE	GROUP	3	HERBICIDE
-------------------	--------------	----------	------------------

MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. THIS WARRANTY EXTENDS TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNING, OR CAUTIONS. BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. CONTROL SOLUTIONS, INC. and the seller offer this product, and the Buyer and User accept it, subject to the foregoing CONDITIONS OF SALE AND WARRANTY.

Optional Marketing Language:

[PREVENT!]

[PREVENT! 15,000], [PREVENT! 10,000], [PREVENT! 5,000]

[Weed Grass Preventer and Turf Food]

[With Iron and Sulfur Plus Slow Release Nitrogen]

[Covers [5,000], [10,000], [15,000] Square Feet]

[For Pre-Emergence Control of: [Crabgrass], [Goosegrass], [Foxtail], [Barnyardgrass] [& other listed Annual Grass and Broadleaf Weeds] in Established Turfgrass]

[STOPS MANY WEED GRASSES BEFORE THEY GROW]

[PREVENTS CRABGRASS, ANNUAL BLUEGRASS, FOXTAIL]

[Loveland Turf Network]

[Loveland Serving the Environment of Man]

[GREEN GRASS [Cleans our Air, Cools our Earth, Controls Erosion, Fights Pollution, Purifies our Water]]

[Call [the Loveland Turf Network] For Fertilizers - Seed - Sod - Rock – Trees]

[We're the if not, Why not Golf Course Family of the Midwest - Pioneers in grass for over three generations. Suppliers to cities, golf course, parks, home and industry. The 1st purchase sells the 2nd!]

[Visit www.grasspad.com] Omaha, Nebraska: 402/331-5262 Kansas/Missouri: 800-922-4100]

Optional State Specific Language:

[Help do your part in protecting our rivers and lakes by following label directives. Do not apply near water storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back onto your lawn. Do not apply to frozen ground.]

{Label may also include state specific language regarding Nitrogen application}

[Text in brackets and italics are notes to the label editor.]
 [Text in brackets alone are optional marketing statements or alternative statements.]

PRODIAMINE	GROUP	3	HERBICIDE
-------------------	--------------	----------	------------------

Sub label B: Homeowner label

PRODIAMINE 0.20 PLUS

[FOR PRE-EMERGENCE CONTROL OF GRASS AND BROADLEAF WEEDS IN LAWNS AND ORNAMENTAL TURF

(excluding golf course putting greens)]

[LANDSCAPE ORNAMENTALS, ESTABLISHED PERENNIALS AND WILDFLOWER PLANTINGS]

ACTIVE INGREDIENT:

Prodiamine (CAS # 29091-21-2) 0.20%

OTHER INGREDIENTS: 99.80%

TOTAL: 100.00%



ADAMA

Consumer &
Professional
Solutions

Manufactured for:
Control Solutions, Inc.
5903 Genoa Red Bluff
Pasadena, TX 77507

EPA Reg. No: 53883-173

EPA Est. No: _____

Net Contents: _____

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See back panel for further precautions and First Aid information

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
<p>Have the product container or label with you when calling a poison control center (800-222-1222) or doctor, or going for treatment. You may also contact SafetyCall® International (866) 897-8050 for emergency medical treatment information.</p>	

NON-TARGET ORGANISM ADVISORY STATEMENT: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize drift.

[Text in brackets and italics are notes to the label editor.]

[Text in brackets alone are optional marketing statements or alternative statements.]

PRODIAMINE	GROUP	3	HERBICIDE
------------	-------	---	-----------

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

DIRECTION FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

[Not for use on plants being grown for sale or other commercial use, for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.] *[When label omits all applications other than turf, this section will be omitted.]*

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product is a pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in established lawns and ornamental turf (excluding golf course putting greens), landscape ornamentals and established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seeds, germination and root development. When used according to directions, it will control; barnyardgrass; bluegrass; annual (*Poa annua*); carpetweed; chickweed, common; chickweed, mouseear (from seed); crabgrass (large, smooth); crowfootgrass; cupgrass, woolly; foxtails, annual; goosegrass¹; henbit; itchgrass; johnsongrass (from seed); junglerice; knotweed; kochia; lambsquarter, common; lovegrass; panicum (Texas, fall, browntop); pigweed; purslane, common; pusley, Florida; rescuegrass³, shepherd's purse², signalgrass, broadleaf; speedwell, Persian; sprangletop; spurge, prostrate; witchgrass; woodsorrel, yellow (from seed).

¹In many areas [where not prohibited by fertilizer use restrictions**] a single application of 7.6 to 17.2 pounds/1,000 square feet (330 to 750 pounds/acre) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 7.6 to 11.5 pounds/1,000 square feet (330 to 500 pounds/acre), followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see "Minimum and Maximum Application Rate by Turfgrass Species" table in the Rates of Application section of this label).

²For this weed, apply in late summer, fall or winter prior to germination.

³Suppression only.

[**Consult your State Cooperative Extension Service for current information on annual application limitations for lawn fertilizers in your region.]

WHEN TO APPLY TO ESTABLISHED TURF: This product must be applied prior to germination of weed seeds, as it will not control established weeds.

THIS MAP SHOWS APPROXIMATE CRABGRASS GERMINATION DATES

[Text in brackets and italics are notes to the label editor.]
 [Text in brackets alone are optional marketing statements or alternative statements.]

PRODIAMINE	GROUP	3	HERBICIDE
-------------------	--------------	----------	------------------



APPLICATION DIRECTIONS: Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days application.

USE RESTRICTIONS:

- Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer.
- Do not apply to newly set sod until the following year.
- Do not apply to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.
- Do not apply through any type of irrigation system.
- Do not apply aurally.
- Do not use on food producing trees, vines or plants.

USE PRECAUTIONS:

The following precautions apply to the use of this product in turf grasses and lawns. Application of this product may thin emerged annual bluegrass and newly overseeded grasses.

- Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil).
- Application to turf stressed by drought, low fertility, or pest damage may result in turf injury.
- Disturbing the herbicide barrier with cultural practices (such as core aeration) may result in reduced weed control.

RATES OF APPLICATION: This product is labeled for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not exceed the maximum yearly rate listed:

MINIMUM AND MAXIMUM APPLICATION RATES (by Turf Grass Species)		
Turf Species	Minimum Rate	Maximum Annual Rate¹
Bahiagrass ^[3] , Bermudagrass ² Centipedegrass, Zoysiagrass, Seashore Paspalum, St. Augustinegrass Tall Fescue (including turf- Type)	4 lbs/1,000 sq. ft. (0.008 lbs/ai) (174 lbs/acre)	17.2 lbs/1,000 sq. ft. per year (0.0344 lbs/ai) (750 lbs/acre per year)
Buffalograss Kentucky Bluegrass Perennial Ryegrass	4 lbs/1,000 sq. ft. (0.008 lbs/ai) (174 lbs/acre)	11.5 lbs/1,000 sq. ft. per year (0.023 lbs/ai) (500 lbs/acre per year)

[Text in brackets and italics are notes to the label editor.]
 [Text in brackets alone are optional marketing statements or alternative statements.]

PRODIAMINE	GROUP	3	HERBICIDE
-------------------	--------------	----------	------------------

Creeping Red Fescue	4 lbs/1,000 sq. ft. (0.008 lbs/ai) (174 lbs/acre)	8.5 lbs/1,000 sq. ft. per year (0.017 lbs/ai) (370 lbs/acre per year)
Creeping Bentgrass	4 lbs/1,000 sq. ft. (0.008 lbs/ai) (174 lbs/acre)	7.6 lbs/1,000 sq. ft. per year (0.0152 lbs/ai) (330 lbs/acre per year)

¹These are the maximum rates per calendar year.

²May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 6 pounds/1,000 square feet (260 pounds/acre). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.

³[In Florida: Do not exceed 2 applications per year to Bahiagrass and Centipedegrass in the north. Do not exceed 3 applications per year to Centipedegrass in the central and south regions. Consult your State Cooperative Extension Service for current information on annual application limitations for lawn fertilizers in your region.]

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates labeled for each turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control.

WHEN TO OVERSEED AFTER APPLICATION: This product will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/reseeding results. (See map above.)

USE RATE	MONTHS BEFORE OVERSEEDING		
	North	Transition	South
4 pounds/1,000 sq. ft. (174 pounds/acre)	4	4	4
7.6 pounds/1,000 sq. ft. (330 pounds/acre)	5	4	4
8.5 pounds/1,000 sq. ft. (370 pounds/acre)	6	5	5
9.2 pounds/1,000 sq. ft. (400 pounds/acre)	-	6	6
11.5 pounds/1,000 sq. ft. (500 pounds/acre)	-	7	7
13 pounds/1,000 sq. ft. (570 pounds/acre)	-	-	9
14.9 pounds/1,000 sq. ft. (650 pounds/acre)	-	-	10
17.2 pounds/1,000 sq. ft. (750 pounds/acre)	-	-	12

LANDSCAPE ORNAMENTALS: (including established perennials and wildflower plantings): This product may be applied for residual pre-emergence weed control in ornamentals. Apply at the rate of 7.6 to 17.2 pounds/1,000 square feet (330 to 750 pounds/acre) (0.0152 – 0.0344 lbs/ai) in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 17.2 pounds/1,000 square feet (750 pounds/acre) (0.0344 lbs/ai) per year.

[Spreader Settings Chart is Optional]

SPREADER SETTINGS to apply at the listed pounds per 1,000 square feet						
	4 lbs.	6 lbs.	7.6 lbs.	8.5 lbs.	11.5 lbs.	17.2 lbs.
Spreader A	X	X	X	X	X	X
[Note to reviewer: May contain additional spreader brands/models, spreader settings and spreader use instructions as appropriate for the product]						

[Text in brackets and italics are notes to the label editor.]

[Text in brackets alone are optional marketing statements or alternative statements.]

PRODIAMINE	GROUP	3	HERBICIDE
-------------------	--------------	----------	------------------

APPLICATION TIMING AND INFORMATION: This product may be applied to newly transplanted and established ornamentals as a broadcast, over the top, or directed application. Around new transplants, delay application to allow soil to settle and water thoroughly before applying. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union. This product is a pre-emergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when applied to soil free of clods, weeds and debris such as leaves. Prior to application, control existing vegetation by hand weeding, cultivation or the use of an appropriate post-emergence herbicide.

TOLERANT ORNAMENTAL SPECIES: This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

SCIENTIFIC NAME	COMMON NAME
<i>Abies spp.</i>	Fir species** (Balsam, Frasier, Noble, etc.)
<i>Abelia grandiflora</i>	Abelia: Sherwood
<i>Acer palmatum</i>	Japanese Maple
<i>A. Platanoides</i>	Norway Maple
<i>Achillea spp.</i>	Yarrow: King Edward
<i>Actindia chinensis</i>	Kiwi*
<i>Agapanthus africanus</i>	Lily-of-the-Nile (African Lily)
<i>Agapanthus orientalis</i>	
<i>Akebia quintata</i>	Five-leaf or Chocolate Vine
<i>Allium cernuum</i>	Lady's Leek, Nodding Onion
<i>Anemone hybrida</i>	Japanese Anemone
<i>Aquilegia spp.</i>	Aquilegia: Red and Gold
<i>Arctostaphylos densiflora</i>	Vine Hill Manzanita
<i>Arctotheca calendula</i>	Cape Weed
<i>Artemisia spp.</i>	Wormwood; Silver Mound, Castle
<i>Aster spp.</i>	Aster: Bonnie Blue, Purple Dome
<i>Aster X Frikartii</i>	
<i>Athrium Filiz-femina</i>	Lady Fern; Fern Lady
<i>Aucuba japonica</i>	Japanese Aucuba
<i>Begonia spp.</i>	Fibrous Begonia: Hardy Grandis
<i>Berberis gladwynensis</i>	Barberry
<i>B. Julianae</i>	Wintergreen Barberry
<i>B. mentorensis</i>	Mentor Barberry
<i>B. Thunbergii</i>	Japanese Barberry
<i>B. Verruculosa</i>	Warty Barberry
<i>Bergenia cordifolia</i>	
<i>Boltonia asteroides</i>	Snowbank
<i>Buddleia davidii</i>	Butterfly-Bush (Dwarf Blue); Royal Red
<i>Buxus microphylla</i>	Japanese Boxwood
<i>Callistemon viminalis</i>	Weeping Bottlebrush
<i>Calluna vulgaris</i>	Scotch Heather
<i>Campanula carpatica</i>	Tussock Bellflower, (White Clips)

[Text in brackets and italics are notes to the label editor.]

[Text in brackets alone are optional marketing statements or alternative statements.]

PRODIAMINE	GROUP	3	HERBICIDE
------------	-------	---	-----------

<i>Campis X Tagliabuana</i>	Trumpet Flower, Madame Galen
<i>Carpobrotus edulis</i>	Hottentot Fig (Ice Plant)
<i>Cassia artemisoides</i>	Feathery Cassia
<i>Ceanothus rigidus</i>	Wild Lilac
<i>Ceratostigma plumbagonoides</i>	
<i>Chamaecyparis pisifera</i>	False Cypress
<i>Cleyera japonica</i>	Cleyera
<i>Citrus spp.</i>	Citrus species*
<i>Coreopsis spp.</i>	Coreopsis (Calliopsis) Early Sunrise, Moonbeam
<i>Cornus florida</i>	Flowering Dogwood
<i>C. Stolonifera</i>	American Dogwood
<i>Cortaderia selloana</i>	Pampas Grass
<i>Cotoneaster apiculatus</i>	Cranberry Cotoneaster
<i>C. buxifolius</i>	Cotoneaster
<i>C. dammeri</i>	Bearberry Cotoneaster
<i>C. Microphyllus</i>	Rockspray Cotoneaster
<i>Cretaegus spp.</i>	Hawthorne
<i>Crocosmia spp.</i>	Lucifer
<i>Cupressus sempervirens</i>	Italian Cypress
<i>Delosperma alba</i>	White Trailing Ice Plant
<i>Delosperma spp.</i>	Cooperi Pink
<i>Delphinium spp.</i>	Larkspur: Blue Elf
<i>Dianthus deltoides</i>	Dianthus: Maiden Pinks "Zing"
<i>D. gratianopolitanus</i>	Cheddar Pink
<i>Dodonea viscosa</i>	Hop Bush
<i>Echinacea purpurea</i>	Coneflower, Purple; Magnus
<i>Elaeagnus pungens</i>	Silverberry
<i>Euonymus fortunei</i>	Wintercreeper
<i>E. japonica</i>	Evergreen Euonymus
<i>E. kiautschovica</i>	Spreading Euonymus
<i>Fatsia japonica</i>	Japanese Aralia
<i>Forsythia intermedia</i>	Border Forsythia
<i>Forsythia suspensa</i>	Weeping Forsythia
<i>F. viridissima</i>	Greenstem Forsythia
<i>Gaillardia spp.</i>	Gaillardia, Blanket Flower: "Goblin"
<i>Gardenia jasminoides</i>	Gardenia, Cape-Jasmine
<i>Gaura spp.</i>	
<i>Gentiana dahurica</i>	Gentian
<i>Geranium cinereum</i>	Cranesbill
<i>Gladiolus spp.</i>	Gladiolus species**
<i>Gypsophilia repens</i>	Baby's Breath
<i>Hedera helix</i>	English Ivy
<i>Hellanthemum spp.</i>	Sunrose
<i>Hemerocallis spp.</i>	Daylily: Aztec Gold, Stella de Oro, Tender Love
<i>Hibiscus</i>	Rose of Sharon**
<i>Hibiscus spp.</i>	Mallow: Disco Belle White
<i>Hibiscus rosa-sinensis</i>	Chinese Hibiscus
<i>Hosta plantaginea</i>	Hosta, Plantain Lily (Fragrant)
<i>H. Sieboldiana</i>	Hosta, "Seersucker"
<i>Houttuynia cordata var. variegata</i>	

[Text in brackets and italics are notes to the label editor.]

[Text in brackets alone are optional marketing statements or alternative statements.]

PRODIAMINE	GROUP	3	HERBICIDE
------------	-------	---	-----------

<i>Hydrangea macrophylla</i>	Bigleaf Hydrangea
<i>Ilex Bernyi</i>	Holly
<i>Ilex cornuta</i>	Chinese Holly
<i>I. crenata</i>	Japanese Holly
<i>I. opaca</i>	American Holly
<i>I. vomitoria</i>	Yaupon Holly, Schillings
<i>Inula ensifolia</i>	
<i>Iris ensata</i>	Sword-Leaved Iris; Jodlesong
<i>Iris siberica</i>	Siberian Iris; Cabernet
<i>Iris spp.</i>	Iris species**
<i>Jasminum nudiflorum</i>	Winter Jasmine
<i>Juniperus chinensis</i>	Chinese Juniper; Nick's Compact, Parsonnii
<i>J. conferta</i>	Shore Juniper; Blue Pacific
<i>J. davurica</i>	Parsoni
<i>J. horizontalis</i>	Creeping Juniper
<i>Juglans spp.</i>	Walnut*
<i>Justicia brandegeana</i>	Shrimp Plant
<i>Lagerstromia indica</i>	Crape Myrtle
<i>Lagerstromia indica X Fauriei</i>	Crape Myrtle; Tuscarora
<i>Lantana montevidensis</i>	Weeping Lantana
<i>Lavender spp.</i>	Lavender; Munstead
<i>Leonopodium alpinum</i>	Edelweiss
<i>Ligustrum amurense</i>	Amur Privet
<i>Ligustrum sinense</i>	Chinese Privet; Variegata
<i>L. japonicum</i>	Japanese Privet
<i>L. lucidum</i>	Glossy Privet (Waxleaf)
<i>Lilium spp.</i>	Lily; Jazz
<i>Liriope muscari</i>	Big Blue Lily, Silver Mound, Evergreen Giant
<i>Liriope muscari var. variegata</i>	Liriope, Variegated
<i>Liriope spicata</i>	Liriope, Creeping
<i>Lobelia Cardinalis</i>	Cardinal Flower, Indian Pink
<i>Lonicera japonica</i>	Japanese Honeysuckle
<i>L. tatarica</i>	Tatarian Honeysuckle
<i>Lorapetalum chinense</i>	Burgundy
<i>Lythrum spp.</i>	Loosestrife; Modern Pink
<i>Magnolia spp.</i>	Magnolia species**
<i>Maleophora luteola</i>	Ice Plant
<i>Malus spp.</i>	Crabapple*
<i>Miscanthus sinensis</i>	Yaku Jima, Silberfeder
<i>Nandina domestica</i>	Heavenly Bamboo
<i>Narcissus spp.</i>	Narcissus species**
<i>Nerium oleander</i>	Oleander
<i>Olea europaea</i>	Olive*
<i>Ophiopogon japonicus</i>	Mondo Grass
<i>Omanthus heterophyllus</i>	Osmanthus (False Holly); Gulf Tide
<i>Osteospermum fruticosum</i>	Trailing African Daisy
<i>Oxydendrum arboreum</i>	Sourwood
<i>Paeonia suffruticosa</i>	Tree Peony
<i>Pennisetum setaceum</i>	Fountain Grass (Dwarf)
<i>Perovskia atriplicifolia</i>	
<i>Persea americana</i>	Avocado*
<i>Photinia Fraseri</i>	Fraser's Photinia (Redtip)

[Text in brackets and italics are notes to the label editor.]

[Text in brackets alone are optional marketing statements or alternative statements.]

PRODIAMINE	GROUP	3	HERBICIDE
------------	-------	---	-----------

<i>Physotegia Virginiana</i>	Dragonhead, False; Vivid
<i>Picea spp.</i>	Spruce species** (Colorado Blue, Norway, etc.)
<i>Pieris japonica</i>	Lily-of-the-Valley Bush
<i>Pinus brutia</i>	Calabrian Pine
<i>P. canariensis</i>	Canary Island Pine
<i>P. Elliottii</i>	Slash Pine
<i>P. halepensis</i>	Aleppo Pine
<i>P. nigra</i>	Austrian Pine
<i>P. palustris</i>	Longleaf Pine
<i>P. radiata</i>	Monterey Pine
<i>P. strobus</i>	Eastern White Pine
<i>P. sylvestris</i>	Scotch Pine
<i>P. taeda</i>	Loblolly Pine
<i>P. Thunbergiana</i>	Japanese Black Pine
<i>P. Virginiana</i>	Virginia Pine
<i>Pistachio spp.</i>	Pistachio*
<i>Pittosporum rhombifolium</i>	Queensland Pittosporum
<i>P. tobira</i>	Japanese Pittosporum
<i>Podocarpus macrophyllus</i>	Japanese Yew
<i>Prunus laurocerasus</i>	English Laurel
<i>Prunus spp.</i>	Almond, Apricot, Nectarine, Peach, Plum and Prune*
<i>Pseudotsuga menziesii</i>	Douglas Fir**
<i>Pyracantha coccinea</i>	Firethorn (Scarlet)
<i>P. fortuneana</i>	Firethorn
<i>P. koidzumii</i>	Firethorn
<i>Pyrus spp.</i>	Bradford Pear spp.
<i>Quercus Shumardii</i>	Oak, Shumard's Red
<i>Quercus spp.</i>	Oak species
<i>Raphiolepis indica</i>	Indian Hawthorne
<i>Raphiolepis umbellata</i>	Yedda Hawthorne
<i>Rhododendron</i> (including Azalea)	'Coral Bells,' 'Formosa,' 'Hino-crimson,' 'PJM' 'Delaware Valley White,' 'Roseum Elegans,' 'Flame Creeper,' 'Girard Crimson', 'Wakeiebisu 'White Gumbo', 'George L. Tabor'
<i>Rosa banksiae</i>	Lady Bank's Rose
<i>Rosmarinus officinalis</i>	Rosemary*
<i>Rudbeckia spp.</i>	Black-Eyed Susan; Goldsturm
<i>Santolina virens</i>	Santolina
<i>Saxifraga spp.</i>	Saxifrage; Purple Dome
<i>Scabiosa spp.</i>	Pincushion Flower
<i>Sedum album</i>	Stonecrop
<i>Sedum caudicola</i>	Stonecrop; Lidakense
<i>Sedum dasyphyllum</i>	Stonecrop
<i>Sedum spurium</i>	Stonecrop; Dragon's Blood
<i>Spiraea bumalda</i>	Spirea; Anthony Waterer
<i>Syzygium paniculatum</i>	Australian Brushcherry
<i>Taxus cuspidata</i>	Japanese Yew
<i>T. media</i>	Yew
<i>Teucrium spp.</i>	Germander
<i>Thalictrum diptercarpum</i>	Meadow Rue

[Text in brackets and italics are notes to the label editor.]

[Text in brackets alone are optional marketing statements or alternative statements.]

PRODIAMINE	GROUP	3	HERBICIDE
-------------------	--------------	----------	------------------

<i>Thuja occidentalis</i>	American Arborvitae
<i>Trachelospermum asiaticum</i>	Star Jasmine
<i>Tsuga canadensis</i>	Canada Hemlock
<i>Tulipa spp.</i>	Tulip species
<i>Veronica spp.</i>	Veronica, Speedwell; Sunny Border
<i>Viburnum japonicum</i>	Japanese Viburnum
<i>V. odoratissimum</i>	Sweet Viburnum
<i>V. plicatum</i>	Japanese Snowball
<i>V. rigidum</i>	Canary Island Viburnum
<i>V. suspensum</i>	Arrowwood Viburnum
<i>V. tinus</i>	Laurustinus
<i>V. trilobium</i>	Cranberry Bush
<i>V. wrightii</i>	Leatherleaf Viburnum
<i>Vinca major</i>	Greater Periwinkle, Vinca
<i>Vinca Minor</i>	Common Periwinkle, Vinca
<i>Vitis spp.</i>	Grape*
<i>Weigela florida</i>	Old Fashioned Weigela
<i>Yucca aloifolia</i>	Spanish Bayonet, Yucca
<i>Y. filametosa</i>	Adam's Needle, Yucca

*Do not use on food producing trees, vines or plants.

**Not for use on container grown plants.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a dry place. Store in locked area out of reach of children and domestic animals.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **IF EMPTY:** Place in trash or offer for recycling, if available. **IF PARTLY FILLED:** Call your local solid waste agency for disposal instructions. Never place any unused product down any indoor or outdoor drain or toilet.

CONDITIONS OF SALE AND WARRANTY

The DIRECTIONS FOR USE of this product are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Injury/ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of Control Solutions, Inc. or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

Control Solutions, Inc. warrants that this product conforms to the chemical description on the label, and is reasonably fit for the purposes set forth in the DIRECTIONS FOR USE, when it is used in accordance with such directions, subject to the inherent risks mentioned above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, CONTROL SOLUTIONS, INC. NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESSED OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. THIS WARRANTY EXTENDS TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNING, OR CAUTIONS. BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING

[Text in brackets and italics are notes to the label editor.]

[Text in brackets alone are optional marketing statements or alternative statements.]

PRODIAMINE	GROUP	3	HERBICIDE
-------------------	--------------	----------	------------------

FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. CONTROL SOLUTIONS, INC. and the seller offer this product, and the Buyer and User accept it, subject to the foregoing CONDITIONS OF SALE AND WARRANTY.

Optional Marketing Language:

[PREVENT!]

[PREVENT! 15,000], [PREVENT! 10,000], [PREVENT! 5,000]

[Weed Grass Preventer and Turf Food]

[With Iron and Sulfur Plus Slow Release Nitrogen]

[Covers [5,000], [10,000], [15,000] Square Feet]

[For Pre-Emergence Control of: [Crabgrass], [Goosegrass], [Foxtail], [Barnyardgrass] [& other listed Annual Grass and Broadleaf Weeds] in Established Turfgrass]

[STOPS MANY WEED GRASSES BEFORE THEY GROW]

[PREVENTS CRABGRASS, ANNUAL BLUEGRASS, FOXTAIL]

[Loveland Turf Network]

[Loveland Serving the Environment of Man]

[GREEN GRASS [Cleans our Air, Cools our Earth, Controls Erosion, Fights Pollution, Purifies our Water]]

[Call [the Loveland Turf Network] For Fertilizers - Seed - Sod - Rock – Trees]

[We're the if not, Why not Golf Course Family of the Midwest - Pioneers in grass for over three generations. Suppliers to cities, golf course, parks, home and industry. The 1st purchase sells the 2nd!]

[Visit www.grasspad.com] Omaha, Nebraska: 402/331-5262 Kansas/Missouri: 800-922-4100]

Optional State Specific Language:

[Help do your part in protecting our rivers and lakes by following label directives. Do not apply near water storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back onto your lawn. Do not apply to frozen ground.]

{Label may also include state specific language regarding Nitrogen application}