01/09/2006

1/10



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs Registration Division (7505C) 1200 Pennsylvania Ave., N.W. Washington, D.C. 20460

PA Reg. Number:	Date of Issuance

53883-184

JAN 9 2006

NOTICE OF PESTICIDE:

X Registration
Reregistration
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

PPZ 1.55

Date:

1/9/06

Name and Address of Registrant (include ZIP Code):

Control Solutions, Inc. 5903 Genoa-Red Bluff Pasadena, TX 77507-1041

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby conditionally registered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

- 1. Submit and/or cite all data required for registration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
 - 2. Make the following label changes before you release the product for shipment:
 - a. Revise the EPA Registration Number to read, "EPA Reg. No. 53883-184."

Continued on page 2

Signature of Approving Official:

Mary L. Waller, Product Manager (21)

Fungicide Branch, Registration Division (7505C)

EPA Form 8570-6

Page 2 EPA Reg. No. 53883-184.

- 3. Submit acceptable Storage Stability (OPPTS No. 830.6317) and Corrosion Characteristics (OPPTS No. 830.6320) studies within 15 months of the date of this registration.
- 4. Submit one copy of the revised final printed label for the record before you release the product for shipment.

A copy of the product chemistry (dated 10/18/05) and acute toxicity (dated 12/07/05) reviews are included for your records.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

Sincerely,

Mary L. Waller

Product Manager (21)

Fungicide Branch

Registration Division (7505C)

Mary Llaller

Enclosure



ACCEPTED with COMMENTS In EPA Letter Dated:

JAN 9 2006

PPZ 1.55

Fungicide

53883-84

[] a	lenotes	optional	marketing	language
------	---------	----------	-----------	----------

- [Broad spectrum and systemic disease control on roses, flowers, home lawns, trees, and shrubs]
- [Water- Based]
- [Controls black spot, blights, brown patch, powdery mildew, rust, scab, and more.]
- [Prevents and controls major diseases on roses, flowers, lawns, trees, and shrubs]
- [Concentrate- Just mix with water and spray]
- [For use in pressure type or hose-end sprayers]
- [Enough to treat a 1,333 sq. ft. lawn]
- Absorbed into the plant-After spray has dried, cannot be washed off by sprinklers or rain.]
- [Not affected by rain after spray has dried]
- [Systemic disease control]
- [Systemic Action- Can't be washed off by rain after spray has dried]
- [Prevents a large variety of diseases on home lawns and ornamentals]
- [Prevents lawn diseases before they happen]
- [Broad spectrum disease control on home lawns and ornamentals]
- [Works up to 4 weeks for certain diseases]
- [Encourages increased turf footing]
- [Prevents and controls...
 - Black Spot

- Brown Patch
- and more!!

- Dollar Spot
- Powdery Mildew

Rust

• Leaf Spot]

KEEP OUT OF REACH OF CHILDREN WARNING

See additional precautionary statements on back panel.

Active Ingredient:	
Propiconazole (CAS No. 60207-90-1)	1.55%
Other Ingredients:	98.45%
Total:	100.00%

NET	CONTENTS	

DIDOT AID				
FIRST AID				
Have the product container or label with you when calling a poison control center or doctor, or				
	going for treatment.			
IF IN EYES	Hold eye open and rinse slowly and gently with water for 15-20 minutes.			
	• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye			
	Call a poison control center or doctor for treatment advice			
IF	Call a poison control center or doctor immediately for treatment			
SWALLOWED	advice.			
	 Have person sip a glass of water if able to swallow. 			
	Do not induce vomiting unless told to do so by a poison control			
	center or doctor.			
	Do not give anything by mouth to an unconscious person.			
IF INHALED	Move person to fresh air.			
	If person is not breathing, call 911 or an ambulance, then give			
	artificial respiration, preferably mouth-to-mouth if possible.			
	• Call a poison control center or doctor for further treatment advice.			
IF ON SKIN OR	Take off contaminated clothing.			
CLOTHING	Rinse skin immediately with plenty of water for 15-20 minutes.			
11	Call a poison control center or doctor for treatment advice.			
Note to Physician:	If ingested, induce emesis or lavage stomach. Treat symptomatically.			

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

Warning- Causes substantial, but temporary eye injury. Do not get in eyes. Wear goggles or safety glasses. Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with skin or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Do not allow people or pets to contact treated areas until spray is dry.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment wash water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product must not be used in the production of agricultural commodities grown for resale.

READ ALL INSTRUCTIONS INSIDE BEFORE USE.



FOLLOW ALL DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL. POOR DISEASE CONTROL OR INJURY TO DESIRABLE PLANTS MAY RESULT FROM IMPROPER USE OF THIS PRODUCT.

DO NOT ALLOW PEOPLE OR PETS INTO TREATED AREAS UNTIL SPRAY HAS DRIED.

GENERAL INFORMATION

PPZ 1.55 is a water- based systemic fungicide which effectively prevents and controls major diseases on roses, flowers, lawns, trees, and shrubs.

PPZ 1.55 is very effective when used for disease prevention or applied at the first sign of disease. Best results occur when treating early in the season. If disease is already present, PPZ 1.55 will not eliminate it.

PPZ 1.55, when used as directed, is safe to plants and controls the diseases listed on this label. The use on plants or diseases not listed on the label may result in plant injury and poor disease control.

Do not apply this product to African violets, begonias, Boston ferns, or geraniums.

Contact your local County Extension Service if you need additional information about lawn and ornamental diseases.

MIXING INSTRUCTIONS

- Shake well before using.
- Use a concentrate plus water-style sprayer, a pump-up pressure sprayer, or a dial-style hose-end sprayer for application. It is not recommended to use with hose-end sprayers that only have an on/off switch or those which are continuously filled with water during application.

Food utensils, such as measuring cups and spoons, must not be used for food purposes after use in measuring pesticides.

NOTE: ½ fl. oz. = 1 Tablespoon 1 fl. oz. = 2 Tablespoons

Concentrate Plus Water Style Sprayer

- -Determine the amount of concentrate needed from the chart below.
- -Pour concentrate into the sprayer jar to the specified "fl. oz." level.
- -Next add the necessary amount of water to the "gals." level.
- -Put the sprayer top back on the jar and shake well before spraying.

Pump-Up Spraver

- -Fill the sprayer with the appropriate amount of water.
- -Add the amount of concentrate stated in the chart below.
- -Shake well before application.



Dial-Style Hose-End Sprayer

- -Put the sprayer on the "fl. oz./gallon" level stated in the chart below.
- -Use only enough product necessary for the area that is to be treated.
- -Close the sprayer.
- -Attach the sprayer to the hose and apply the product to the lawn.

(For example: To treat a 1,000 sq. ft. lawn for Powdery Mildew pour 12 fl. oz. into sprayer jar, and set sprayer dial at 3 fl. oz. per gallon)

APPLICATION RATES:

ROSES AND FLOWERS

-Apply product to the point of run off.

-Before watering, allow spray to fully dry.

Plant	Disease	How Much Product To Apply	Comments
Roses	Powdery Mildew	1 Tablespoon (½ fl. oz.) per gal. of water	For best results, thorough coverage is needed. Begin application at bud break. Treat every 10 days.
	Black Spot, Rust	2 tablespoons (1 fl. oz.) per gal. of water	Treat every 10 days. Note: Under severe conditions, 100% control of black spot may not be achieved.
Calendula	Powdery Mildew	l Tablespoon (½ fl. oz.) per gal. of water	Apply every 30 days.
Chrysanthemums	Ray Blight	2 Tablespoons (1 fl. oz.) per gal. of water	Apply every 7 days.
Delphinium	Powdery Mildew	1 Tablespoon (½ fl. oz.) per gal. of water	Apply every 14 days.
Snap Dragons	Rust	2 Tablespoons (1 fl. oz.) per gal. of water	Apply every 10 days.
Sweet William	Leaf Spot	1 Tablespoons (½ fl. oz.) per gal. of water	Apply every 14 days.

LAWNS

-Apply the product to the lawn after mowing or allow the area sprayed to dry completely prior to mowing.

-Prior to application, measure the length and width of area to be treated. Length x width = square feet. (For example, an area 30 ft. long and 20 ft. wide is 600 sq. ft.)

Established Lawns

Disease	How Much Product to Apply	Comments
Brown Patch	5 fl. oz. in 1 gal. of water per 250 sq. ft. of lawn	Repeat treatment every 10 days if necessary. First application should take place in May or June before disease is present. For southern lawns, make applications at first sign of disease in spring or fall.
Dollar Spot, Powdery Mildew, Rust, Anthracnose	3 fl. oz. in 1 gal. of water per 250 sq. ft. of lawn	Apply every 14 days.
Red Thread, Dichondra Rust on Dichondra Lawns	5 fl. oz. in 1 gal. of water per 250 sq. ft. of lawn	Repeat every 21 days, if needed, when conditions are favorable for disease to occur.
Stripe Smut	3 fl. oz. in 1 gal. of water per 250 sq. ft. of lawn	Apply in early Spring before grass starts to grow or in Fall after grass is dormant.

New Seedings

How Much Product to Apply	Comments
5 fl. oz. in 2 gals. of water per 500 sq ft. of	For faster root development and top growth, apply at the 2 to 3-leaf stage of growth.
lawn	



TREES AND SHRUBS

-Apply product to the point of run off.
-Before watering plants, allow spray to fully dry.

Plant	Disease	How Much Product to Apply	Comments
Crab Apple	Scab, Rust	½ Tablespoon (¼ fl. oz.) per gal. of water	Apply when buds just begin to break open. Scab-apply every
Note: Do not use fruit from treated crab apples for			14 days. Rust-make 3 applications, beginning when buds just begin to break open. Apply at 7-day intervals.
food purposes.	Powdery Mildew	1 ½ Tablespoons (¾ fl. oz.) per gal. of water	Apply every 14 days
Crape Myrtle	Powdery Mildew	3 tablespoons (1½ fl. oz.) per gal. of water	Apply every 21 days.
Dogwood	Anthracnose	1 Tablespoon (½ fl. oz.) per gal. of water	Apply every 14-28 days.
Douglas Fir, Shasta Fir	Needle Rust	3 Tablespoons (1 ½ fl. oz.) per gal. of water	Douglas Fir-apply every 21 days. Shasta Fir-apply in May-June before disease occurs.
Jeffrey Pine	Tip Blight	3 Tablespoons (1½ fl. oz.) per gal. of water	Make the first application in mid-late winter (NovJan.) Apply every 2 months up to 3 times
Oak, Red Tip Photinia	Leaf Spot	3 Tablespoons (1½ fl. oz.) per gal. of water	Apply every 14 days.

•

NONBEARING FRUIT AND NUT TREE

-Nonbearing fruit and nut trees are those trees that will not produce fruit for at least one year after application of this product.

-Apply product to the point of run off.

Plant	Disease	How Much Product	Comments
		to Apply	
Apple	Scab	½ Tablespoon (¼ fl. oz.)	Begin applications when
		per gal. of water	leaves start growing in the
	•		Spring and spray every 14
			days.
	Rust	½ Tablespoon (1/4 fl. oz.)	Make 3 applications
		per gal. of water	beginning when buds just
			begin to break open. Apply
			at 7-day intervals
	Powdery Mildew	1 tablespoon (½ fl. oz.)	Begin applications in early
		per gal. of water	Spring and spray every 14
· 			days.
Cherry	Powdery Mildew,	1 Tablespoon (½ fl. oz)	Begin applications in early
	Leaf Spot	per gal. of water	Spring and spray every 21
			days.
Citrus	Greasy Spot	4 Tablespoons (2 fl. oz.)	Apply June- August every 21
		per gal. of water	days.
Pecan	Scab, Powdery	2 Tablespoons (1 fl. oz.)	Begin applications at bud
÷	Mildew, Downy Spot,	per gal. of water	break and make 3
	Zonate Leaf Spot,		applications at 14-day
	Fungal Leaf Scorch,		intervals.
	Brown Spot		
Walnut	Anthracnose	2 Tablespoons (1 fl. oz)	Apply every 14 days.
		per gal. of water	

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in cool, dry area away from children and domestic animals. **Pesticide Disposal:** If empty: Do not reuse this container. Do not rinse unless required for recycling. Place in trash. If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Unless otherwise instructed, place in trash. Never pour unused product down the drain or on the ground.

WARRANTY STATEMENT

Control Solutions, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with the Directions For Use. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use



or application, all of which are beyond the control of Control Solutions, Inc. To the extent allowed by law, Control Solutions, Inc. shall in no event be liable for consequential, special, or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. In addition to the foregoing, no purchaser of this product (other than an end user) shall be entitled to any reimbursement for any loss suffered as a result of any suspension or cancellation of the registration for this product by the U.S. Environmental Protection Agency. Except, as expressly provided herein, Control Solutions, Inc. makes no warranties, guarantees, or representations of any kind, either expressed or implied, or by usage of trade, statutory or otherwise, with regard to the product sold, including, but not limited to merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage. The exclusive remedy of any buyer or user of this product for any and all losses, injuries, or damage resulting from or in any way arising from the use, handling, or application of this product, whether in contract, warranty, tort, negligence, strict liability, or otherwise, shall be damages not exceeding the purchase price paid for this product or, at Control Solutions, Inc. election, the replacement of this product. [optional warranty statement]

Note: Buyer assumes all responsibilities for safety and use not in strict accordance with directions. *[optional warranty statement]*

Control Solutions, Inc. 5903 Genoa-Red Bluff Pasadena, TX 77507-1041 EPA Reg. No. 53883-XXX EPA Est. No.

٠.