

PLEASE NOTE

This image contains more than one label approved for this product on this date.

53883-195

03/11/2008

1/8



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

March 11, 2008

Jane M. Miller,
Agent to Control Solutions, Inc.
5903 Genoa-Red Bluff
Pasadena, Texas 77507

Subject: Amendment: Add Me-Too Uses to Labeling
Lambda LP Soil & Turf Insect Killer Granules
EPA Reg. Nos. 53883-195
Application dated December 01, 2007


Dear Ms. Miller:

The above referenced labeling, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable subject to the following comments:

1. First page, directly under product's name: delete the last phrase "*and similar type areas*" from the end of statement, or list the "other similar type areas".
2. On the front panel, add the referral statement "See additional precautionary statements and directions for use on side/back panel".
3. Throughout the label, delete all references to, and use instructions for "fire ants" and "red imported fire ants". These uses/pests are not listed on the cited me-too product.
4. Under Optional Marketing Statements, delete "*Controls 30+ insects*", or list all the insects and change statement to "*Controls listed insects*".
5. Delete "*interior plantscapes*" from the first statement under General Information: Lawn and Perimeter Pest Control.
6. The container disposal statements for pesticide products have changed per PR Notice 2007-4; revise appropriately.

Submit a formulators exception statement; this form is required with amendment to add new sites/pests. Please revise labeling accordingly and submit final printed label to the Agency prior to releasing the product for shipment. See enclosed copy of the stamped label for your record. If you have any questions please contact Dr. B.A. Akinlosotu at (703) 605-0653.

Sincerely,



George T. LaRocca,
Product Manager 13
Insecticide Branch
Registration Division (7505P)

Enclosure
baa/

Lambda LP
Soil & Turf Insect Killer Granules

[alt. name: Cyonara LP Lawn & Perimeter Insect Control Granules]

For Use around buildings and structures for the control of listed pests, including on lawns, around ornamental trees and shrubs, around residential, institutional, public, commercial, agricultural and industrial buildings; and parks, recreational areas, athletic fields, and similar type areas.

[Optional Marketing Statements -

- 10 pounds treats up to 5,000 sq. ft.
- Kills in 24 hours
- Kills [Controls] many types of insects, including ants, fleas, chinch bugs & grubs
- Broadcast with spreader
- Resealable Bag
- For Perimeter Treatment Around Your House Foundation
- Kills Ants, Cockroaches, Fleas, Ticks and other listed insects
- Convenient Shaker Bag
- No Spreader
- Controls Chinch Bugs, Ticks, and Fleas
- Controls 30+ insects
- Use on lawns and as a perimeter treatment around your house foundation]

ACCEPTED
with COMMENTS
in EPA Label Dated

MAR 11 2008
Under the Federal Insecticide, Fungicide, and Rodenticide Act
53883-195

Active Ingredient:

Lambda-cyhalothrin 0.04%

Other Ingredients: 99.96%

Total 100.00%

Net Weight _____

KEEP OUT OF REACH OF CHILDREN
CAUTION/PRECAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall® International for emergency medical treatment at (866) 897-8050.	
If swallowed:	<ul style="list-style-type: none"> • Call a Poison Control Center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a Poison Control Center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a Poison Control Center or doctor for treatment advice.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION - Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly after handling and before eating or smoking. Food utensils such as spoons or measuring cups must not be used for food purposes after use in measuring pesticides.

ENVIRONMENTAL HAZARDS

This product is highly toxic to fish and aquatic vertebrates. Do not apply directly to water.

READ ENTIRE LABEL BEFORE USE

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Keep children and pets off treated area until product has been watered in and lawn is dry.

Restrictions:

- Do not apply this product to edible growing crops or stored raw agricultural commodities used for food or feed.
- Do not allow applications to contact water inhabited by fish, such as in ornamental fish ponds that are located around structures being treated.
- Do not apply this product by aerial application.
- Use of this pesticide adjacent to water may affect aquatic organisms. To protect these organisms, do not apply this pesticide within 25 ft. of lakes, reservoirs, rivers, permanent streams, marches or natural ponds, estuaries, and commercial fish farm ponds.
- In New York State, this product may not be applied within 100 feet of a coastal marsh, or stream that drains directly into a coastal marsh.
- Observe all restrictions and precautions which appear on the labels of these products.
- Do not apply more than 0.36 lbs. of the a.i. (800 pounds of Lambda LP) per acre per year.

IMPORTANT: Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, outdoor ornamental gardens or parks, or on lawns or grounds.

GENERAL INFORMATION

LAWN AND PERIMETER PEST CONTROL

Lambda LP insecticide may be used for applications to maintain interior plantscapes or outdoor areas where turf and ornamental ground covers are grown such as residential landscaped areas and non residential landscapes around institutional, public, commercial and institutional buildings, parks, recreational areas, and athletic fields. Broadcast Lambda LP Insecticide with suitable application equipment to ensure uniform coverage over the treatment area. The application rates listed in the **Lawn and Perimeter Use Rate** table will provide excellent control of the respective pests under typical conditions.

NOTE:

- Use higher rate for longer length of control.
- Watering (irrigation or rainfall) following application will dissolve the granules and release the insecticide.
- Make no more than 6 general applications per year. More frequent treatments should be limited to spot treatments.
- Uniform coverage is necessary to provide the most effective level of control.
- Consult your state university or local Cooperative Extension Service office for specific pest control application timing in your area.

5/8

- Application to turf should be started prior to the establishment of high insect pest populations and significant turf damage. Make reapplications as necessary to keep pest populations under control, using higher rates as pest pressure increases.

Recommended Application Rates for Lawn and Perimeter Use Rate

	PEST	A.I. lb per acre	Lambda LP Rate Per 1000 ft2	Lambda LP Rate Per Acre
Band Treatment Around Foundation	Ants Beetles Crickets Earwigs Firebrats Millipedes Mites Silverfish Sowbugs	0.04-0.06	2 - 3 lbs.	88 - 131 lbs.
Overall Lawn Application	Ants Armyworms Centipedes Crickets Cutworms Earwigs Grasshoppers Green June Beetles Japanese Beetle (Adult) Leathoppers Millipedes Mites Pillbugs Sod webworms Spider Mites (Twospotted, Spruce) Spittlebugs Springtails and Ticks (including Ticks which may transmit Lyme Disease)	0.04-0.06	2 - 3 lbs.	88 - 131 lbs.
Overall Lawn Application	Billbugs Chinch Bugs Dung beetles* Green June beetle grubs* Hyperodes weevils Japanese beetle grubs* Mole Crickets	0.06-0.078	3 - 4 lbs	131 - 176 lbs.
Spot Treatment	Ants		Apply 1-2 teaspoons around each ant hill	

6/8

Fire Ants		Mound Treatment: Apply ½ cup over, and 2 ft. around each fire ant mound	
-----------	--	---	--

* Suppression only.

SPECIFIC USE INSTRUCTIONS
Treatment Around Foundations

Around house foundation to control pests normally found around or near foundations of houses, as well as in lawns, and certain of these pests that may enter houses.

Pests	Use Rate	Use Instructions
Ants (Argentine, Southern, Field, Allegheny Mound, Florida Carpenter, Black Carpenter, Cornfield, Honey, Pavement, Red Imported Fire Ant, Nuisance), Fleas Roaches (German and American), Palmetto bugs Waterbugs Millipedes Mites Silverfish Sowbugs (Pillbugs) Crickets Earwigs	Apply at the rate of 2 lb to 3 lb per 1,000 sq. ft. (an area 5' x 200') oz. or 3.2 oz. to 4.8 oz. per 100 sq. ft. (10' x 10').	Apply a 3 to 10 foot band of granules next to the house foundation and completely distribute the granules. Water lightly immediately following application. Treat when insects first appear or when you notice lawn damage. For best results, leaf litter and debris should be removed from the foundation area. Branches and vegetation that touch the foundation should be cut back. Other potential pest harborage sites, such as deep mulch, should be raked away before application.

Overall Lawn Application

Pests	Use Rate	Use Instructions
Ants (Argentine, Southern, Field, Allegheny Mound, Florida Carpenter, Black Carpenter, Cornfield, Honey, Pavement, Red Imported Fire Ant, Nuisance) Armyworms (Fall & Yellow-striped) Centipedes European Crane Flies Crickets Cutworms Earwigs Fleas Lawn moths (sod webworms) Grasshoppers Japanese beetle (adults) Pillbugs Spittle bugs Mites Green June Beetles Leafhoppers Millipedes Springtails and Ticks (including Ticks which may transmit Lyme Disease)	Apply at the rate of 2 lb to 3 lb per 1,000 sq. ft. (an area 20' x 50') or 3.2 oz. to 4.8 oz. per 100 sq. ft. (10' x 10').	Apply with a fertilizer spreader (see spreader setting chart). Apply when grass is dry and water the lawn lightly immediately following application. Treat when insects first appear or when you notice lawn damage.

7/8

<p>Billbugs Grubs (Japanese beetle grubs, Green June Beetle Grubs) <i>Ataenuis spretulus</i> (dung beetles) Hyperodes Weevils Mole Crickets Chinch bugs</p>	<p>Apply 3 lb. per 1,000 sq. ft. of lawn.</p>	<p>Water grass completely immediately following application.</p> <p>Treat when insects first appear or when you notice lawn damage.</p> <p>For hyperodes weevils, apply in April-May.</p> <p>For billbug control, apply in May when activity is first observed or when chewed or brown grass indicates damage from these insects.</p> <p>For grub control, except on dung beetle, treat any time between late July and early October.</p> <p>For dung beetle grub control, apply in early spring or early summer (March-July) when damage is first noticed. Immediately following application, completely water grass with ¼ to ½ inch of water. If a second generation occurs, treat again in late summer or early fall.</p> <p>For chinch bug control, apply at higher rates during the summer when both nymphs and adults may be present.</p> <p>For mole crickets apply in the early spring for control of adults when actively invading grass areas. If the soil is dry, water before application to bring mole crickets closer to the surface and in contact with the granules. Apply late in the day and water in following application. For control of early nymphal stages, apply at the time of peak egg hatch. As the nymphs mature, higher rates may be necessary.</p>
---	---	--

Spot Treatment

Pests	Use Rate	Use Instructions
<p>Ants (Argentine, Southern, Field, Allegheny Mound, Florida Carpenter, Black Carpenter, Cornfield, Honey, Pavement, Red Imported Fire Ant, Nuisance)</p>	<p>Mound Treatment: Apply 1-2 teaspoons over and around each ant hill.</p>	<p>Water lightly immediately following application.</p> <p>NOTE: Use both broadcast and spot treatment for difficult to large ant infestations. If soil is dry, water lightly before application to bring the ants closer to the surface for better control.</p>
<p>Fire Ants</p>	<p>Mound Treatment: Apply ½ cup over, and 2 ft. around, each fire ant mound.</p>	<p>Immediately following application, water in completely. Apply a minimum of 1 gallon of water to each mound area. To prevent the disturbance of ants, water gently. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness. NOTE: For best results, apply in cool weather, 65°-80° F, or in early morning or late evening hours. Treat new mounds as they appear. Fast acting. Works within 24 hours.</p>

8/8

Spreader Settings for Lawn Application

[INSERT SPREADER SETTING CHART]

NOTE: The spreader settings stated are approximate. Precisely weigh a few pounds into your spreader and apply to a measured area. To assure correct rate of application, adjust spreader settings if needed.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Reclose container tightly after each use. Store in a cool, dry area which is inaccessible to children and pets.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Paper and plastic bags: completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by other procedures allowed by state and local authorities.

WARRANTY STATEMENT

CONTROL SOLUTIONS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated in the Directions for Use when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or CONTROL SOLUTIONS, INC., and Buyer and User assume the risk of any such use.

Control Solutions, Inc.
5903 Genoa-Red Bluff
Pasadena, TX 77507-1041

EPA Reg. No. 53883-195

EPA Est. No. xxxxxx-xx-xxx

NEXT

LABEL

53883-195

03/11/2008

1/5



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

March 11, 2008

Jane M. Miller,
Agent to Control Solutions, Inc.
5903 Genoa-Red Bluff
Pasadena, Texas 77507

Subject: Supplemental Labeling to Add Me-Too Uses.
Lambda LP Lawn & Perimeter Insect Control Granules
EPA Reg. Nos. 53883-195
Submissions/Applications dated December 01, 2007 & January 24, 2008

Dear Ms. Miller:

The above referenced amendment, submitted in connection with registrations under FIFRA Section 3(c)(7)(B), as amended, is acceptable subject to the following comments.

This accepted supplemental labeling is subject to the same label revisions as indicated in our letter (dated March 11, 2008) pertaining to the main label, "Lambda LP Soil & Turf Insect Killer Granules", such as:

- Throughout the label, delete all references to, and use instructions for "**fire ants**" and "**red imported fire ants**". These uses/pests are not listed on the cited me-too product.
- Delete "**interior plantscapes**" from the first statement under General Information: Lawn and Perimeter Pest Control.
- The container disposal statements for pesticide products have changed per PR Notice 2007-4; revise appropriately.

Revise the labeling accordingly and submit copy of the final printed label to the Agency prior to releasing the product for shipment. At your next label printing, or within eighteen (18) months, whichever comes first, please incorporate this supplemental labeling into the product's main labeling. See enclosed stamped label for your records. If you have any questions concerning this action, please contact Dr. B. A. Akinlosotu at (703) 605-0653.

Sincerely,

A handwritten signature in black ink, appearing to read "George T. LaRocca".

George T. LaRocca,
Product Manager 13
Insecticide Branch
Registration Division (7505P)

Enclosure:

2/5

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR **LAMBDA LP** BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using **LAMBDA LP** as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

LAMBDA LP
Lawn & Perimeter Insect Control Granules

EPA Reg. No. 53883-195

**KEEP OUT OF REACH OF CHILDREN
CAUTION/PRECAUCION**

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

MAR 11 2008
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
53883-195

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

In case of an emergency with this product contact SafetyCall® International at (866) 897-8050.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of pesticide application.

Keep children and pets off treated area until product has been watered in and lawn is dry.

Restrictions:

- Do not apply this product to edible growing crops or stored raw agricultural commodities used for food or feed.
- Do not allow applications to contact water inhabited by fish, such as in ornamental fish ponds that are located around structures being treated.
- Do not apply this product by aerial application.
- Use of this pesticide adjacent to water may affect aquatic organisms. To protect these organisms, do not apply this pesticide within 25 ft. of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds.
- In New York State, this product may not be applied within 100 feet of a coastal marsh, or stream that drains directly into a coastal marsh.
- Observe all restrictions and precautions which appear on the labels of these products.
- Do not apply more than 0.36 lbs. of the a.i. (800 pounds of LAMBDA LP) per acre per year.

IMPORTANT: Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior landscapes, outdoor ornamental gardens or parks, or on lawns or grounds.

GENERAL INFORMATION

LAWN AND PERIMETER PEST CONTROL

Lambda LP insecticide may be used for applications to maintain interior plantscapes or outdoor areas where turf and ornamental ground covers are grown such as residential landscaped areas and non residential landscapes around institutional, public, commercial and institutional buildings, parks, recreational areas, and athletic fields. Broadcast Lambda LP Insecticide with suitable application equipment to ensure uniform coverage over the treatment area. The application rates listed in the **Lawn and Perimeter Use Rate** table will provide excellent control of the respective pests under typical conditions.

NOTE:

- Use higher rate for longer length of control.
- Watering (irrigation or rainfall) following application will dissolve the granules and release the insecticide.
- Make no more than 6 general applications per year. More frequent treatments should be limited to spot treatments.
- Uniform coverage is necessary to provide the most effective level of control.
- Consult your state university or local Cooperative Extension Service office for specific pest control application timing in your area.
- Application to turf should be started prior to the establishment of high insect pest populations and significant turf damage. Make reapplications as necessary to keep pest populations under control, using higher rates as pest pressure increases.

Recommended Application Rates for Lawn and Perimeter Use

	PEST	A.I. lb per acre	Lambda LP Rate Per 1000 ft ²	Lambda LP Rate Per Acre
Band Treatment Around Foundation	Ants Crickets Firebrats Mites Sowbugs Beetles Earwigs Millipedes Silverfish	0.04-0.06	2 - 3 lbs.	88 - 131 lbs.
Overall Lawn Application	Ants Armyworms Centipedes Crickets Cutworms Earwigs Grasshoppers Green June Beetles Japanese Beetle (Adult) Leafhoppers Millipedes Mites Pillbugs Sod webworms Spider Mites (Twospotted, Spruce) Spittlebugs Springtails and Ticks (including Ticks which may transmit Lyme Disease)	0.04-0.06	2 - 3 lbs.	88 - 131 lbs.

Overall Lawn Application	Billbugs Chinch Bugs Dung beetles* Green June beetle grubs* Hyperodes weevils Japanese beetle grubs* Mole Crickets	0.06-0.078	3 - 4 lbs	131 - 176 lbs.
	Spot Treatment Ants		Apply 1-2 teaspoons around each ant hill	
	Fire Ants		Mound Treatment: Apply ½ cup over, and 2 ft. around each fire ant mound	

*Suppression only.

SPECIFIC USE INSTRUCTIONS

Treatment Around Foundations

Pests Controlled: Ants (Argentine, Southern, Field, Allegheny Mound, Florida Carpenter, Black Carpenter, Cornfield, Honey, Pavement, Red Imported Fire Ant, Nuisance), Fleas, Roaches (German and American), Palmetto bugs, Waterbugs, Millipedes, Mites, Silverfish, Sowbugs (Pillbugs), Crickets, Earwigs

Application Rate: Apply at the rate of 2 lb to 3 lb per 1,000 sq. ft. (an area 5' x 200') oz. or 3.2 oz. to 4.8 oz. per 100 sq. ft. (10' x 10').

Application Instructions: For best results, leaf litter and debris should be removed from the foundation area. Branches and vegetation that touch the foundation should be cut back. Other potential pest harborage sites, such as deep mulch, should be raked away before application.

Overall Lawn Application

Chinch Bug and Mole Cricket Control -

For **chinch bug control**, apply at higher rates during the summer when both nymphs and adults may be present.

For **mole crickets** apply in the early spring for control of adults when actively invading grass areas. If the soil is dry, water before application to bring mole crickets closer to the surface and in contact with the granules. Apply late in the day and water in following application. For control of early nymphal stages, apply at the time of peak egg hatch. As the nymphs mature, higher rates may be necessary.

Spot Treatment

Pests Controlled: Ants (Argentine, Southern, Field, Allegheny Mound, Florida Carpenter, Black Carpenter, Cornfield, Honey, Pavement, Red Imported Fire Ant, Nuisance)

Mound Treatment: Apply 1-2 teaspoons over and around each ant hill.

NOTE: Use both broadcast and spot treatment for difficult to large ant infestations. If soil is dry, water lightly before application to bring the ants closer to the surface for better control.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Reclose container tightly after each use. Store in a cool, dry area which is inaccessible to children and pets.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Paper and plastic bags: completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by other procedures allowed by state and local authorities.

Read the "WARRANTY STATEMENT" on the label for Lambda LP before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Control Solutions, Inc.
5903 Genoa-Red Bluff
Pasadena, TX 77507-1041