817



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

> OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

August 18, 2009

Amy Warren, Regulatory Affairs Specialist Control Solutions, Inc. 5903 Genoa-Red Bluff Pasadena, Texas 77507

Subject: Remove Contradictory Language in DFU Lambda LP Soil & Turf Insect Killer Granules EPA Reg. Nos. 53883-195 Application dated May 27, 2009

Dear Ms. Warren:

The above referenced labeling, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

Please note that we have approved the amended labeling for this product with the understanding that it is consistent with the Agency's previously stamped accepted labels. See enclosed copy of the stamped label for your record. If you have any questions concerning this action please contact Dr. B.A. Akinlosotu at (703) 605-0653.

Sincerely imberly Nesci.

Product Manager 13 Insecticide Branch Registration Division (7505P)

Enclosure

Lambda LP Soil & Turf Insect Killer Granules

[alt. name: Cyonara LP Lawn & Perimeter Insect Control Granules] For outdoor use around the home only.

[Optional Marketing Statements -

- 10 pounds treats up to 5,000 sq. ft.
- Kills in 24 hours
- Kills [Controls] many types of insects, including ants, fleas, chinch bugs & grubs
- Broadcast with spreader
- Resealable Bag
- For Perimeter Treatment Around Your House Foundation
- Kills Ants, Cockroaches, Fleas, Ticks and other listed insects Under the Federal Insecticide, Fungicide,
- Convenient Shaker Bag
- No Spreader
- Controls Chinch Bugs, Ticks, and Fleas]

ACCEPTED AUG 1 8 2009 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under:

Active Ingredier Lambda-cyhaloth	nt: EPA. Reg. No: <u>5383-199</u> irin	
Other Ingredien	rotal	
Net Weight	ت رزن س ۹ 	
	KEEP OUT OF REACH OF CHILDREN دمین CAUTION	
	FIRST AID	
If swallowed:	 Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor 	
	Do not give anything by mouth to an unconscious person.	
If in eyes:	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. 	

Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
 Call a Poison Control Center or doctor for treatment advice.

Have the product container or label with you when calling a Poison Control Center or doctor, or

going for treatment. You may also call Poison Control Center 1-800-222-1222.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION - Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly after handling and before eating or smoking. Food utensils such as spoons or measuring cups must not be used for food purposes after use in measuring pesticides. ENVIRONMENTAL HAZARDS

This product is highly toxic to fish and aquatic vertebrates. Do not apply directly to water.

READ ENTIRE LABEL BEFORE USE

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Keep children and pets off treated area until product has been watered in and lawn is dry.

In New York State, this product may not be applied to lawns within 100 feet of a coastal marsh, or stream that drains directly into a coastal marsh.

GENERAL INFORMATION

For use outdoors, on lawns and as a barrier band treatment around house foundation. Kills ants (including Fire Ants), armyworms (Fall and Yellow-striped), billbugs, chinch bugs, crickets, cutworms, earwigs, European crane flies, fleas, grubs, roaches (German and American), sod webworms, ticks and other insect pests listed in the tables below.

To use this product follow these 3 easy steps -(1) Determine the size of the job and measure the area to be treated. (2) Determine the proper setting for your spreader using the spreader setting chart. (3) Treat as directed in rate chart. Complete coverage is important. Water in granules immediately following application.

[Optional: If using Shaker bag, use the following rather than the paragraph above]

To use this product follow these 2 easy steps- (1) Determine the size of the job and measure the area to be treated. (2) Treat as directed in rate chart. Complete coverage is important. Water in granules immediately following application.

NOTE:

• Make no more than 6 applications per year.

- Do not apply within 24 hours of rain

.

- Do not allow product to run off from target area.
- · Do not mow for 24 hours following treatment.

Band Treatment

Around house foundation to control pests normally found around or near foundations of houses, as well as in lawns, and certain of these pests that may enter houses.

Pests	Use Rate	Use Instructions
Ants (Argentine, Southern,	Apply at the rate of	Apply a 3 to 5 foot band of granules next to the
Field, Allegheny Mound,	2 lb per 1,000 sq. ft.	house foundation and completely distribute the
Florida Carpenter, Black	(an area 5' x 200') oz. or	granules. Water lightly immediately following
Carpenter, Cornfield, Honey,	3.2 oz. per 100 sq. ft.	application.
Pavement, Red Imported Fire	(10' x 10').	
Ant, Nuisance), Fleas,		Treat when insects first appear or when you notice
Roaches (German and		lawn damage.
American), Palmetto bugs,		
Waterbugs, Millipedes, Mites,		
Silverfish, Sowbugs		
(Pillbugs), Crickets, Earwigs		

Overall Lawn Application

Pests	Use Rate	Use Instructions
Ants (Argentine, Southern, Field, Allegheny Mound, Florida Carpenter, Black Carpenter, Cornfield, Honey, Pavement, Red Imported Fire Ant, Nuisance), Armyworms (Fall & Yellow-striped), Centipedes, European	For application with other types of equipment, apply at the rate of 2 lb per 1,000 sq. ft. (an area 20' x 50') or 3.2 oz. per 100 sq. ft. (10' x 10').	Apply with a fertilizer spreader (see spreader setting chart). Apply when grass is dry and water the lawn lightly immediately following application. Treat when insects first appear or when you notice lawn damage.

Cutworms, Earwigs, Fleas, Lawn moths (sod webworms), Grasshoppers, Japanese beetle (adults), Pillbugs, Spittle bugs, Mites, Green June Beetles, Leafhoppers, Millipedes, Springtails and Ticks (including Ticks which may transmit Lyme Disease) Billbugs, Grubs (Japanese beetle grubs, Green June	Apply 3 lb. per 1,000 sq. ft. of lawn.	Water grass completely immediately following application.
Beetle Grubs, <i>Ataenuis</i> <i>spretulus</i> (dung beetles), Hyperodes Weevils, Mole Crickets, Chinch bugs		 Treat when insects first appear or when you notice lawn damage. For hyperodes weevils, apply in April-May. For billbug control, apply in May when activity is first observed or when chewed or brown grass indicates damage from these insects. For grub control, except on dung beetle, treat any time between late July and early October. For dung beetle grub control, apply in early spring or early summer (March-July) when damage is first noticed. Immediately following application, completely water grass with ¼ to ½ inch of water. If a second generation occurs,

Spot Treatment

/+ 1

Pests	Use Rate	Use Instructions
Ants (Argentine, Southern, Field, Allegheny Mound, Florida Carpenter, Black Carpenter, Cornfield, Honey, Pavement, Red Imported Fire Ant, Nuisance)	Mound Treatment: Apply 1 ½ teaspoons over and around each ant hill.	Water lightly immediately following application.
Fire Ants	Mound Treatment: Apply ½ cup over, and 2 ft. around, each fire ant mound.	Immediately following application, water in completely. Apply a minimum of 1 gallon of water to each mound area. To prevent the disturbance of ants, water gently. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness. NOTE: For best results, apply in cool weather, 65°-80° F, or in early morning or late evening hours. Treat new mounds as they appear. Fast acting. Works within 24 hours.

Spreader Settings for Lawn Application

[INSERT SPREADER SETTING CHART]

NOTE: The spreader settings stated are approximate. Precisely weigh a few pounds into your spreader and apply to a measured area. To assure correct rate of application, adjust spreader settings if needed.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use close tightly.

In case of spill: Avoid contact, isolate area and keep out animals and unprotected persons. Confine spills.

To confine spill: Cover to prevent dispersal. Place damaged package in a holding container and identify contents.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spilled bait, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance.

Container Disposal: Non-refillable container. Do not reuse or refill this container. Empty container through use according to label instructions. Offer for recycling, if available. If recycling is not available; then dispose of container in a sanitary landfill or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Note: Buyer assumes all responsibilities for safety and use not in strict accordance with directions.

Control Solutions, Inc. 5903 Genoa-Red Bluff Pasadena, TX 77507-1041

EPA Reg. No. 53883-195

EPA Est. No. xxxxxx-xx-xxx