

53883-195

1/4/2010

10 of 12



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Control Solutions, Inc.
c/o Jane M. Miller
5903 Genoa-Red Bluff Road
Pasadena, TX 77507-1041

JAN - 4 2010

Subject: Amendment to Reinstate Uses on Labels
Lambda LP Insecticide Soil & Turf Insect Killer Granules, and Lambda LP Lawn
& Perimeter Insect Control Granules (Supplemental Labeling)
EPA Reg. No. 53883-195
Your Submission Dated October 9, 2009
Decision No: 422194

Dear Ms. Miller:


The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable subject to the comments below. Two (2) copies of the finished labeling must be submitted prior to releasing the product for shipment. A stamped copy of each label is enclosed for your records.

1. Delete the word "vertebrates" from the Environmental Hazards section on page 2 of the label.
2. Under the Precautionary Statements section on page 2 of the label, you must replace the sentence starting with "Wash hands thoroughly..." with the following statement:

"Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet."
3. On page 6 of the primary label, and on page 4 of the supplemental label, replace the word "may" with "must" in the pesticide disposal statement in the Storage and Disposal section.

At your next label printing, or within eighteen (18) months, whichever comes first, you must incorporate the supplemental labeling into the main product labeling. If you have any questions regarding this action, please contact Rosanna Louie at (703) 308-0037.

Sincerely,



Kimberly Nesci
Product Manager 11, Insecticide Branch
Registration Division (7505P)

Enclosures:

- Lambda LP Soil & Turf Insect Killer Granules Label, Stamped *Accepted with Comments*
- Lambda LP Lawn & Perimeter Insect Control Granules Supplemental Label, Stamped *Accepted with Comments*

3 of 12

**Lambda LP
Soil & Turf Insect Killer Granules**

[alt. name: Cyonara LP Lawn & Perimeter Insect Control Granules]

For Use around buildings and structures for the control of listed pests, including on lawns, around ornamental trees and shrubs, around residential, institutional, public, commercial, agricultural and industrial buildings; and parks, recreational areas, and athletic fields.

[Optional Marketing Statements -

- 10 pounds treats up to 5,000 sq. ft.
- Kills in 24 hours
- Kills [Controls] many types of insects, including ants, fleas, chinch bugs & grubs
- Broadcast with spreader
- Resealable Bag
- For Perimeter Treatment Around Your House Foundation
- Kills Ants, Cockroaches, Fleas, Ticks and other listed insects
- Convenient Shaker Bag
- No Spreader
- Controls Chinch Bugs, Ticks, and Fleas
- Use on lawns and as a perimeter treatment around your house foundation]

**ACCEPTED
with COMMENTS
In EPA Letter Dated:
JAN - 4 2010**

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
53883-195**

Active Ingredient:

Lambda-cyhalothrin 0.04%

Other Ingredients: 99.96%

Total 100.00%

Net Weight _____

**KEEP OUT OF REACH OF CHILDREN
CAUTION/PRECAUCION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

See additional Precautionary Statements and Directions for Use on side/back panel.

FIRST AID	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall® International for emergency medical treatment at (866) 897-8050.	
If swallowed:	<ul style="list-style-type: none"> • Call a Poison Control Center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a Poison Control Center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a Poison Control Center or doctor for treatment advice.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION - Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly after handling and before eating or smoking. Food utensils such as spoons or measuring cups must not be used for food purposes after use in measuring pesticides.

ENVIRONMENTAL HAZARDS

This product is highly toxic to fish and aquatic vertebrates organisms. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind and rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

READ ENTIRE LABEL BEFORE USE

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Keep children and pets off treated area until product has been watered in and lawn is dry.

Restrictions:

- Apply this product directly to the lawn or garden area. Water treated area as directed on this label. Do not water to the point of run off.
- Do not make applications during rain.
- Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.
- Do not apply this product to edible growing crops or stored raw agricultural commodities used for food or feed.
- Do not allow applications to contact water inhabited by fish, such as in ornamental fish ponds that are located around structures being treated.
- Do not apply this product by aerial application.
- Use of this pesticide adjacent to water may affect aquatic organisms. To protect these organisms, do not apply this pesticide within 25 ft. of lakes, reservoirs, rivers, permanent streams, marches or natural ponds, estuaries, and commercial fish farm ponds.
- In New York State, this product may not be applied within 100 feet of a coastal marsh, or stream that drains directly into a coastal marsh.
- Observe all restrictions and precautions which appear on the labels of these products.
- Do not apply more than 0.36 lbs. of the a.i. (800 pounds of Lambda LP) per acre per year.

IMPORTANT: Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in outdoor ornamental gardens or parks, or on lawns or grounds.

GENERAL INFORMATION

LAWN AND PERIMETER PEST CONTROL

Lambda LP insecticide may be used for applications to outdoor areas where turf and ornamental ground covers are grown such as residential landscaped areas and non residential landscapes around institutional, public, commercial and institutional buildings, parks, recreational areas, and athletic fields. Broadcast Lambda LP Insecticide with suitable application equipment to ensure uniform coverage over the treatment area. The application rates listed in the **Lawn and Perimeter Use Rate** table will provide excellent control of the respective pests under typical conditions.

NOTE:

- Use higher rate for longer length of control.
- Watering (irrigation or rainfall) following application will dissolve the granules and release the insecticide.
- Make no more than 6 general applications per year. More frequent treatments should be limited to spot treatments.
- Uniform coverage is necessary to provide the most effective level of control.
- Consult your state university or local Cooperative Extension Service office for specific pest control application timing in your area.
- Application to turf should be started prior to the establishment of high insect pest populations and significant turf damage. Make reapplications as necessary to keep pest populations under control, using higher rates as pest pressure increases.

Recommended Application Rates for Lawn and Perimeter Use Rate

	PEST	A.I. lb per acre	Lambda LP Rate Per 1000 ft²	Lambda LP Rate Per Acre
Band Treatment Around Foundation	Ants Beetles Crickets Earwigs Firebrats Millipedes Mites Silverfish Sowbugs	0.04-0.06	2 - 3 lbs.	88 - 131 lbs.
Overall Lawn Application	Ants Armyworms Centipedes Crickets Cutworms Earwigs Grasshoppers Green June Beetles Japanese Beetle (Adult) Leafhoppers Millipedes Mites Pillbugs Sod webworms Spider Mites (Twospotted, Spruce) Spittlebugs Springtails and Ticks (including Ticks which may transmit Lyme Disease)	0.04-0.06	2 - 3 lbs.	88 - 131 lbs.
Overall Lawn Application	Billbugs Chinch Bugs Dung beetles* Green June beetle grubs* Hyperodes weevils Japanese beetle grubs* Mole Crickets	0.06-0.078	3 - 4 lbs	131 - 176 lbs.

60 of 12

Spot Treatment	Ants		Apply 1-2 teaspoons around each ant hill.
	Fire Ants		Mound Treatment: Apply ½ cup over, and 2 ft. around each fire ant mound.

* Suppression only.

SPECIFIC USE INSTRUCTIONS

Treatment Around Foundations

Around house foundation to control pests normally found around or near foundations of houses, as well as in lawns, and certain of these pests that may enter houses.

Pests	Use Rate	Use Instructions
Ants (Argentine, Southern, Field, Allegheny Mound, Florida Carpenter, Black Carpenter, Cornfield, Honey, Pavement, Red Imported Fire Ant, Nuisance), Fleas Roaches (German and American), Palmetto bugs Waterbugs Millipedes Mites Silverfish Sowbugs (Pillbugs) Crickets Earwigs	Apply at the rate of 2 lb to 3 lb per 1,000 sq. ft. (an area 5' x 200') oz. or 3.2 oz. to 4.8 oz. per 100 sq. ft. (10' x 10').	Apply a 3 to 10 foot band of granules next to the house foundation and completely distribute the granules. Water lightly immediately following application. Treat when insects first appear or when you notice lawn damage. For best results, leaf litter and debris should be removed from the foundation area. Branches and vegetation that touch the foundation should be cut back. Other potential pest harborage sites, such as deep mulch, should be raked away before application.

Overall Lawn Application

Pests	Use Rate	Use Instructions
Ants (Argentine, Southern, Field, Allegheny Mound, Florida Carpenter, Black Carpenter, Cornfield, Honey, Pavement, Red Imported Fire Ant, Nuisance) Armyworms (Fall & Yellow-striped) Centipedes European Crane Flies Crickets Cutworms Earwigs Fleas Lawn moths (sod webworms) Grasshoppers Japanese beetle (adults) Pillbugs Spittle bugs Mites Green June Beetles Leafhoppers Millipedes Springtails and Ticks (including Ticks which may transmit Lyme Disease)	Apply at the rate of 2 lb to 3 lb per 1,000 sq. ft. (an area 20' x 50') or 3.2 oz. to 4.8 oz. per 100 sq. ft. (10' x 10').	Apply with a fertilizer spreader (see spreader setting chart). Apply when grass is dry and water the lawn lightly immediately following application. Treat when insects first appear or when you notice lawn damage.

<p>Billbugs Grubs (Japanese beetle grubs, Green June Beetle Grubs) <i>Ataenus spretulus</i> (dung beetles) Hyperodes Weevils Mole Crickets Chinch bugs</p>	<p>Apply 3 lb. per 1,000 sq. ft. of lawn.</p>	<p>Water grass completely immediately following application.</p> <p>Treat when insects first appear or when you notice lawn damage.</p> <p>For hyperodes weevils, apply in April-May.</p> <p>For billbug control, apply in May when activity is first observed or when chewed or brown grass indicates damage from these insects.</p> <p>For grub control, except on dung beetle, treat any time between late July and early October.</p> <p>For dung beetle grub control, apply in early spring or early summer (March-July) when damage is first noticed. Immediately following application, completely water grass with ¼ to ½ inch of water. If a second generation occurs, treat again in late summer or early fall.</p> <p>For chinch bug control apply at higher rates during the summer when both nymphs and adults may be present.</p> <p>For mole crickets apply in the early spring for control of adults when actively invading grass areas. If the soil is dry, water before application to bring mole crickets closer to the surface and in contact with the granules. Apply late in the day and water in following application. For control of early nymphal stages, apply at the time of peak egg hatch. As the nymphs mature, higher rates may be necessary.</p>
--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Spot Treatment

Pests	Use Rate	Use Instructions
<p>Ants (Argentine, Southern, Field, Allegheny Mound, Florida Carpenter, Black Carpenter, Cornfield, Honey, Pavement, Red Imported Fire Ant, Nuisance)</p>	<p>Mound Treatment: Apply 1-2 teaspoons over and around each ant hill.</p>	<p>Water lightly immediately following application.</p> <p>NOTE: Use both broadcast and spot treatment for difficult to large ant infestations. If soil is dry, water lightly before application to bring the ants closer to the surface for better control.</p>
<p>Fire Ants</p>	<p>Mound Treatment: Apply ½ cup over, and 2 ft. around, each fire ant mound.</p>	<p>Immediately following application, water in completely. Apply a minimum of 1 gallon of water to each mound area. To prevent the disturbance of ants, water gently. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness. NOTE: For best results, apply in cool weather, 65°-80° F, or in early morning or late evening hours. Treat new mounds as they appear. Fast acting. Works within 24 hours.</p>

Spreader Settings for Lawn Application

80412

[INSERT SPREADER SETTING CHART]

NOTE: The spreader settings stated are approximate. Precisely weigh a few pounds into your spreader and apply to a measured area. To assure correct rate of application, adjust spreader settings if needed.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry area which is inaccessible to children and pets. Reclose container tightly after each use.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Completely empty container into application equipment. Then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Spills: If the container is leaking or material is spilled, carefully sweep up spilled product and wash exposed area. In spill or leak incidents, keep unauthorized people away.

WARRANTY STATEMENT

CONTROL SOLUTIONS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated in the Directions for Use when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or CONTROL SOLUTIONS, INC., and Buyer and User assume the risk of any such use.

[Note: Buyer assumes all responsibilities for safety and use not in strict accordance with directions.]

Control Solutions, Inc.
5903 Genoa-Red Bluff
Pasadena, TX 77507-1041

EPA Reg. No. 53883-195

EPA Est. No. xxxxxx-xx-xxx

90A12

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR LAMBDA LP BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using LAMBDA LP as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

NOTE: This supplemental label expires on June 15, 2011 and must not be used or distributed after this date.

LAMBDA LP

Lawn & Perimeter Insect Control Granules

EPA Reg. No. 53883-195

**KEEP OUT OF REACH OF CHILDREN
CAUTION/PRECAUCION**

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

JAN - 4 2010

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
53883-195**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

In case of an emergency with this product contact SafetyCall® International at (866) 897-8050.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of pesticide application.

Keep children and pets off treated area until product has been watered in and lawn is dry.

Restrictions:

- Apply this product directly to the lawn or garden area. Water treated area as directed on this label. Do not water to the point of run off.
- Do not make applications during rain.
- Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.
- Do not apply this product to edible growing crops or stored raw agricultural commodities used for food or feed.
- Do not allow applications to contact water inhabited by fish, such as in ornamental fish ponds that are located around structures being treated.
- Do not apply this product by aerial application.
- Use of this pesticide adjacent to water may affect aquatic organisms. To protect these organisms, do not apply this pesticide within 25 ft. of lakes, reservoirs, rivers, permanent streams, marches or natural ponds, estuaries, and commercial fish farm ponds.
- In New York State, this product may not be applied within 100 feet of a coastal marsh, or stream that drains directly into a coastal marsh.
- Observe all restrictions and precautions which appear on the labels of these products.
- Do not apply more than 0.36 lbs. of the a.i. (800 pounds of LAMBDA LP) per acre per year.

IMPORTANT: Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in outdoor ornamental gardens or parks, or on lawns or grounds.

GENERAL INFORMATION

LAWN AND PERIMETER PEST CONTROL

Lambda LP insecticide may be used for applications to outdoor areas where turf and ornamental ground covers are grown such as residential landscaped areas and non residential landscapes around institutional, public, commercial and institutional buildings, parks, recreational areas, and athletic fields. Broadcast Lambda LP Insecticide with suitable application equipment to ensure uniform coverage over the treatment area. The application rates listed in the **Lawn and Perimeter Use Rate** table will provide excellent control of the respective pests under typical conditions.

NOTE:

- Use higher rate for longer length of control.
- Watering (irrigation or rainfall) following application will dissolve the granules and release the insecticide.
- Make no more than 6 general applications per year. More frequent treatments should be limited to spot treatments.
- Uniform coverage is necessary to provide the most effective level of control.
- Consult your state university or local Cooperative Extension Service office for specific pest control application timing in your area.
- Application to turf should be started prior to the establishment of high insect pest populations and significant turf damage. Make reapplications as necessary to keep pest populations under control, using higher rates as pest pressure increases.

Recommended Application Rates for Lawn and Perimeter Use

	PEST	A.I. lb per acre	Lambda LP Rate Per 1000 ft2	Lambda LP Rate Per Acre
Band Treatment Around Foundation	Ants Crickets Firebrats Mites Sowbugs	Beetles Earwigs Millipedes Silverfish	0.04-0.06	2 - 3 lbs. 88 - 131 lbs.

Overall Lawn Application	Ants Armyworms Centipedes Crickets Cutworms Earwigs Grasshoppers Green June Beetles Japanese Beetle (Adult) Leafhoppers Millipedes Mites Pillbugs Sod webworms Spider Mites (Twospotted, Spruce) Spittlebugs Springtails and Ticks (including Ticks which may transmit Lyme Disease)	0.04-0.06	2 - 3 lbs.	88 – 131 lbs.
Overall Lawn Application	Billbugs Chinch Bugs Dung beetles* Green June beetle grubs* Hyperodes weevils Japanese beetle grubs* Mole Crickets	0.06-0.078	3 - 4 lbs	131 - 176 lbs.
Spot Treatment	Ants		Apply 1-2 teaspoons around each ant hill	
	Fire Ants		Mound Treatment: Apply ½ cup over, and 2 ft. around each fire ant mound	

*Suppression only.

SPECIFIC USE INSTRUCTIONS

Treatment Around Foundations

Pests Controlled: Ants (Argentine, Southern, Field, Allegheny Mound, Florida Carpenter, Black Carpenter, Cornfield, Honey, Pavement, Red Imported Fire Ant, Nuisance), Fleas, Roaches (German and American), Palmetto bugs, Waterbugs, Millipedes, Mites, Silverfish, Sowbugs (Pillbugs), Crickets, Earwigs

Application Rate: Apply at the rate of 2 lb to 3 lb per 1,000 sq. ft. (an area 5' x 200') oz. or 3.2 oz. to 4.8 oz. per 100 sq. ft. (10' x 10').

Application Instructions: For best results, leaf litter and debris should be removed from the foundation area. Branches and vegetation that touch the foundation should be cut back. Other potential pest harborage sites, such as deep mulch, should be raked away before application.

Overall Lawn Application

Chinch Bug and Mole Cricket Control -

For **chinch bug control**, apply at higher rates during the summer when both nymphs and adults may be present.

For **mole crickets** apply in the early spring for control of adults when actively invading grass areas. If the soil is dry, water before application to bring mole crickets closer to the surface and in contact with the granules. Apply late in the day and water in following application. For control of early nymphal stages, apply at the time of peak egg hatch. As the nymphs mature, higher rates may be necessary.

Spot Treatment

Pests Controlled: Ants (Argentine, Southern, Field, Allegheny Mound, Florida Carpenter, Black Carpenter, Cornfield, Honey, Pavement; Red Imported Fire Ant, Nuisance)

Mound Treatment: Apply 1-2 teaspoons over and around each ant hill.

NOTE: Use both broadcast and spot treatment for difficult to large ant infestations. If soil is dry, water lightly before application to bring the ants closer to the surface for better control.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry area which is inaccessible to children and pets. Reclose container tightly after each use.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Completely empty container into application equipment. Then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Spills: If the container is leaking or material is spilled, carefully sweep up spilled product and wash exposed area. In spill or leak incidents, keep unauthorized people away.

Read the "WARRANTY STATEMENT" [NOTE] on the label for Lambda LP before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Control Solutions, Inc.
5903 Genoa-Red Bluff
Pasadena, TX 77507-1041