

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

August 15, 2024

SENT BY EMAIL

Lisa Adamson
ladamson@controlsolutionsinc.com
CONTROL SOLUTIONS, INC.

Subject: Labeling Notification per Pesticide Registration Notice (PRN) 98-10 - Revise label with

optional graphics and a claim

Product Name: LAMBDA 0.5% CONCENTRATE

Admin Number: 53883-197 EPA Receipt Date: 07/22/2024 Action Case Number: 00622778

Dear Lisa Adamson:

The U.S. Environmental Protection Agency is in receipt of your application for notification under Pesticide Registration Notice 98-10 for the above referenced product. The EPA has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The labeling submitted with this application has been stamped "Notification" and will be placed in our records.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains statements or claims substantially differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

If you have questions, please contact Elizabeth Andrews via email at andrews.elizabeth@epa.gov.

Sincerely,

Elizabeth Andrews, Risk Manager

IVB1, RD

Office of Pesticide Programs

Sub-label A: Concentrate Packaging

LAMBDA-CYHALOTHRIN GROUP 3A INSECTICIDE

LAMBDA 0.5% CONCENTRATE

[alt. name Martin's Cyonara™ Home & Garden Insect Control] [alt. name Martin's Cyonara™ Home & Garden Insect Control Ready To Spray]

For outdoor use around the home only.

Active Ingredient: Lambda-cyhalothrin	0.5% 99.5%
Total:	100.0%
Contains 0.04 lbs. Lambda-cyhalothrin per gallon.	
KEEP OUT OF REACH OF CHILDREN CAUTION	

[See attached [back panel] booklet for additional precautionary statements. **OR** Open for Precautionary Statements, First Aid and Complete Directions for Use.]

EPA Reg. No. 53883-197

EPA Est. No.

NOTIFICATION

53883-197

Net Contents ____

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

8/15/2024

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

	First Aid
Have the product c	ontainer or label with you when calling a poison control center or doctor, or
going for treatment	· ·
If swallowed:	Call a Poison Control Center or doctor immediately for treatment advice.
	 Have a person sip a glass of water if able to swallow.
	 Do not induce vomiting unless told to do so by a Poison Control Center or doctor.
	 Do not give anything by mouth to an unconscious person.
If in eyes:	Hold eye open and rinse slowly and gently with water for 15-20 minutes.
	 Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
	 Call a poison control center or doctor for treatment advice.
You may also conta information.	act SafetyCall® International (866) 897-8050 for emergency medical treatment

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. Do not contaminate water when disposing of equipment washwaters. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. This product is highly toxic to bees exposed to direct treatement or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatement area.

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

RESTRICTIONS

- Do not apply as a broadcast application using a backpack sprayer on landscaping and turf.
- Do not apply to residential lawns and turf in residential settings (e.g. homes, parks, schools, athletic fields, or any other area frequented by the general public.
- Treat surfaces to ensure thorough coverage but avoid runoff.
- To treat insects harbored in voids and cracks-and-crevices, applications must be made in such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants.
- Do not allow children or pets into the treated area until dry.
- Do not apply when the wind speed is greater than 10mph.
- Do not treat pets with this product.
- Do not apply this product in a way that will contact adults, children, or pets, either directly or through drift.
- Spot treatments must not exceed two square feet in size (for example, 2 ft. by 1 ft. or 4 ft. by 0.5 ft.).
- All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:
 - 1. Application to pervious surfaces such as soil, lawn, turf, and other vegetation;
 - 2. Perimeter band treatments of 7 feet wide or less from the base of a man-made structure to pervious surfaces (e.g., soil, mulch, or lawn);
 - 3. Applications to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure;
 - 4. Applications to vertical surfaces (such as the side of a man-made structure) directly above impervious surfaces (e.g., driveways, sidewalks, etc.), up to 2 feet above ground level;
 - 5. Applications to vertical surfaces directly above pervious surfaces, such as soil, lawn, turf, mulch or other vegetation) only if the pervious surface does not drain into ditches, storm drains, gutters, or surface waters.
- For soil or foliar applications, do not apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.
- Do not apply the product into fish pools, ponds, streams, or lakes. Do not apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.
- Do not allow the product to enter any drain during or after application.
- Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment.
- Do not apply or irrigate to the point of runoff.

 Do not make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours). Rainfall within 24 hours after application may cause unintended runoff of pesticide application.

In New York State, this product may not be applied to lawns within 100 ft. of a coastal marsh, or stream that drains directly into a coastal marsh.

READ THE ENTIRE LABEL BEFORE USE.

PRODUCT INFORMATION

Lambda 0.5% Concentrate kills ants, fleas, chinch bugs, centipedes, mosquitoes, fire ants, spiders and other listed insects.

MIXING AND APPLICATION

- 1. Apply as directed.
- 2. In a CLEAN sprayer, measure and mix the amount of product and water as stated in the tables below.
- 3. When handling pesticide products, wear chemical resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, and viton ≥ 14 mils. Do not use kitchen utensils such as measuring cups and spoons for food purposes after they have been used with pesticides.
- 4. Spray as directed to achieve complete coverage.
- 5. Flush sprayer with clean water following each use.

Treat when insects begin to appear and repeat treatment to maintain control only as directed. Apply as directed, keeping in mind that complete coverage is crucial. Make certain to follow the important use limitations such as, the amount of spray to apply and the PHI (Pre-Harvest Interval) for the particular vegetable you are treating.

MEASUREMENT CONVERSIONS

1 tablespoon = 3 teaspoons

1 fl. oz. = 6 teaspoons or 2 tablespoons

6 fl. oz. = $\frac{3}{4}$ cup

1 ½ tablespoons in 3 gallons of water = 1 ½ teaspoons in 1 gallon of water

2 tablespoons in 3 gallons of water = 2 teaspoons in 1 gallon of water

NOTE: This product is non-staining to most home siding depending on age and cleanliness. Prior to applying this product to areas that may contact home siding (vinyl siding in particular), test in a hidden area and recheck in a few hours. If any staining is observed, do not use this product.

PESTS OUTSIDE THE HOME

Lambda 0.5% Concentrate is for use on ornamental plants next to building foundations and to perimeters and surfaces of buildings.

BARRIER APPLICATION INSTRUCTIONS: Treat when insects begin to appear and repeat treatment to maintain control as directed. Apply a continuous band of insecticide around the foundation of the building up to a maximum height of 2 ft to significantly decrease possible access by crawling pests. For optimum results, remove any leaf litter and debris around the foundation. Cut back any branches or vegetation that come into contact with the foundation. Rake back deep mulch and remove any other material that may serve as harborage sites for pests.

Outdoor Surfaces Use

Pests	Rate	Comments
Ants*(except for fire,	Mix 0.5 fl. oz.	Apply a 3 ft. band around the building and
pharaoh, harvester	(1 tablespoon) in 1	upwards along the foundation to 2 feet.
and carpenter ants)	gallon of water to treat	Apply as a coarse spray to thoroughly and
Centipedes	250 sq. ft.	uniformly wet the foundation and/or band area
Cockroaches		so that the insecticide will reach the soil or
Crickets		thatch level where pests may be active.
Fleas		
Kudzu bugs		1 gallon will treat 250 sq. ft. of surface area.
Millipedes		
Mosquitoes		Repeat treatment every 8 weeks (2 months)
Palmetto bugs		at the 2 fl. oz. per 1,000 sq. ft. rate as needed.
Scorpions		
Sowbugs		
Pillbugs		
Spiders		
Waterbugs		

^{*}Argentine, Southern, Field, Allegheny mound, Florida carpenter, Cornfield, Honey, Pavement, Nuisance, Odorous, Pyramid, Black turf, White-footed, Crazy, Little Black, Ghost, Thief, Acrobat, Citronella, Big-headed, Lasius.

Restrictions:

- Do not apply more than 2 fl. oz (0.00063 lbs. Lambda-cyhalothrin) per 1,000 sq. ft. per application.
- Do not make more than 14 applications per year per location.

FIRE ANT MOUNDS

Mix 1.0 fl. oz. (2 tablespoons) in 1 gallon of water and throughly wet each mound and a 4 ft. diameter surrounding the mound. To achieve optimum results, treat in cool weather (65°-80°F), or in the early morning or late evening hours. Apply to new mounds as they appear and do not disrupt mounds before treatment. Kills the entire mound and the queen in 24 hours.

[MOSQUITO CONTROL:

Application Instructions: Measure the area to be sprayed and follow the mixing instructions in the tables below. Treat when insects begin to appear and repeat treatment to maintain control as directed. Apply as directed, keeping in mind that complete coverage is crucial.

Outdoor Surfaces: ornamental plants next to building foundations, buildings, screens. Mix 0.5 oz (1 tablespoon) in 1 gallon of water. Apply up to a 7 ft. band around the building and upwards along the foundation to 2 feet . Thoroughly and uniformly wet the foundation and/or band area so that the insecticide will reach the soil or thatch level where pests may be active. One gallon treats 250 square feet. Repeat treatments every 8 weeks (2 months).]

ORNAMENTAL TREES, SHRUBS AND FLOWERS

APPLICATION INSTRUCTIONS: Apply thoroughly as a foliar cover spray to both sides of plant leaves and all plant surfaces where insects are present. For optimum results, apply early prior to high insect populations. Repeat applications as directed to maintain control.

NOTE:

- For **Scale** control, treat the whole plant, including the stem, trunks, and twigs.
- For **Bagworm** control, apply when bagworm larvae start to hatch.

Use Sites	Rate	Insects Controlled
For use on ornamental trees, shrubs and flowers: Ageratum, Ash, Aster, Arborvitae, Arizona Cypress, Azaleas, Begonia, Birch, Boxwood, Box Elder, Camellias, Cherry (ornamental), Non-bearing Citrus, Chrysanthemums, Carnations, Coleus, Common Ninebark, Conifers, Douglas Fir, Elm, English Ivy, Euonymus, Exacum, Weeping Fig, Fir, Gladioli, Gold Bells, Hawthorn, Holly, Honeysuckle, Hypoestes, Ivy, Juniper, Lilac, Locust, Maple, Marigold, Mimosa, Mock Orange, Nannyberry, Oak, Orchid, Palm, Pansy, Pea Shrub, Petunia, Philodendron, Pine, Ornamental Plum, Poinsettia, Poplar, Tulip Poplar, Rhododendron, Roses, Snapdragon, Snowberry,	Mix 0.25 fl. oz. (1/2 tablespoon) in 1 gallon of water to treat 250 sq. ft.	Ants*, Armyworms, Aphids*, Azalea caterpillar, Bagworms, Black vine weevil (Adult), California oakworm, Boxelder bugs, Budworms, Cankerworms, Cicadas, Cockroaches, Crickets, Cutworms, Eastern tent caterpillar, Elm leaf beetles, European sawfly, Fall webworm, Flea beetles, Forest tent caterpillar, Gypsy moth larvae, Japanese beetles (Adults), June beetles (Adults), Lace bugs, Ladybeetles, Leaf feeding caterpillars, Leafhoppers, Leafminers (Adults), Leaf rollers, Leaf skeletonizers, Kudzu bugs, Midges, Moths, Oleander moth larvae, Pillbug, Pine sawfly, Pine shoot beetles, Pinetip moths, Root weevil, Sawfly, Scale insects (crawlers), Spiders, Spittlebugs, Striped beetles, Striped oakworm, Thrips, Tip moth, Tussock moth larvae
Spruce, Taxus, Willow, Zinnia.	Mix 0.5 fl. oz. (1 tablespoon) in 1 gallon of water to treat 250 sq. ft	Broadmites, Brown softscale, California redscale (crawler), Clover mites, Mealybugs, Pineneedle scale (crawler), Spider mites, Whiteflies

^{*} Allegheny mound, Argentine, Black carpenter, Cornfield, Field, Florida carpenter, Harvester, Honey, Pavement, Southern, Sugar, Nuisance, Odorous, Pyramid, Black turf, White-footed, Crazy, Little Black, Ghost, Thief, Acrobat, Citronella, Big-headed, Lasius.

- Do not apply more than 4 fl. oz (0.00063 lbs. Lambda-cyhalothrin) per 1,000 sq. ft. per application.
- Do not make more than 14 applications per year per location at 2 fl. oz per 1,000 sq. ft. rate.
- Do not make more than 7 applications per year per location at 4 fl. oz per 1,000 sq. ft. rate.

VEGETABLES

APPLICATION INSTRUCTIONS: Apply when insects first appear or when damage occurs. For complete coverage, spray all plant surfaces that are infested with pests. Follow the important use limitations such as, the amount of spray to apply and the PHI (Pre-Harvest Interval) for the particular vegetable you are treating.

RATE: Measure the area to be sprayed. Mix $0.5 \, \text{fl.}$ oz. (1 tablespoon) in 1 gallon of water per 300 sq. ft.

Vegetables	Insects Controlled	PHI*
Broccoli Brussels Sprouts Cabbage Cavalo Broccoli Cauliflower Chinese Broccoli Chinese Cabbage Chinese Mustard Kohlrabi	Alfalfa Looper, Aphid, Armyworm, Beet Armyworm, Cabbage Looper, Cabbage Webworm, Corn Earworm, Cutworm, Diamondback Moth, Fall Armyworm, Flea Beetle, Grasshopper, Imported Cabbageworm, Japanese Beetle (Adult), Leafhopper, Lygus Bug, Meadow Spittlebug, Plant Bug, Southern Cabbageworm, Spider Mite, Stink Bug, Thrips, Yellow-striped Armyworm, Whitefly	1
Corn (sweet and pop)	Armyworm, Bean Leaf Beetle, Beet Armyworm, Cereal Leaf Beetle, Chinch Bug, Corn Earworm, Corn Leaf Aphid, Cutworm, English Grain Aphid, European Corn Borer, Grasshopper, Green Bug, Green Cloverworm, Hop Vine Borer, Japanese Beetle (Adult), Meadow Spittlebug, Mexican Corn Rootworm Beetle (Adult), Northern Corn Rootworm Beetle (Adult), Sap Beetle (Adult), Southern Corn Rootworm Beetle (Adult), Southwestern Corn Borer, Stalk Borer, Stink Bug, Tobacco Budworm, Western Bean Cutworm, Western Corn Rootworm Beetle (Adult)	21
Corn (sweet only)	Aphid, Aster Leafhopper, Corn Silkfly (Adult), Southern Armyworm, Spider Mite, Tarnished Plant Bug, Western Bean Cutworm	21
Lettuce (head and leaf)	Aphid, Alfalfa Looper, Armyworm, Beet Armyworm, Cabbage Looper, Corn Earworm, Cutworm, Diamondback Moth, European Corn Borer, Fall Armyworm, Flea Beetle, Grasshopper, Green Cloverworm, Imported Cabbageworm, Japanese Beetle (Adult), Leafhopper, Meadow Spittlebug, Plant Bug, Saltmarsh Caterpillar, Southern Armyworm, Spider Mite, Stink Bug, Tobacco Budworm, Vegetable Weevil (Adult), Whitefly	1
Onion (bulb) and Garlic	Aphid, Armyworm, Cutworm, Flower Thrips, Leafminer (Adult), Onion Maggot (Adult), Onion Thrips, Plant Bug, Seedcorn Maggot (Adult), Stink Bug, Tobacco Thrips, Western Flower Thrips	14
Tomato and Tomatillo	Aphid, Beet Armyworm, Cabbage Looper, Colorado Potato Beetle, Cutworm spp., European Corn Borer, Fall Armyworm, Flea Beetle, Grasshopper, Hornworm, Leafhopper, Leafminer, Meadow Spittlebug, Plant Bug, Southern Armyworm, Stink Bug, Tobacco Budworm, Tomato Fruitworm, Tomato Pinworm, Yellow-striped Armyworm, Whitefly	5

- Do not apply more than 1.67 fl. oz (0.00053 lbs. Lambda-cyhalothrin) per 1,000 sq. ft. per application.
- Do not make more than 9 applications per year per location.
- Do not make more than 5 applications per year per location for corn.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in cool, dry area away from children and domestic animals. **CONTAINER HANDLING:** If empty: Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. If recycling is not available: then dispose of container in a sanitary landfill. If partly filled: Call your local solid waste agency for disposal instructions. Unless otherwise instructed, place in trash. Never pour unused product down the drain or on the ground.

LIMITATION OF WARRANTY AND LIABILITY

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Control Solutions, Inc. or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Control Solutions, Inc. and Seller harmless for any claims relating to such factors.

Control Solutions, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Control Solutions, Inc., and Buyer and User assume the risk of any such use. To the extent permitted by applicable law, CONTROL SOLUTIONS, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

IT IS CONTROL SOLUTIONS INC., AND SELLERS INTENTION THAT in no event shall Control Solutions or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF CONTROL SOLUTIONS, INC., AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF CONTROL SOLUTIONS, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.

Control Solutions, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Control Solutions, Inc.

Control Solutions, Inc. 5903 Genoa-Red Bluff Road Pasadena, TX 77507-1041 LAMBDA-CYHALOTHRIN GROUP 3A INSECTICIDE

LAMBDA 0.5% CONCENTRATE

[alt. name Martin's Cyonara™ Home & Garden Insect Control] [alt. name Martin's Cyonara™ Home & Garden Insect Control Ready To Spray]

For outdoor use around the home only.

Active Ingredient:	
Lambda-cyhalothrin	0.5%
Other Ingredients:	99.5%
Total:	100.0%
Contains 0 04 lbs. Lambda-cyhalothrin per gallon	

KEEP OUT OF REACH OF CHILDREN CAUTION

[See attached [back panel] booklet for additional precautionary statements. **OR** Open for Precautionary Statements, First Aid and Complete Directions for Use.]

EPA Reg. No. 53883-197

EPA Est. No.

Net Contents ____

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

	First Aid
Have the product of going for treatmen	container or label with you when calling a poison control center or doctor, or t.
If swallowed:	 Call a Poison Control Center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor.
	 Do not give anything by mouth to an unconscious person.
If in eyes:	 Hold eye open and rinse slowly and gently with water for 15-20 minutes.
	 Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
	 Call a poison control center or doctor for treatment advice.
You may also cont information.	act SafetyCall® International (866) 897-8050 for emergency medical treatment

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. Do not contaminate water when disposing of equipment washwaters. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. This product is highly toxic to bees exposed to direct treatement or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatement area.

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

PRODUCT INFORMATION

This product provides control against listed backyard nuisance pests: mosquitoes, ants, cockroaches, fleas, [and other listed pests]. Simply connect to hose and spray for 8 weeks protection.

RESTRICTIONS:

- Do not apply as a broadcast application using a backpack sprayer on landscaping and turf.
- Do not apply to residential lawns and turf in residential settings (e.g. homes, parks, schools, athletic fields, or any other area frequented by the general public.
- Treat surfaces to ensure thorough coverage but avoid runoff.
- To treat insects harbored in voids and cracks-and-crevices, applications must be made in such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants.
- Do not allow children or pets into the treated area until dry.
- Do not apply when the wind speed is greater than 10mph.Do not apply to residential lawns and turf in residential settings (e.g. homes, parks, schools, athletic fields, or any other area frequented by the general public.
- Do not treat pets with this product.
- Do not apply this product in a way that will contact adults, children, or pets, either directly or through drift.
- Spot treatments must not exceed two square feet in size (for example, 2 ft. by 1 ft. or 4 ft. by 0.5 ft.).
- All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:
 - 1. Application to pervious surfaces such as soil, lawn, turf, and other vegetation;
 - 2. Perimeter band treatments of 7 feet wide or less from the base of a man-made structure to pervious surfaces (e.g., soil, mulch, or lawn);
 - 3. Applications to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure;
 - 4. Applications to vertical surfaces (such as the side of a man-made structure) directly above impervious surfaces (e.g., driveways, sidewalks, etc.), up to 2 feet above ground level;
 - 5. Applications to vertical surfaces directly above pervious surfaces, such as soil, lawn, turf, mulch or other vegetation) only if the pervious surface does not drain into ditches, storm drains, gutters, or surface waters.

- For soil or foliar applications, do not apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.
- Do not apply the product into fish pools, ponds, streams, or lakes. Do not apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.
- Do not allow the product to enter any drain during or after application.
- Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment.
- Do not apply or irrigate to the point of runoff.
- Do not make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours). Rainfall within 24 hours after application may cause unintended runoff of pesticide application.

In New York State, this product may not be applied to lawns within 100 ft. of a coastal marsh, or stream that drains directly into a coastal marsh.

APPLICATION INSTRUCTIONS

Determine the area to be treated. For trees, shrubs and flowers, apply as a thorough cover spray. Treat when insects begin to appear and repeat treatment to maintain control only as directed. Apply as directed, keeping in mind that complete coverage is crucial. Follow the important use limitations such as, the amount of spray to apply and the PHI (Pre-Harvest Interval) for the particular vegetable you are treating.

- One quart (32 oz.) (0.01 lbs. Lambda-cyhalothrin) to treat up to 8,000-16,000 sq. ft.
- Do not make more than 7 applications per year per location at the one quart (32 oz.) per 8,000 sq. ft. rate
- Do not make more than 14 applications per year per location at the one quart (32 oz.) per 16,000 sq. ft. rate

Ready Spray Nozzle Instructions

- 1. Shake container well before use.
- 2. Connect the Ready to Spray nozzle to a hose, making sure that the dial on the nozzle is in the "OFF" position with the (green) safety tab in the valve notch.
- 3. Turn on water. Begin treatment on the furthest area and work back toward the faucet so that you do not come in contact with treated areas.
- 4. To START spraying, direct nozzle toward treatment area and bend the (green) safety tab back (located at the right of the yellow dial) and hold while turning the yellow dial counter clockwise until it stops. Water will automatically mix with the product.
- 5. Spray desired area until wet to control insect pests. Evenly spray the area by walking at a steady pace and sweep evenly, slightly overlapping treated areas.
- 6. To STOP spraying, turn the dial counter clockwise **quickly** until it stops and the safety tab engages into the notch on the valve. Turn the water off. To ease pressure before removing nozzle from the hose, bend the safety tab back and dial "ON" until water stops spraying.
- 7. Make sure the dial is in the "OFF" position with (green) safety tab in the valve notch to store unused product. Keep in a cool area away from heat, sunlight or open flame.

SPECIFIC USE DIRECTIONS

How to Use the	• [Connect] [Hose]
Ready to Spray	• [1.]Shake Well.
	• [2.] Make sure control lever is in the OFF position. Attach to hose.
	• [3.] Turn on water. Extend hose to furthest place around perimeter or in

garden, so you can work backwards and limit exposure to spray. [Spray] [Apply mix] • [1.] Aim towards direction of application and away from you. • [2.] Turn control lever to ON to begin spraying. • [3.] Spray evenly and as directed for specified type of treatment area.] [Garden] [trees][shrubs][flowers][perimeter] • [Finish] [Stop] [Store] • [1.] When finished turn control lever to OFF. • [2.] Turn faucet OFF. Discharge residual water pressure by turning control lever ON and OFF again. • [3.] Disconnect Sprayer and Store / Discard as instructed. APPLICATION • Apply where insects are expected to appear or when insects or damage are INSTRUCTIONS seen. Repeat as necessary according to site-specific directions listed below. · Apply in calm weather. Begin spraying at the farthest point of the treatment area and work backward to avoid contact with wet treated surfaces. • Spray thoroughly to wet plant surfaces up to the point of runoff. • Sprays must be dry in treated areas before people or pets re-enter. • Do not apply to trees taller than 10 feet. Consider hiring a licensed professional. Follow the important use limitations such as, the amount of spray to apply and the PHI (Pre-Harvest Interval) for the particular vegetable you are Keep in a cool area away from heat, sunlight or open flame.

(NOTE TO REVIEWER: Table above is optional and if used, will replace the text directly above. Final label will only contain either the box or the text above the box.}

NOTE: This product is non-staining to most home siding depending on age and cleanliness. Prior to applying this product to areas that may contact home siding (vinyl siding in particular), test in a hidden area and recheck in a few hours. If any staining is observed, do not use this product.

PESTS OUTSIDE THE HOME

Lambda 0.5% Concentrate is for use on ornamental plants next to building foundations, to surfaces of buildings and screens.

BARRIER APPLICATION INSTRUCTIONS: Treat when insects begin to appear and repeat treatment to maintain control as directed. Apply a continuous band of insecticide around the foundation of the building and to a maximum height of 2 ft. to significantly decrease possible access by crawling pests. For optimum results, remove any leaf litter and debris around the foundation. Cut back any branches or vegetation that come into contact with the foundation. Rake back deep mulch and remove any other material that may serve as harborage sites for pests.

Outdoor Surfaces Use

Pests	Comments
Ants*(excluding fire,	Apply a 3 ft. band around the building and upwards along the
pharaoh, harvester, and	foundation to 2 feet. Apply as a coarse spray to thoroughly and
carpenter ants)	uniformly wet the foundation and/or band area so that the insecticide
Centipedes	will reach the soil or thatch level where pests may be active.
Cockroaches	
Crickets	32 ounces (1 bottle) treats 16,000 sq. ft.

Fleas	
Millipedes	Repeat treatment every 8 weeks (2 months).
Mosquitoes	
Palmetto bugs	
Scorpions	
Sowbugs	
Pillbugs	
Spiders	
Waterbugs	

^{*}Argentine, Southern, Field, Allegheny mound, Florida carpenter, Conrfield, Honey, Pavement, Nuisance, odorous, pyramid, black turf, white-footed, Crazy, Little Black, Ghost, Thief, Acrobat, Citronella, Big-headed, Lasius.

Do not make more than 14 applications per year per location at the 1 quart (32 fl. oz.).
 per 16,000 sq. ft. rate.

FIRE ANT MOUNDS

Throughly wet each mound and a 4 ft. diameter surrounding the mound. To achieve optimum results, treat in cool weather (65°-80°F), or in the early morning or late evening hours. Apply to new mounds as they appear and do not disrupt mounds before treatment. Kills the entire mound and the queen in 24 hours.

• **Restrictions:**Do not make more than 14 applications per year per location at the one quart (32 oz.) per 16,000 sq. ft. rate

[MOSQUITO CONTROL:

Application Instructions: Measure the area to be sprayed. Treat when insects begin to appear and repeat treatment to maintain control as directed. Apply as directed, keeping in mind that complete coverage is crucial.

Restrictions:

• Do not make more than 14 applications per year per location at the one quart (32 oz.) per 16,000 sq. ft. rate

Outdoor Surfaces: ornamental plants next to building foundations, buildings, screens. Apply a 3 ft. band around the building and upwards along the foundation to 2 feet. Thoroughly and uniformly wet the foundation and/or band area so that the insecticide will reach the soil or thatch level where pests may be active. Repeat treatments every 8 weeks (2 months).]

ORNAMENTAL TREES, SHRUBS AND FLOWERS

APPLICATION INSTRUCTIONS: Apply thoroughly as a foliar cover spray to both sides of plant leaves and all plant surfaces where insects are present. Penetrate dense foliage. Wet leaves and branches to the point of runoff. For optimum results, apply early prior to high insect populations. Repeat applications as directed to maintain control.

NOTE:

- For **Scale** control, treat the whole plant, including the stem, trunks, and twigs.
- For **Bagworm** control, apply when bagworm larvae start to hatch.

Use Sites	Insects Controlled
For use on ornamental trees, shrubs and	Ants*(excluding fire, pharaoh, harvester and

flowers including:

Ageratum, Ash, Aster, Arborvitae, Arizona Cypress, Azaleas, Begonia, Birch, Boxwood, Box Elder, Camellias, Cherry (ornamental), Non-bearing Citrus, Chrysanthemums, Carnations, Coleus, Common Ninebark, Conifers, Douglas Fir, Elm, English Ivy, Euonymus, Exacum, Weeping Fig, Fir, Gladioli, Gold Bells, Hawthorn, Holly, Honeysuckle, Hypoestes, Ivy, Juniper, Lilac, Locust, Maple, Marigold, Mimosa, Mock Orange, Nannyberry, Oak, Orchid, Palm, Pansy, Pea Shrub, Petunia, Philodendron, Pine, Ornamental Plum, Poinsettia, Poplar, Tulip Poplar, Rhododendron, Roses, Snapdragon, Snowberry, Spruce, Taxus, Willow, Zinnia.

carpenter), Armyworms, Aphids, Azalea caterpillar, Bagworms, Black vine weevil (Adult), California oakworm, Boxelder bugs, Budworms, Cankerworms, Cicadas, Cockroaches, Crickets, Cutworms, Eastern tent caterpillar, Elm leaf beetles, European sawfly, Fall webworm, Flea beetles, Forest tent caterpillar, Gypsy moth larvae, Japanese beetles (Adults), June beetles (Adults), Lace bugs, Ladybeetles, Leaf feeding caterpillars, Leafhoppers, Leafminers (Adults), Leaf rollers, Leaf skeletonizers, Kudzu bugs, Midges, Moths, Oleander moth larvae, Pillbug, Pine sawfly, Pine shoot beetles, Pinetip moths, Root weevil, Sawfly, Scale insects (crawlers), Spiders, Spittlebugs, Striped beetles, Striped oakworm, Thrips, Tip moth, Tussock moth larvae Broadmites, Brown softscale, California redscale (crawler), Clover mites, Mealybugs, Pineneedle scale (crawler), Spider mites, Whiteflies

* Allegheny mound, Argentine, Black carpenter, Cornfield, Field, Florida carpenter, Harvester, Honey, Pavement, Southern, Sugar, Nuisance, Odorous, Pyramid, Black turf, White-footed, Crazy, Little Black, Ghost, Thief, Acrobat, Citronella, Big-headed, Lasius.

Restrictions:

- One quart (32 oz.) (0.01 lbs. Lambda-cyhalothrin) to treat up to 8,000-16,000 sq. ft.
- Do not make more than 7 applications per year per location at the one quart (32 oz.) per 8,000 sq. ft. rate
- Do not make more than 14 applications per year per location at the one quart (32 oz.) per 16,000 sq. ft. rate

VEGETABLES

APPLICATION INSTRUCTIONS: Apply when insects first appear or when damage occurs. For complete coverage, spray all plant surfaces that are infested with pests and penetrate dense foliage. Make certain to follow the important use limitations such as, the amount of spray to apply and the PHI (Pre-Harvest Interval) for the particular vegetable you are treating.

Vegetables	Insects Controlled	PHI*
Broccoli Brussels Sprouts Cabbage Cavalo Broccoli Cauliflower Chinese Broccoli Chinese Cabbage Chinese Mustard Kohlrabi	Alfalfa Looper, Aphid, Armyworm, Beet Armyworm, Cabbage Looper, Cabbage Webworm, Corn Earworm, Cutworm, Diamondback Moth, Fall Armyworm, Flea Beetle, Grasshopper, Imported Cabbageworm, Japanese Beetle (Adult), Leafhopper, Lygus Bug, Meadow Spittlebug, Plant Bug, Southern Cabbageworm, Spider Mite, Stink Bug, Thrips, Yellow-striped Armyworm, Whitefly	1

Corn (sweet and pop)	Armyworm, Bean Leaf Beetle, Beet Armyworm, Cereal Leaf Beetle, Chinch Bug, Corn Earworm, Corn Leaf Aphid, Cutworm, English Grain Aphid, European Corn Borer, Grasshopper, Green Bug, Green Cloverworm, Hop Vine Borer, Japanese Beetle (Adult), Meadow Spittlebug, Mexican Corn Rootworm Beetle (Adult), Northern Corn Rootworm Beetle (Adult), Sap Beetle (Adult), Southern Corn Rootworm Beetle (Adult), Southwestern Corn Borer, Stalk Borer, Stink Bug, Tobacco Budworm, Western Bean Cutworm, Western Corn Rootworm Beetle (Adult)	21
Corn (sweet only)	Aphid, Aster Leafhopper, Corn Silkfly (Adult), Southern Armyworm, Spider Mite, Tarnished Plant Bug, Western Bean Cutworm	21
Lettuce (head and leaf)	Aphid, Alfalfa Looper, Armyworm, Beet Armyworm, Cabbage Looper, Corn Earworm, Cutworm, Diamondback Moth, European Corn Borer, Fall Armyworm, Flea Beetle, Grasshopper, Green Cloverworm, Imported Cabbageworm, Japanese Beetle (Adult), Leafhopper, Meadow Spittlebug, Plant Bug, Saltmarsh Caterpillar, Southern Armyworm, Spider Mite, Stink Bug, Tobacco Budworm, Vegetable Weevil (Adult), Whitefly	1
Onion (bulb) and Garlic	Aphid, Armyworm, Cutworm, Flower Thrips, Leafminer (Adult), Onion Maggot (Adult), Onion Thrips, Plant Bug, Seedcorn Maggot (Adult), Stink Bug, Tobacco Thrips, Western Flower Thrips	14
Tomato and Tomatillo	Aphid, Beet Armyworm, Cabbage Looper, Colorado Potato Beetle, Cutworm spp., European Corn Borer, Fall Armyworm, Flea Beetle, Grasshopper, Hornworm, Leafhopper, Leafminer, Meadow Spittlebug, Plant Bug, Southern Armyworm, Stink Bug, Tobacco Budworm, Tomato Fruitworm, Tomato Pinworm, Yellow-striped Armyworm, Whitefly	5
* PHI = Pre-Harvest Interval. Number of days to wait between the last application and harvest.]		

- One quart (32 oz.) (0.01 lbs. Lambda-cyhalothrin) to treat up to 16,000 sq. ft.
- Do not make more than 9 applications per year per location for vegetable treatment
- Do not make more than 5 applications per year per location for corn

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in cool, dry area away from children and domestic animals. **CONTAINER HANDLING:** If empty: Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. If recycling is not available: then dispose of container in a sanitary landfill. If **partly filled:** Call your local solid waste agency for disposal instructions. Unless otherwise instructed, place in trash. Never pour unused product down the drain or on the ground.

LIMITATION OF WARRANTY AND LIABILITY

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended

consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Control Solutions, Inc. or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Control Solutions, Inc. and Seller harmless for any claims relating to such factors.

Control Solutions, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Control Solutions, Inc., and Buyer and User assume the risk of any such use. To the extent permitted by applicable law, CONTROL SOLUTIONS, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

IT IS CONTROL SOLUTIONS INC., AND SELLERS INTENTION THAT in no event shall Control Solutions or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF CONTROL SOLUTIONS, INC., AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF CONTROL SOLUTIONS, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.

Control Solutions, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Control Solutions, Inc.

Control Solutions, Inc. 5903 Genoa-Red Bluff Road Pasadena, TX 77507-1041

[Optional Marketing Statements- Applicable to Both Sub-Labels]

- [Protects Vegetables, Roses, Flowers, Trees, Shrubs.]
- [Controls Ants (excluding pharaoh, harvester and carpenter ants), Aphids, Chinch Bugs, Cutworms, Flea Beetles, Mites, & many more.]
- [Kills Garden Insects, including chinch bugs, fleas, fire ants, mosquitoes, aphids, Japanese beetles, whiteflies, ants (excluding pharaoh, harvester and carpenter ants), mites, and other listed insects.]
- [For perimeter insect control.]
- [For Barrier use.]
- [For Garden Insects.]
- [Kills and Repels Insects.]
- [Kills Pests Outside the Home for Up to 8 Weeks (2 months)* *excluding red imported fire ants and other public health pests]
- [Kills Pests on Outdoor Surfaces for Up to 8 Weeks (2 months)* * excluding red imported fire ants and other public health pests]
- [Kills listed bugs outside so they can't come inside]
- Kills Bugs* Outside Before They Come Inside *see label for complete pest list
- [Low odor (formula).]
- [Fast Acting Formula.]
- [Works (Kills) in 24 hours.]
- [Kills 500+ Pests on contact.]
- [Long-lasting insect protection.]
- [Non-staining.]
- [Kills Fire Ants in 24 hours.]
- [Kills the Fire Ant Mound.]
- [Kills the Queen and the Mound.]
- [Controls Fire Ants for up to 4 Weeks.]
- [Fast acting with up to 8 weeks Residual* *excluding public health pests]
- Keeps killing for up to [2 months][8 weeks]* *excluding public health pests
- [Controls Mosquitoes, Chinch Bugs, Fleas, Aphids, Fire Ants & Mites]
- [Ready To Spray](when sold in ready to spray packaging)
- Kills Home Invading Insects
- Kills garden [invading] pests[insects] [including] [ants (excluding pharaoh, harvester and carpenter ants),] [aphids] [armyworms] [beetles] [cutworms] [Japanese beetles] [chinch bugs] [caterpillars] [sod worms] [white flies] and other listed pests [insects]
- Insect Protection
- Not harmful to lawn grasses* *When used as directed
- Works two ways: (1) Kills (and) (2) Protects
- Kills and prevents over 500 listed [insects][pests]
- Penetrates [mulch][thatch]
- Kills above and below the surface for [up to] 2 months
- Kills [Surface]] and [Soil] Insects
- Use on gardens [Outdoor Surfaces][plants] [vegetables][flowers] [trees]
 [Shrubs][HomePerimeter] [shrubs][ornamentals] [ornamentals]
- [For Home Fruit & Vegetable Gardens, Ornamental & Flower Gardens, and [Perimeters][Around the Home]]
- Can be used as an outdoor barrier treatment
- [Also][Can be used][for use] on vegetable gardens
- Tough on [insert listed insects], gentle on [your] garden [plants] [shrubs] [vegetables] [fruits] [edibles] [flowers][trees][ornamentals] [home][brick][wood] [siding]
- Won't harm [your] garden [plants] [shrubs] [vegetables] [fruits] [edibles]
 [flowers][trees][ornamentals][home][brick] [wood][siding]
- Protects [edibles][fruits and vegetables] from insect damage

- Protects over 20 listed fruits and vegetables
- Easy to use
- Kills listed nuisance pests outdoors [before they can come inside]
- Kill [annoying] [damaging] [insects] [pests] and reclaim your garden [today!]
- Reclaim your[backyard][today]
- Don't let insects [bugs] feed on your[landscape][and][backyard] another day
- Pretreat your [backyard] [outdoor spaces] before [the BBQ][outdoor fun]
- Pretreat your [backyard] [outdoor spaces] against [insert listed pest]
- Protect your backyard from [nuissance][pests]
- Protect your container garden from [pests][insert listed pest]
- Protect your [roses][flowers] from [damaging][harmful] pests
- Not harmful to grasses
- · Won't harm plants or blooms
- Kills listed common backyard [insects][bugs]
- Kills [harmful] [damaging] [insert listed pest]
- Stops damaging insects
- Kills pests by contact
- Visible results
- Kills by contact
- Starts working immediately by contact*. *excluding public health pests
- Fast acting for your outdoor weekend [plus 2 months control] * *excluding public health pests
- 8 weekends of [pest][insert listed pest][protection][control] * *excluding public health pests
- Lasts {up to} [4] [6] [8] weeks [2] [months] * *excluding public health pests
- Up to [4] [6] [8] weeks [2] month control [protection]* *excluding public health pests
- Protects your [lawn][backyard] for up to [2] months! * *excluding public health pests
- One application protects up to [8 weeks][2 months] * *excluding public health pests
- One Application [2 month][8 weeks][control][protection] * *excluding public health pests
- Kills [insert listed pest from appendix 1]
- Controls [insert listed pest from appendix 1]
- Kills Ants (excluding pharaoh, harvester and carpenter), Roaches, Spiders and Other Home Invading Pests as listed
- Kills [insert listed pest from appendix 1] on contact
- Effective control against [insert listed pest from appendix 1]
- Proven [insert listed pest from appendix 1] killer
- Kills mosquitoes that may transmit West Nile Virus
- Kills mosquitoes that may transmit [the] Zika Virus
- Kills mosquitoes that may transmit the Chikungunya virus
- Kills mosquitoes that may transmit yellow fever
- Kills mosquitoes that may transmit dengue fever
- Kills mosquitoes that may transmit encephalitis
- Kills Fleas that may transmit Flea Allergy Dermatitis (FAD)/ Flea-Bite Anemia/ Tapeworms/ Haemobartonellosis [to cats/dogs]
- Kills Fleas that may transmit Bartonellosis [to cats/dogs/people]
- Kills Fleas that may transmit Mycoplasma to cats
- Kills mosquitoes that may transmit Heart worm [to cats/dogs]
- Kills mosquitoes that may transmit Subcutaneous dirofilariasis [to cats/dogs/people]
- [Satisfaction] [Guaranteed*] [kill*] *or your money back
- Guaranteed results* *or your money back
- Guaranteed to work* *or your money back
- Trusted Brand since [insert year]
- Established Brand since [insert year]
- Est. [insert year]

- Trusted brand for over XX years
- Formulated and packaged in the USA
- Protects your [/backyard/garden] from bugs/insects [so you [and your kids/pets/dog] can enjoy your/the [outdoors/event/party/barbeque/birthday party/playtime/home/yard]]
- Kills bugs/insects [which destroy roses, flowers, tree, shrubs, edibles, landscape, backyard] [and/or] [which] [annoy people/kids/pets.]
- People and pets may re-enter the treated area once the product has dried.
- People and pets may enter [treated area][backyard] after spray has dried.
- Kills [nuissance][annoying] backyard pests, as listed
- Protecting your [backyard] [outdoors] has never been easier
- Protect your family from [insert listed pests][listed nuissance pests]
- Bug Killer
- Insect Killer
- Ready to Spray
- Hate The Bugs. Love The Garden. Get [brand]!
- Got [insert listed pest here]? Get [brand]!
- Shake well before using
- Water based formula
- Bonus Size
- Combo Pack
- Wow (Pack)
- Value size
- Value/Twin pack
- Great value
- Kills [insert listed pest], [insert listed pest], [insert listed pest] and 500 other listed pests
- Kills over 450 listed pests
- Kills 450[plus][+] [insects][pests]
- Economical use to cover large areas
- Easily covers large areas
- Read the entire label before applying

[If you have a medical emergency, need consultation on diagnosis, therapy and analysis please call (800) 420-9347]

[If you have questions relating to product use or application, please call (866) 945-5033]

[If you have a product spill or transportation emergency call (800) 424-9300]

[For Consumer Refund: TechPac, LLC PO Box 95437 Palatine, IL 60095-0437]

[Optional Marketing Statements- Applicable to Sub-Label A]

- [Broad Spectrum (Insect Killer) (Insecticide) (Concentrate).]
- [Concentrated Insect Spray.]
- [Makes up to 128 gallons.]
- Concentrate
- [Includes] easy to use measuring cap
- Easy-measure cap

[Optional Marketing Statements- Applicable to Sub-Label B]

- No mixing or measuring
- Easy-to-use sprayer
- No [mixing][measuring] required
- Simply connect to hose and spray [for 8 weeks protection] *. *excluding public health

pests

- [Simply] attach to [garden] hose and [spray] [go]
 Automatically mixes as you spray
- [Connect and Spray (optional for hose-end sprayer).]


Optional Images and associated text for applicable RTS Sprayer packaging:


SPRAY STREAM **ROTATE IN OFF POSITION** MIX H2O OFF





Quick [Reference Table][Facts] {NOTE to Reviewer: Table below is optional}

Quick [Reference Table][Facts] (NOTE to Reviewer: Table below is optional)		
[Product Facts]	[Insert listed claim(s)]	
PRODUCT FACTS		
[Where to Use]	Fruit and Vegetable Gardens, Ornamental Plants and	
WHERE TO USE	Shrubs, Flowers and Ground Covers, Around Home Perimeter [Insert listed claim(s)]	
Crack and crevice use only		
[Fruits & Vegetables]	[Insert listed fruits & vegetables]	
FRUITS & VEGETABLES	[And other listed fruits and vegetables]	
Kills	[Insert listed pests]	
KILLS	[And other listed [pests] [insects] [bugs]] [Insert listed claim(s)]	
















Appendix 1. [Complete] Pest List:

PEST LIST: {NOTE to Reviewer: This pest list is optional}

- 1. Alfalfa Looper
- 2. American Plum Borer
- 3. Ants Acrobat
- 4. Ants Big-headed
- 5. Ants Black turf
- 6. Ants Citronella
- 7. Ants Crazy
- 8. Ants Field
- 9. Ants Ghost
- 10. Ants Lasius
- 11. Ants Lawn
- 12. Ants Little Black
- 13. Ants Thief
- 14. Ants White-footed
- 15. Ants Allegheny mound
- 16. Ants Argentine
- 17. Ants Cornfield
- 18. Ants Honey
- 19. Ants Nuisance
- 20. Ants Odorous
- 21. Ants Pavement
- 22. Ants Pyramid
- 23. Ants Southern
- 24. Angular-winged katydid
- 25. Aphid
- 26. Aphids Buckhorn
- 27. Aphids Giant Willow28. Aphids Greenbug
- 29. Aphids Green Apple
- 30. Aphids Pea
- 31. Aphids Potato
- 32. Aphids Root
- 33. Aphids Rose
- 34. Aphids Walnut
- 35. Aphids Wooly Alder
- 36. Aphids Wooly Apple
- 37. Aphids Wooly Elm
- 38. Aphids Wooly Pear
- 39. Armyworm
- 40. Arrow-shaped Micrathena
- 41. Aster Leafhopper
- 42. Azalea caterpillar
- 43. Bagworms
- 44. Bean Leaf Beetle
- 45. Beet Armyworm
- 46. Boxelder bugs
- 47. Broadmites
- 48. Brown softscale

- 49. Budworms
- 50. Cabbage Looper
- 51. Cabbage Webworm
- 52. California Katydid
- 53. California Oakworm
- 54. California redscale (crawler)
- 55. Cankerworms
- 56. Celery looper
- 57. Centipedes
- 58. Cereal Leaf Beetle
- 59. Chinch Bug
- 60. Chinch Bug Hairy
- 61. Chinch Bug Southern
- 62. Chinch Bug False
- 63. Cicadas
- 64. Clover Mite
- 65. Colorado Potato Beetle
- 66. Corn Earworm
- 67. Corn Leaf Aphid
- 68. Corn Rootworms
- 69. Corn Rootworms Northern Larvae
- 70. Corn Rootworms Southern Larvae
- 71. Corn Rootworms Western Larvae
- 72. Corn Rootworms Mexican Larvae
- 73. Corn Silkfly (Adult)
- 74. Crickets
- 75. Crickets Arizona
- 76. Crickets Black horned Tree
- 77. Crickets California Tree
- 78. Crickets Common short-tailed
- 79. Crickets Eastern striped
- 80. Crickets Fall field
- 81. Crickets House
- 82. Crickets Indies short-tailed
- 83. Crickets Mormon
- 84. Crickets Northern wood
- 85. Crickets Sand field
- 86. Crickets Snowy Tree
- 87. Crickets Southeastern field
- 88. Crickets Southern wood
- 89. Crickets Spring field
- 90. Crickets Spotted Camel
- 91. Crickets Texas field
- 92. Crickets Tropical
- 93. Crickets Variable field
- 94. Crickets Vocal field
- 95. Crickets Western striped
- 96. Cutworms
- 97. Cutworms Army
- 98. Cutworms Black
- 99. Cutworms Bronzed
- 100. Cutworms Clover
- 101. Cutworms Granulate
- 102. Cutworms Greasy
- 103. Cutworms Green
- 104. Cutworms Spotted

- 105. Cutworms Variegated
- 106. Cutworms Western Bean
- 107. Eastern tent caterpillar
- 108. Elm leaf beetles
- 109. English Grain Aphid
- 110. European Corn Borer
- 111. Fall Armyworm
- 112. Fall webworm
- 113. Flea beetles
- 114. Flea beetles Cabbage
- 115. Flea beetles Corn
- 116. Flea beetles Grape
- 117. Flea beetles Pale striped
- 118. Flea beetles Potato
- 119. Flea beeltes Redheaded
- 120. Flea beetles Strawberry
- 121. Flea beetles Tobacco
- 122. Flower Thrips
- 123. Forest tent caterpillar
- 124. Gladiator Katydid
- 125. Grand Western Cicada
- 126. Grasshopper
- 127. Grasshopper Alutacea Bird
- 128. Grasshopper American Bird
- 129. Grasshopper Aztec Pygmy
- 130. Grasshopper Banded Range
- 131. Grasshopper Carolina
- 132. Grasshopper Creosote Bush
- 133. Grasshopper Differential
- 134. Grasshopper Dragon Lubber
- 135. Grasshopper Great crested
- 136. Grasshopper Green valley
- 137. Grasshopper Horse Lubber
- 138. Grasshopper Lubber
- 139. Grasshopper Painted
- 140. Grasshopper Pallid-winged
- 141. Grasshopper Panther Spotted
- 142. Grasshopper Pygmy
- 143. Grasshopper Red legged
- 144. Grasshopper Southeastern Lubber
- 145. Grasshopper Spur-throated
- 146. Grasshopper Three banded
- 147. Grasshopper Toothpick
- 148. Grasshopper Two striped
- 149. Green Bug
- 150. Green Cloverworm
- 151. Hop Vine Borer
- 152. Hornworm
- 153. Imported Cabbageworm
- 154. Japanese Beetle (Adult)
- 155. June beetles (Adults)
- 156. Katydid Fork tailed Bush
- 157. Katydid Keeled Shield-back
- 158. Katydid Mescalero Shield-back
- 159. Katydid Oblong-winged
- 160. Katydid Short-legged Shield back

- 161. Katydid True (Northern)
- 162. Kudzu Buas
- 163. Lace bugs
- 164. Ladybeetles
- 165. Ladybeetles Asian
- 166. Ladybeetles Convergent
- 167. Ladybeetles Seven-spotted168. Ladybeetles Thirteen-spotted
- 169. Ladybeetles Twice-stabbed
- 170. Ladybeetles Two-spotted
- 171. Leaf feeding caterpillars
- 172. Leaf rollers
- 173. Leaf skeletonizers
- 174. Leafhoppers
- 175. Leafhoppers Aster
- 176. Leafhoppers Bluntnosed
- 177. Leafhoppers Corn Blotch
- 178. Leafhoppers Eastern Grape
- 179. Leafhoppers Elm
- 180. Leafhoppers Mountain
- 181. Leafhoppers Potato
- 182. Leafhoppers Redbanded
- 183. Leafhoppers Rice
- 184. Leafhoppers Rose
- 185. Leafhoppers Sharpnosed
- 186. Leafhoppers Six-spotted
- 187. Leafhoppers Southern Garden
- 188. Leafhoppers Variegated
- 189. Leafhoppers Variegated larvae
- 190. Leafhoppers Virginia creeper
- 191. Leafhoppers Western grape
- 192. Leafhoppers Whitebanded elm
- 193. Leafminer (Adult)
- 194. Leafminers (Adults) Alder
- 195. Leafminers (Adults) Boxwood
- 196. Leafminers (Adults) Corn blotch
- 197. Leafminers (Adults) Holly
- 198. Leafminers (Adults) Oak
- 199. Leafminers (Adults) Spruce needle
- 200. Leafminers (Adults) Tentiform
- 201. Leafminers (Adults) Vegetable
- 202. Leafrollers Oblique banded
- 203. Leafrollers Red-banded
- 204. Leafrollers Variegated
- 205. Lesser Peach Tree Borer
- 206. Lygus Bug
- 207. Meadow Spittlebug
- 208. Mealybugs
- 209. Mealybugs Citrus
- 210. Mealybugs Comstock
- 211. Mealybugs Grape (crawlers)
- 212. Mealybugs Hawthorne
- 213. Mealybugs Longtailed
- 214. Mealybugs Madiera
- 215. Mealybugs Maple
- 216. Mealybugs Obscure

- 217. Mexican Corn Rootworm Beetle (Adult)
- 218. Midges
- 219. Millipedes
- 220. Millipedes Almond
- 221. Millipedes Flat
- 222. Millipedes Florida
- 223. Millipedes North American
- 224. Millipedes Soft
- 225. Mites Banks grass
- 226. Mites Broad
- 227. Mites carmine
- 228. Mites European
- 229. Mites Grain
- 230. Mites Pacific
- 231. Mites Pecan Leaf Scorch
- 232. Mites Pecan Scorch
- 233. Mites Russet
- 234. Mites Spruce
- 235. Mites Spider
- 236. Mites Two-Spotted Spider
- 237. Moths
- 238. Moths Abrupt Epiblema
- 239. Moths Alligator Weed Stem Borer
- 240. Moths Arch
- 241. Moths Artichoke plume
- 242. Moths Aster-head Phaneta
- 243. Moths Bald Cypress Coneworm
- 244. Moths Banded Sunflower
- 245. Moths Bidens Borer
- 246. Moths Black Dash Epinotia
- 247. Moths Black-Blotched Bactra
- 248. Moths Black Bordered Lemon
- 249. Moths Black-headed Birch Leafolder
- 250. Moths Bougainvillea Caterpillar
- 251. Moths Broad-patch Carolella
- 252. Moths Buff-tipped Phaneta
- 253. Moths Cactus
- 254. Moths Carmine Snout
- 255. Moths Catkinworm
- 256. Moths Clover Hayworm
- 257. Moths Codling
- 258. Moths Constricted Sonia
- 259. Moths Corn Earworm
- 260. Moths Crescent-Winged Caudellia
- 261. Moths Dagger
- 262. Moths Dame's Rocket
- 263. Moths Darker Moodna
- 264. Moths Delightful Bird Dropping
- 265. Moths Diamondback
- 266. Moths Dotted Ecdytolopha
- 267. Moths Eublemmine
- 268. Moths Evergreen Coneworm
- 269. Moths Exposed Bird Cropping
- 270. Moths Eye-spotted
- 271. Moths Filbertworm
- 272. Moths Fingered Dagger

- 273. Moths Four-spotted Bird-Dropping
- 274. Moths Fruit-tree Leafroller
- 275. Moths Fruitworm
- 276. Moths Goldenrod Gall Moth
- 277. Moths Grape Berry
- 278. Moths Grape leaffolder
- 279. Moths Gray Dagger
- 280. Moths Gray Half Spot
- 281. Moths Green Leuconycta
- 282. Moths Gypsy
- 283. Moths Half Yellow
- 284. Moths Hickory Leafroller
- 285. Moths Hickory Shuckworm
- 286. Moths Hypenines
- 287. Moths Hypenodine
- 288. Moths Iris Borer
- 289. Moths Jack Pine Budworm
- 290. Moths Kimball's Leafroller
- 291. Moths Labyrinth
- 292. Moths Lacuna
- 293. Moths Lantana Leaftier
- 294. Moths Lantana
- 295. Moths Lesser Loblotty Pineconeworm
- 296. Moths Locust Leafroller
- 297. Moths Locust Twig Borer
- 298. Moths Macrame
- 299. Moths Maple Bud Borer
- 300. Moths Maple Tip Borer
- 301. Moths Maple Webworm
- 302. Moths Maple Basswood Leafroller
- 303. Moths Marbled-Green Leuconycta
- 304. Moths Nantucket Pine Tip
- 305. Moths Oak Leafroller
- 306. Moths Oak Leaftier
- 307. Moths Oriental
- 308. Moths Pale Glyph
- 309. Moths Palm Bud
- 310. Moths Pecan Bud
- 311. Moths Pine Tube
- 312. Moths Sallow
- 313. Moths Sculptured
- 314. Moths Sunflower
- 315. Moths Sunflower Bud
- 316. Moths Tip
- 317. Moths -Verbena
- 318. Moths Waterlily
- 319. Moths Wedgling
- 320. Moths Yellow Headed Cutworm
- 321. Moths Zimmerman pine
- 322. Northern Corn Rootworm Beetle (Adult)
- 323. Oleander moth larvae
- 324. Onion Maggot (Adult)
- 325. Onion Thrips
- 326. Palmetto bugs
- 327. Peach Tree Borer
- 328. Peach Twig Borer

- 329. Pillbug
- 330. Pine sawfly
- 331. Pine shoot beetles
- 332. Pinetip moths
- 333. Root weevil
- 334. Saltmarsh Caterpillar
- 335. Sap Beetle (Adult)
- 336. Sawflies
- 337. Sawflies European
- 338. Sawflies European Pine
- 339. Sawflies Redheaded
- 340. Sawflies Redheaded Pine
- 341. Scale insects (crawlers)
- 342. Scales Beech
- 343. Scales Black
- 344. Scales Calico
- 345. Scales California Red
- 346. Scales Cottony Cushion
- 347. Scales Cottony Maple
- 348. Scales European Elm
- 349. Scales Euonymous
- 350. Scales Fletcher
- 351. Scales Hemispherical
- 352. Scales Magnolia
- 353. Scales Pine needle/Crawlers
- 354. Scales San Jose
- 355. Scales Brown Soft
- 356. Seedcorn Maggot (Adult)
- 357. Southern Armyworm
- 358. Southern Cabbageworm
- 359. Southern Corn Rootworm Beetle (Adult)
- 360. Southwestern Corn Borer
- 361. Sowbugs
- 362. Soybean looper
- 363. Spiders American House 364. Spiders Ant mimic
- 365. Spiders Barn
- 366. Spiders Black and Yellow Argiope
- 367. Spiders Bola
- 368. Spiders Branch yip
- 369. Spiders Brown Daddy Long Legs
- 370. Spiders Brownish Grey Fishing
- 371. Spiders Burrowing Wolf
- 372. Spiders California Trapdoor
- 373. Spiders Carolina Wolf
- 374. Spiders Crablike Spiny Orb Weaver
- 375. Spiders Daddy Long Legs
- 376. Spiders Daring Jumping
- 377. Spiders Desert Loxosceles
- 378. Spiders Desert Tarantula
- 379. Spiders Dimorphic Jumping Spider
- 380. Spiders Eastern Daddy Long Legs
- 381. Spiders Elegant Crab
- 382. Spiders Elongate Long- Jawed Orb Weavers
- 383. Spiders Forest Wolf
- 384. Spiders Garden (Cross)

- 385. Spiders Golden Huntsman
- 386. Spiders Goldenrod
- 387. Spiders Golden-silk (Calico)
- 388. Spiders Grass
- 389. Spiders Green Lynx
- 390. Spiders Green Lyssomanes
- 391. Spiders Hammock
- 392. Spiders Huntsman
- 393. Spiders Inconspicuous Crab
- 394. Spiders Jumping
- 395. Spiders Jumping Lynx
- 396. Spiders Long-jawed Orb Weaver
- 397. Spiders Mabel Orchard
- 398. Spiders Marbled Orb Weaver
- 399. Spiders Metaphid Jumping
- 400. Spiders Nursery Web
- 401. Spiders Orb Weavers
- 402. Spiders Rabid Wolf
- 403. Spiders Red spotted Crab
- 404. Spiders Selenopid Crab
- 405. Spiders Shamrock
- 406. Spiders Silver Argiope
- 407. Spiders Six Spotted Fishing
- 408. Spiders Six Spotted Orb Weaver
- 409. Spiders Spitting
- 410. Spiders -Thin legged Wolf Spider
- 411. Spiders Thrice banded Crab
- 412. Spiders Turret
- 413. Spiders Venusta Orchard
- 414. Spiders Wandering
- 415. Squash Stalk Borer
- 416. Squash Vine Borer
- 417. Spittlebugs
- 418. Stalk Borer
- 419. Stink Bug
- 420. Stink Bug Green
- 421. Stink Bug Two spotted
- 422. Striped beetles
- 423. Striped oakworm
- 424. Tarnished Plant Bug
- 425. Thrips
- 426. Thrips-Grape Thrips
- 427. Thrips-Grass Thrips
- 428. Thrips-Greenhouse Thrips
- 429. Thrips-Introduced Basswood Thrips
- 430. Thrips-Iris Thrips
- 431. Thrips-Lily Bulb Thrips
- 432. Thrips-Melon Thrips
- 433. Thrips-Mullein Thrips
- 434. Thrips-Pear Thrips
- 435. Thrips-Privet Thrips
- 436. Thrips-Redbanded Thrips
- 437. Thrips-Tobacco Thrips
- 438. Tobacco Budworm
- 439. Tomato Fruitworm

- 440. Tomato Pinworm
- 441. True Armyworm
- 442. Tussock moth larvae
- 443. Waterbugs
- 444. Weevils Alfalfa
- 445. Weevils Arborvitae
- 446. Weevils Asiatic Oak
- 447. Weevils Black Vine
- 448. Weevils Blue Grass
- 449. Weevils Blue Green Citrus Root
- 450. Weevils Carrot
- 451. Weevils Cow Pea Curculio
- 452. Weevils Cranberry
- 453. Weevils -Curcuilo
- 454. Weevils Diaprepes Root
- 455. Weevils Granary
- 456. Weevils Northern Pine
- 457. Weevils Obscure Root
- 458. Weevils Orchard
- 459. Weevils Orchid
- 460. Weevils Pea
- 461. Weevils Pea Leaf
- 462. Weevils Pecan
- 463. Weevils Pepper
- 464. Weevils Pine Shoot
- 465. Weevils Plum Curculio
- 466. Weevils Southern Blue Green Citrus Root
- 467. Weevils Strawberry Root
- 468. Weevils Sunflower headclipping
- 469. Weevils Sunflower seed
- 470. Weevils Sunflower stem
- 471. Weevils Sweet Potato
- 472. Weevils Twobanded Japanese
- 473. Weevils Vegetable
- 474. Weevils Woods
- 475. Weevils Yellow poplar
- 476. Western Bean Cutworm
- 477. Western Corn Rootworm Beetle (Adult)
- 478. Western Flower Thrips
- 479.
- 480. Whitefly
- 481. Yellow-striped Armyworm