

U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

| Number: | | |
|-----------|--------|-----|
| 53883-199 | SEP 22 | AMC |

Date of Issuance:

Term of Issuance: Conditional

Name of Pesticide Product:

EPA Reg.

IMI 0.5 G Insecticide

NOTICE OF PESTICIDE:

<u>x</u> Registration __ Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Ms. Jane M. Miller Biologic Inc. 115 Obtuse Hill Brookfield, Connecticut 06804

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

egistration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A). Once a pesticide is registered, however, it is not regarded as permanently acceptable. Registration does not eliminate the need for continual reassessment of pesticides. If the Agency determines that, at any time, additional data are required to maintain in effect an existing registration, the Agency will require submission of such data under FIFRA section (3)(c)(2)(B).

1. Revise the EPA Registration Number to read, EPA Reg. No. "53883-199".

Signature of Approving Official:

 ρ

Dani Daniel Insecticide-Rodenticide Branch Registration Division (7505P) ----

SEP 2 2 2009

2. Complete the "Environmental Hazards" statement as follows: "equipment wash waters or rinsate. Runoff may be hazardous to aquatic organisms in water adjacent treated areas.

Replace this sentence at the beginning of the second paragraph of the "Environmental Hazards" statement: "This product contains a chemical with properties and characteristics ..."

- 3. On page 2, under "Turfgrass" delete the site "sodfarms". The above statement says does not permit use on sodfarms nor does the parent label.
- 4. On page 3 under "Rate" correct ½ to 1/8 lb to read ¼ to 1/8 as is stated on the parent label.
- 5. You can only claim what is on the parent label. If you want to make additional claims you must submit data to support the claim. On page one "I Application provides season long control and Kills grubs all season long" does not appear on the parent label, in order to make this claim data must be submitted to support it.
- 6. Submit to the Agency the required one year storage stability (830.6317) and Corrosion Characteristic study (830.6320) for the proposed product. It is recommended that observations are made at 0, 3, 6, 9, and 12 months intervals.
- 7. Under the "Warranty" statement change the sentence "To the extent allowed by law" to read "To the extent consistent with applicable law..."
- 8. Submit two copies of your final printed label before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sect. 6(e). Your release for shipment of the product constitute acceptance of these conditions.

A stamped copy of the label is enclosed for your records. If you have any questions regarding this notice, please contact me at (703) 305-5409.

Enclosure:

ACCEPTED with COMMENTS In EPA Letter Dated: 3/8
SEP 2 2 2006

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the postleide registered under EPA Reg. No.

IMI 0.5 G Insecticide

For systemic insect control in turfgrass and landscape ornamentals.

Optional marketing language in brackets: [large picture of a grub on the label] [Include a picture of the grub life cycle]

[10 Pounds Treats 7000 Sq Ft]
[Controls Billbugs, Green June Beetles and Mole Crickets]
[1 Application Provides Season Long Control]
[Apply anytime between Spring throughout Summer]
[Kills Grubs all Season long]
[<PRODUCT NAME> contains imidacloprid, the active ingredient used in <BRAND NAME>TM or ®.]
[<PRODUCT NAME> is not manufactured or distributed by <BASIC REGISTRANT / BRAND HOLDER>, seller of <BRAND>TM or ®.]
[<BRAND>TM or ® is a trademark of <TRADEMARK HOLDER>>.

ACTIVE INGREDIENT:

| *Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]-N- | |
|--|--------------|
| nitro-2-imidazolidinimine | 0.5% |
| OTHER INGREDIENTS: | 99.5% |
| Total: | 100.0% |
| EPA Reg. No. 53883-XXX EPA Est No. XXXXX-XX-XXX | Net contents |

STOP - Read the label before use. Keep out of reach of children.

CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

| | FIRST AID |
|---------------------------|--|
| If swallowed | Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. |
| If on skin or clothing | Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. |
| If in eyes | Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. |
| | ency call Poison Control Center at 800-222-1222. Have a product container or label |
| | lling a poison control center or doctor, or going for treatment. an: No specific antidote is available. Treat the patient symptomatically. |

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not formulate this product into other end-use products.

NOTE: Not for use in commercial greenhouses, nurseries, on sod farms or on grass grown for seed. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

TURFGRASS

IMI 0.5 G Insecticide is for use on turfgrass in home lawns, business and office complexes, shopping complexes, multi-family residential complexes, golf courses, airports, cemeteries, parks, playgrounds, athletic fields and sod farms to control soil inhabiting pests as listed in the table below. IMI 0.5 G Insecticide can also be used for the suppression of cutworms and chinchbugs in turfgrass areas.

APPLICATION INSTRUCTIONS

Effective control can be achieved when applications are made before or during the egg laying period. The timing of applications can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. For best control apply prior to egg hatch of the target pests, followed by sufficient irrigation or rainfall to move the active ingredient through the thatch.

NOTE: For optimum control, irrigation or rainfall should occur within 24 hours after application to move the active ingredient through the thatch. Do not make applications when turfgrass areas are waterlogged or the soil is saturated with water. Adequate distribution of the active ingredient cannot be achieved when these conditions exist. The area to be treated must be able to accept vertical penetration of rainfall or irrigation water. Avoid mowing turf or lawn area until after sufficient irrigation or rainfall has occurred so that uniformity of application will not be affected.

Do not apply more 80 lb (0.4 lb of active ingredient) per acre per year.

APPLICATION EQUIPMENT

Apply IMI 0.5 0 Insecticide uniformly over the area being treated with granular application equipment. Both drop-type and rotary-type spreaders may be used. Do not use spreaders which apply the material in narrow, concentrated bands. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly and accurately.

Consult your local State Agricultural Experiment Station, or State Extension Turf Specialists for more specific information regarding timing of application.

| RECOMMENDED APPLICATIONS OF IMI 0.5 G INSECTICIDE ON TURFGRASS | | | |
|--|---|--|--|
| PEST | USE DIRECTIONS | RATE | |
| Larvae of: European Crane Fly, (Tipula paludosa) Annual bluegrass weevil, (Hyperodes spp.) Green June Beetle, (Cotinis nitida); May or June beetle, (Phyllophaga spp.); Asiatic garden beetle, (Maladera castanea) Japanese Beetle, (Popllia japonica) Billbug, (Spherophorus spp.) Northern masked chafer and Southern masked chafer, (Cyclocephala borealis, C. immaculata, and/or C. lurida) Black turfgrass ataenius, (Ataenius spretulus and Aphodius spp.) Oriental Beetle, (Anomala orientalis) Cutworms (suppression), (Phyllophaga spp.) European chafer, (Rhizotroqus majalis) | For optimum control of grubs, billbugs, european crane fly and annual bluegrass weevil, make application prior to egg hatch of the target pest. | 60 to 80 lb per acre or 1.2 to 1.8 lb per 1000 sq ft | |
| Chinchbug (suppression) Mole Crickets, (Scapteriscus spp.) | For suppression of chinchbugs, make application prior to hatching of the instar nymphs. For control of mole crickets make application prior to or during the peak egg hatch period. When adults or large nymphs are present and actively tunneling, application should be accompanied by a curative insecticide. | 80 lb per acre or 1.8 lb per 1000 sq ft | |

ORNAMENTALS

IMI 0.5 G Insecticide can be used as a broadcast application on ornamentals in commercial and residential landscapes.

APPLICATION INSTRUCTIONS

IMI 0.5 G Insecticide is a systemic product which is translocated upward within the plant system. For best control, follow applications with sufficient mechanical incorporation, irrigation or rainfall to



move the active ingredient into the soil. Irrigation or rainfall should occur within 24 hours after application. Apply prior to pest infestation to achieve optimum control.

Apply as a broadcast treatment. Prior to application, remove black plastic or other weed barrier material that will not permit water permeability.

NOTE: Avoid runoff or puddling of irrigation water following application. Avoid application of **IMI 0.5 G Insecticide** to areas which are water logged or saturated, which will not allow penetration into the root zone of the plant.

Do not apply more than 80 lb (0.4 lb of active ingredient) per acre per year.

APPLICATION EQUIPMENT

Apply **IMI 0.5 G Insecticide** uniformly over the area being treated with granular application equipment. Both drop-type and rotary-type spreaders may be used to apply **IMI 0.5 0 Insecticide**. Do not use the type of spreaders which would apply the material in narrow, concentrated bands. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly and accurately.

| | CATION OF IMI 0.5 G INSECTICIDE ON ORNAMENT and around industrial and commercial buildings and re | |
|--|--|---------------------------------|
| CROP | PEST | RATE |
| Landscape Plantings (including trees, shrubs evergreens, flowers, foliage plants, and ground covers) | White grub larvae (such as Japanese beetle larvae, Chafers, Phyllophaga spp. Asiatic garden beetle, Oriental beetle) | 1.4 to 1.8 lb per 1000 sq ft |

| Spread | er | Set | tir | ıgs |
|--------|----|-----|-----|-----|
|--------|----|-----|-----|-----|

The following chart can be used as a guide in establishing an accurate rate of IMI 0.5 G Insecticide. Application rates will vary depending upon speed, equipment wear and condition of spreader. Carefully calibrate application equipment prior to commercial use of this product.

| SPREADER | SPEED | SETTINGS | | |
|-------------------------------------|-------|------------|------------|--|
| | (mph) | 60 lb/acre | 80 lb/acre | |
| Cyclone Model CB4000 | 3 | 3 1/4 | 3 1/2 | |
| Earthway Drop | 3 | 9 | 10 | |
| Earthway 10' Rotary | 3 | 11 | 12 | |
| GandyTurf Tender24" | 3 | 16 | 19 | |
| GandyTurf Tender 42" | 3 | 21 | 23 | |
| Lely | 4.5 | 3.5 | + | |
| Lesco Model 6000 | 3 | B 1/2 | С | |
| PS8 Prizelawn CBR-II | 3 | 3 1/4 | 3 ½ | |
| Scotts Proturf R-8A | 3 | Н | Ī | |
| Scott's Proturf SR-1 | 3 | H | | |
| Scott's PF-1, PF-2, PF-3, PF-4 Drop | 3 | 4 | 4 1/2 | |
| Scott's SS-1 Drop | 3 | 4 | 4 1/2 | |
| Spyker Model 34 | 3 | 3.1 | 3.2 | |
| Vicon Model PS-203 | 4.5 | 13 | 16 | |

RESTRICTIONS

Treated areas may be replanted with any crop specified on an imidacloprid label, or with any crop for which a tolerance exists for the active ingredient.

For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12- month plant-back interval should be observed.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Carefully handle the container to prevent spills. If material is spilled, carefully contain any material to prevent non-target contamination. Do not walk through spilled material and dispose of as directed for pesticides below.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill, by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Refer to Precautionary Statements on label for hazards associated with the handling of this material. Contact CHEMTREC at 800-424-9300 for decontamination procedures or any other assistance that may be necessary.

IMPORTANT READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of liability before using this product.

If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following conditions disclaimer of warranties and limitations of liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, because of manner of use and other factors beyond Control Solutions Inc.'s (CSI) control it is impossible for Control Solutions Inc. to eliminate all risks associated with the use of this product As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

WARRANTY STATEMENT

Control Solutions, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Control Solutions, Inc. To the extent allowed by law, Control Solutions, Inc. shall in no event be liable for consequential, special, or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. In addition to the foregoing, no purchaser of this product (other than an end user) shall be entitled to any reimbursement for any loss suffered as a result of any suspension or cancellation of the registration for this product by the U.S. Environmental Protection Agency. Except, as expressly provided herein. Control Solutions, Inc. makes no warranties, quarantees, or representations of any kind, either expressed or implied, or by usage of trade, statutory or otherwise, with regard to the product sold, including, but not limited to merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage. The exclusive remedy of any buyer or user of this product for any and all

losses, injuries, or damage resulting from or in any way arising from the use, handling, or application of this product, whether in contract, warranty, tort, negligence, strict liability, or otherwise, shall be damages not exceeding the purchase price paid for this product or, at Control Solutions, Inc. election, the replacement of this product.

Control Solutions, Inc. 5903 Genoa-Red Bluff Pasadena, TX 77507