

53883-211

01/30/2007

1117



EPA Reg. Number: 53883-211

Date of Issuance:

JAN 30 2007

Term of Issuance: Conditional

U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave. NW
Washington, D.C. 20460

Name of Pesticide Product:
Dithiopyr 0.06% Plus Fertilizer

NOTICE OF PESTICIDE:

X Registration
Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Control Solutions, Inc
5903 Genoa-Red Bluff
Pasadena, TX 77507

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label. Comments on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

- 1. Add the phrase "EPA Registration No. 53883-211 to the label before you release the product for shipment.

COMMENTS CONTINUED ON PAGE 2 OF THIS NOTICE OF REGISTRATION

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product under the enclosed stamped copy of the label constitutes acceptance of these conditions.

Enclosure

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505P)

Signature of Approving Official:

Joanne I. Miller

Date:

JAN 30 2007

2/17

Page 2

EPA Reg. No. 53883-211

Comments Continued:

2. Submit to this Agency a copy of the purchasing order or bill of sales showing the source of the product named on the Formulator's Exemption Statement, and with which this product is to be manufactured; prior to manufacturing this product.
 3. Correct the typographic error in the chemical expression for dithiopyr in the ingredient statement by correcting the spelling of "...pyridinedicarbothioate...".
 4. Submit one (1) copy of the final printed labeling before you release this product for shipment.
 5. Submit and/or cite all data required for the registration of this product when the Agency requires all registrants of similar products to submit data; and submit acceptable responses required for reregistration of this product under FIFRA, section 4.
-

3/17

Dithiopyr 0.06% Plus Fertilizer

Turf and Ornamental Herbicide

For Season-Long Control of Crabgrass and Control or Suppression of Listed Annual Grasses and Broadleaf Weeds in Established Lawns and Ornamental Turf, including Golf Course Fairways, Roughs and Tee Boxes, and in Established Landscape Ornamentals.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

In the state of New York this product may be applied only by commercial applicators at no more than 19.1 lbs. per 1,000 square feet (0.5 pounds AI per acre) per year. Not for use in Nassau or Suffolk Counties.

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

JAN 30 2007
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

Active Ingredient:

Dithiopyr: S,S'-dimethyl 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-3,5-pyridinedicarbothioate 0.06%

Inert Ingredients:..... 99.94%

TOTAL:..... 100.00%

53883-211

EPA REG NO. 53883-XXX

EPA EST. NO. XXXXX-XXX-XXX

Net Weight:

Control Solutions, Inc.
5903 Genoa-Red Bluff Rd.
Pasadena, TX 77507-1041

Guaranteed Fertilizer Analysis: XX-XX-XX

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See [back] [side] [other] panel for further precautions and First Aid information

4/17

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if inhaled. Causes moderate eye irritation. Avoid breathing dust. Remove and wash contaminated clothing before reuse. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Wear long-sleeved shirt and long pants, sock, shoes and chemical-resistant gloves such as natural rubber. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or physician immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or physician. • Do not give anything by mouth to an unconscious person.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.	

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION

Dithiopyr 0.06% Plus Fertilizer provides season-long control of crabgrass and control or suppression of listed annual grasses and broadleaf weeds in established lawns and ornamental turf, including golf course fairways, roughs and tee boxes, and in established landscape ornamentals. This product also provides postemergent crabgrass control up to the 4-leaf stage.

SPREADER SETTINGS to apply at the listed pounds per 1,000 square feet
(to be furnished at printing)

FOR USE ON TURF

Dithiopyr 0.06% Plus Fertilizer provides season-long control of crabgrass and control or suppression of other annual grasses and broadleaf weeds in established lawns and ornamental turf, including golf course fairways, roughs and tee boxes. Dithiopyr 0.06% Plus Fertilizer will not harm nearby established ornamentals identified on this label as "Tolerant Ornamentals" when used according to label directions.

Apply this product only to seeded, sodded, or sprigged lawns and ornamental turf that is well established. Before the use of Dithiopyr 0.06% Plus Fertilizer, the grass must have a developed root system and a uniform stand. Following seeding, sodding or sprigging, the turf must be mowed at least two times. Injury may result if the turf is not well established, or stressed by weather, pests, disease, chemical or other related conditions.

Dithiopyr 0.06% Plus Fertilizer may be applied to established turfgrass species listed on this label. When applied as directed, the following turfgrass species are tolerant to this product:

COOL-SEASON GRASSES	USE PRECAUTIONS
Creeping Bentgrass ¹ (<i>Agrostis palustris</i>), Kentucky Bluegrass (<i>Poa pratensis</i>), Fine Fescue ² (<i>Festuca rubra</i>), Tall Fescue (<i>Festuca arundinacea</i>), Perennial Ryegrass (<i>Lolium perenne</i>),	¹ Turfgrass injury may result from the use of this product on certain varieties of creeping bentgrass such as Cohansey, Carmen, Seaside and Washington. Not all varieties of creeping bentgrass have been testes. Do not apply this product to Colonial Bentgrass (<i>Agrostis tenuis</i>) varieties. ² Turfgrass injury may result from use of this product on the following varieties of Fescue such as Atlanta, Banner, Beauty, Bilgart, CF-2, Enjoy, HF-93, Highlight, Ivalo, Koket, Jamestown, Majenta, Mary, Pennilawn, Tamara, Tatiana, Walfort and Waldina. Not all varieties of fescue have been tested.
WARM-SEASON GRASSES	
Bahiagrass (<i>Paspalum notatum</i>), Bermudagrass (<i>Cynodon dactylon</i>), Buffalograss ³ (<i>Buchloe dactyloides</i>), Carpetgrass (<i>Axonopus compressus</i>), Centipedegrass (<i>Eremochloa ophiuroides</i>), Kikuyugrass (<i>Pennisetum clandestinum</i>), St. Augustinegrass (<i>Stenotaphrum secundatum</i>), Zousiagrass (<i>Zoysia japonica</i>)	³ Do not use this product on seedling Buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

NOTE: It is not possible to evaluate all turfgrass varieties for tolerance to this product. It is suggested that the user evaluate the response to turfgrass varieties not listed on this label on a limited area, at recommended use rates, prior to initiating large-scale use.

RESEEDING, OVERSEEDING OR SPRIGGING

Reseeding, overseeding, or sprigging of treated areas within 10 weeks after single application or within 16 weeks after a split application program may inhibit the establishment of desirable turfgrasses. When reseeding or overseeding, follow good cultural practices such as soil cultivation, irrigation and fertilization. For best results, use mechanical or power seeding equipment (slit seeders) for good seed-to-soil contact.

USE RESTRICTIONS

- Do not apply more than 19.1 lbs./1,000 sq. ft. (0.5 Lbs. Al/acre) per application.
- Do not apply more than 57.3 lbs./1,000 sq. ft. (1.5 lbs. Al/acre) per year.
- In New York, do not apply more than 19.1 lbs. /1,000 sq. ft. per year; and is prohibited from use in Nassau and Suffolk Counties.
- Do not use clippings from treated turf for animal feed or for mulching around vegetables or fruit trees.
- Do not allow domestic animals to feed or forage on treated turf.
- Keep people and pets off treated areas until dust has settled.
- Do not use on golf course putting greens.

CRABGRASS CONTROL

Dithiopyr 0.06% Plus Fertilizer provides pre-emergence control of crabgrass (large, smooth, and southern species) when applied prior to emergence of crabgrass from the ground. Dithiopyr 0.06% Plus Fertilizer provides "early post-emergence" control during the early stages of growth after crabgrass emerges from the ground. This early post-emergence control is limited and applications are only effective if applied before the crabgrass has its 3-leaf stage or first tiller (which usually is about the time when you can first see the crabgrass in the lawn). With this post-emergence activity, the applicator has 2 to 4 weeks longer to make applications for control of crabgrass.

APPLICATION INSTRUCTIONS

For best results, apply Dithiopyr 0.06% Plus Fertilizer within a few days after mowing and delay mowing again for a few days after application. Rainfall or irrigation activates the product. Improved weed control will result if the application is followed by 1/2 inch of rainfall or sprinkler irrigation. Weed control may be hampered if rainfall or irrigation does not occur with 30 days after application. Split, or half rate applications spaced 5 to 10 weeks apart may also enhance weed control. Cultural practices such as aerification and de-thatching should be completed and the turf allowed to recover before application of this product.

APPLICATION TIMING

This product may be applied as a single application, split application, or as a sequential application for crabgrass control in the spring, summer, or fall as described below.

SPRING APPLICATIONS: For single applications in the spring or early summer, follow the appropriate rate listed in "TABLE 1: APPLICATION RATES FOR CRABGRASS CONTROL" below. Rates are based on the location to be used (Region), the mowing height of the turfgrass, and if the timing of the application is considered preemergence or postemergence. The length of residual weed control is dependent on the total rate applied, but may vary depending on weather, weed pressure and location.

Program 1: For preemergence control to turf mowed relatively high, such as home lawns, this program will provide 3 to 5 months of preemergence crabgrass control. Do not use this program for early postemergence control of crabgrass.

Program 2: For preemergence control at sites where the turf is mowed relatively low, such as golf course fairways, and where turfgrass maintenance and weed control applications were not made the previous year. This program will provide 4 to 6 months of preemergence crabgrass control and may also be used for early postemergence control of crabgrass.

Program 3: For preemergence control at locations where turf is mowed relatively low, such as golf course fairways, and where turfgrass maintenance and weed control applications were not made the previous year. This program will provide 4 to 6 months of preemergence crabgrass control and may also be used for early postemergence control of crabgrass at locations where the turf is cut relatively low.

FALL APPLICATIONS: Dithiopyr 0.06% Plus Fertilizer may be applied in late summer or early fall (late August through November). Follow the instructions for "Program 3" for crabgrass control through the

early part of the next spring. Fall applications may be followed by an appropriate spring application to provide season-long control.

[Optional Language:

TURF GRASS REGIONS

Insert map showing regions for crabgrass control as noted in table below]

TABLE 1: APPLICATION RATES FOR CRABGRASS CONTROL

REGION	PROGRAM 1 Preemergence (high-cut turf)	PROGRAM 2 Preemergence (low-cut turf) Postemergence (high-cut turf)	PROGRAM 3 Preemergence (low-cut turf) Postemergence (low-cut turf)
North (all areas except NY)	7.3 lbs./1,000 sq. ft. (317 lbs./acre or 0.19 lb. Al/acre)	9.6 lbs./1,000 sq. ft. (417 lbs./acre or 0.25 lb. Al/acre)	14.5 lbs./1,000 sq. ft. (633 lbs./acre or 0.38 lb. Al/acre)
Transition (DE, KS, KY, MD, MO, VA, southeastern PA, southern areas of IL, IN, OH, and coastal areas of CT, NY and RI.	9.6 lbs./1,000 sq. ft. (417 lbs./acre or 0.25 lb. Al/acre)	14.5 lbs./1,000 sq. ft. (633 lbs./acre or 0.38 lb. Al/acre)	19.1 lbs./1,000 sq. ft. (833 lbs./acre or 0.5 lb. Al/acre)
South (AL, AR, GA, LA, MS, NC, NM, OK, SC, TN and TX)	14.5 lbs./1,000 sq. ft. (633 lbs./acre or 0.38 lb. Al/acre)	9.6 + 9.6* lbs./1,000 sq. ft. (417 + 417 lbs./acre or 0.25 + 0.25 lb. Al/acre)	11.9 + 11.9* lbs./1,000 sq. ft. (517 + 517 lbs./acre or 0.31 + 0.31 lb. Al/acre)
Coastal South (HI, FL, southern coastal areas of AL, GA, LA, MS, NC, SC, and TX)	9.6 + 9.6* lbs./1,000 sq. ft. (417 + 417 lbs./acre or 0.25 + 0.25 lb. Al/acre)	11.9 + 11.9* lbs./1,000 sq. ft. (517 + 517 lbs./acre or 0.31 + 0.31 lb. Al/acre)	14.5 + 14.5* lbs./1,000 sq. ft. (633 + 633 lbs./acre or 0.38 + 0.38 lb. Al/acre)
West AZ, CA, & NV. In this climatically diverse region, use the higher rates in local areas with longer crabgrass seasons.	7.3 + 9.6* lbs./1,000 sq. ft. (317 + 417 lbs./acre or 0.19 + 0.25 lb. Al/acre)	9.6 + 14.5* lbs./1,000 sq. ft. (417 + 633 lbs./acre or 0.25 + 0.38 lb. Al/acre)	9.6 + 9.6* lbs./1,000 sq. ft. (417 + 417 lbs./acre or 0.25 + 0.25 lb. Al/acre)
* Split applications may be made 5-10 weeks apart.			

SPLIT OR SEQUENTIAL APPLICATIONS

Split or sequential applications may be used to provide improved weed control or to provide extended weed control in areas with long growing seasons. Split applications may be made with rates in Table 1 being split between two applications made 5-10 weeks apart. Where longer periods of control are

required, sequential applications may be made in late summer or fall following a spring application or in the spring following a fall application provided the maximum application rate per year is not exceeded. Refer to the "Use Restrictions" section above.

CONTROL OF OTHER GRASS AND BROADLEAF WEEDS

SPRING APPLICATION: When used as directed for preemergence (prior to germination) control of crabgrass in the spring, Dithiopyr 0.06% Plus Fertilizer will also control or suppress the following grasses and broadleaf weeds at the application rates listed in Table 1 above:

GRASSES	
Common Name	Botanical Name
Barley	<i>Hordeum</i> spp.
Barnyardgrass	<i>Echinochloa crus-galli</i>
Bluegrass (annual roughstalk)	<i>Poa annua</i>
Brome	<i>Bromus</i> spp.
Crabgrass, large	<i>Digitaria sanguinalis</i>
Crabgrass, smooth	<i>Digitaria ischaemum</i>
Crabgrass, Southern	<i>Digitaria ciliaris</i>
Crowfootgrass*	<i>Dactyloctenium aegyptium</i>
Dallisgrass (seedling)	<i>Paspalum dilatatum</i>
Foxtail (yellow & green)	<i>Setaria faberi, Setaria Verdi</i>
Goosegrass	<i>Eleusine indica</i>
Kikuyugrass*	<i>Pennisetum clandestinum</i>
Oats, wild	<i>Avena fatua</i>
Ryegrass (annual & perennial)	<i>Lolium</i> spp.
Sandbur	<i>Cenchrus</i> spp.
Smutgrass	<i>Sporobolus indicus</i>
BROADLEAF WEEDS	
Bittercress*	<i>Cardamine</i> spp.
Carpetweed	<i>Mullugo verticillata</i>
Chickweed*	<i>Stellaria</i> spp.
Geranium, Carolina*	<i>Geranium carolinianum</i>
Henbit	<i>Lamium</i> spp.
Knotweed, prostrate	<i>Polygonum aviculare</i>
Lespedeza, common*	<i>Lespedeza striata</i>
Marestail	<i>Conyza Canadensis</i>
Medic, black	<i>Medicago lupulina</i>
Mustard	<i>Brassica</i> spp.
Oxalis, buttercup	<i>Oxalis pes-caprae</i>
Parsley-piert*	<i>Alchemilla arvensis</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Pineappleweed*	<i>Matricaria matricarioides</i>
Purslane, common	<i>Portulaca oleracea</i>
Rocket, London	<i>Sisymbrium irio</i>
Shepherdspurse	<i>Capsella bursa-pastoris</i>
Speedwell, corn*	<i>Veronica arvensis</i>
Spurge, garden	<i>Euphorbia hirta</i>
Spurge, prostrate	<i>Euphorbia humistrata</i>
Spurge, spotted	<i>Euphorbia maculate</i>
Woodsorrel, creeping	<i>Oxalis corniculata</i>
Woodsorrel, yellow	<i>Oxalis stricta</i>
* Suppression only.	

9/17

FALL APPLICATION: When used for preemergence (prior to germination) control of crabgrass in late summer or early fall (late August through November), Dithiopyr 0.06% Plus Fertilizer will also control or suppress the following weeds:

WEEDS CONTROLLED	
Common Name	Botanical Name
Bluegrass (annual)	<i>Poa annua</i>
Bittercress	<i>Cardamine</i> spp.
Chickweed	<i>Stellaria</i> spp.
Geranium (Carolina)	<i>Geranium carolinianum</i>
Henbit	<i>Lamium</i> spp.
Parsley-piert	<i>Alchemilla arvensis</i>
Pineappleweed	<i>Matricaria matricarioides</i>
Shepherdspurse	<i>Capsella bursa-pastoris</i>

APPLICATION EQUIPMENT AND INSTRUCTIONS

This product may be applied with drop or rotary-type spreaders designed to apply granular herbicides. For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps. Avoid the use of spreaders that tend to apply granules in narrow rows or concentrated bands. Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended application rate under actual application conditions. The desired calibration setting may be marked or recorded for future reference.

Apply this product uniformly over the treatment area. More uniformity can be achieved by applying one-half of the required amount of product over the treatment area and then applying the remaining one-half in a different direction (e.g., at a right angle to the previous direction). Check equipment frequently to verify calibration and proper functioning. Clean and lubricate spreader after use.

FOR USE ON LANDSCAPE ORNAMENTALS

Dithiopyr 0.06% Plus Fertilizer provides preemergence control or suppression of annual grass and broadleaf weeds in landscape ornamentals. Refer to "For Use on Turf" section of this label for a complete listing of weeds controlled or suppressed. This product may be applied to soil surfaces or mulch where ornamental plants are grown for aesthetic purposes in ornamental gardens, parks, golf courses, and residential areas.

APPLICATION INSTRUCTIONS

Dithiopyr 0.06% Plus Fertilizer is a preemergence herbicide that will not control established weeds. Make applications prior to weed seed germination to bare ground or mulch. For best results, apply to soil that is free of clods, weeds and debris such as leaves. Before application, any existing vegetation should be removed by hand weeding, cultivation, or with the use of a postemergence herbicide. Once the application is made, do not disturb the soil surface as the herbicide barrier will be broken. Irrigate immediately after application to remove herbicide particles from the foliage to prevent foliar damage to ornamentals.

SPLIT OR SEQUENTIAL APPLICATIONS

Split or sequential applications may be used to provide improved weed control or to provide extended weed control in areas with long growing seasons. Split applications may be made with rates in Table 2 being split between two applications made 5-10 weeks apart. Where longer periods of control are required, sequential applications may be made in late summer or fall following a spring application or in the spring following a fall application provided the maximum application rate per year is not exceeded.

USE RESTRICTIONS

- Only apply this product to established ornamentals.
- DO NOT apply this product directly to the bare roots of ornamental plants as injury may result.

- DO NOT incorporate this product into soil. Dilution of active ingredient and plant root injury may occur.
- DO NOT apply to soil around ornamentals that are under stress due to drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, winter injury, injury from previously applied pesticides or injury due to insects, nematodes, or disease.
- DO NOT apply under conditions that would affect product distribution (such as windy conditions).
- DO NOT apply this product on grasses grown for seed.
- DO NOT graze livestock or feed forage cut from areas treated with this product.

USE RATES

Apply Dithiopyr® 0.06% Plus Fertilizer with a properly calibrated spreader that will provide accurate, even particle distribution. Refer to Table 2 below for product rates when treating a small area.

DO NOT apply more than 19.1 lbs. of this product per 1,000 sq. ft., per application and no more than 57.3 lbs. of this product per 1,000 sq. ft. per year.

In New York State, this product may only be used by Professional Applicators at no more than 19.1 lbs. per 1,000 sq. ft. (0.5 lbs. active ingredient per acre) per year, and is prohibited from use inn Nassau and Suffolk Counties.

TABLE 2: APPLICATION RATES FOR LANDSCAPE ORNAMENTALS

Pounds Product/Acre	Pounds Product/1,000 sq. ft.	Pounds Product/100 Sq. Ft.
833	19.1	1.91 (30.6 oz.)

APPLICATION EQUIPMENT AND INSTRUCTIONS

This product may be applied with drop or rotary-type spreaders designed to apply granular herbicides. For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps. Avoid the use of spreaders that tend to apply granules in narrow rows or concentrated bands. Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended application rate under actual application conditions. The desired calibration setting may be marked or recorded for future reference.

Apply this product uniformly over the treatment area. More uniformity can be achieved by applying one-half of the required amount of product over the treatment area and then applying the remaining one-half in a different direction (e.g., at a right angle to the previous direction). Check equipment frequently to verify calibration and proper functioning. Clean and lubricate spreader after use.

TOLERANT ORNAMENTALS

When applied as directed under the conditions described on this label, the ornamentals listed below have shown tolerance in field trials. However, this product has not been tested on all cultivars of each species or under all possible growing conditions. A limited area involving only a few plants should be treated for tolerance evaluations before large-scale applications.

Common Name	Botanical Name	Tolerant Cultivars
Abelia, Dwarf	<i>Abelia X grandiflora</i>	Nana
Ajuga	<i>Ajuga reptans</i>	Bronze
	<i>Ajuga Genevensis</i>	Bronze Beauty
Almond, flowering	<i>Prunus gladulosa</i>	
Apple*	<i>Malus pumila</i>	
Arborvitae	<i>Thuja occidentalis</i>	Nigra
		Pyramidalis
		Smaragh
		Techny
		Woodwardii
Arborvitae, Golden	<i>Thuja orientalis</i>	
Aster, Chinese	<i>Callistephus chinensis</i>	Dwarf Queen

<p>Ash, Green Ash, Mountain Ash, Purple Azalea</p>	<p><i>Fraxinus pennsylvanica</i> <i>Sorbus aucuparia</i> <i>Fraxinus Americana</i> <i>Rhododendron</i> spp.</p>	<p>Brilliant Buccaneer Carror Chimes (Belgian) Elsie Lee Exbury Fashion Hardijzer Beauty Hershey Red Higasa Hinocrimson Holland (Hybrid) Marion Lee Northern Lights Orange Cup Orchid Lights Snow Southern Charm</p>
<p>Azalea, Flame Azalea, Kirishima Bamboo, Heavenly Barberry</p>	<p><i>Rhododendron calendulaceum</i> <i>Nandina domestica</i> <i>Berberis thunbergii</i></p>	<p>Aurea Dwarf Pygmy Green Koboid Pygmy Red Rose Glow Atropurpurea</p>
<p>Barberry, Purple Basket Flower Bearberry (common) Bee Balm Begonia Birch, River Blackeyed Susan Blanket Flower Blueberry*</p>	<p><i>Berberis vulgaris</i> <i>Gaillardia grandiflora</i> <i>Arctostaphylos uva-ursti</i> <i>Monarda didyma</i> <i>Begonia</i> spp. <i>Betula nigra</i> <i>Rudbeckia hirta</i> <i>Gaillardia</i> spp. <i>Vaccinium</i> spp.</p>	<p>Massachusetts Goldstrum</p>
<p>Bottlebrush Boxwood, Japanese Boxwood, Weller Broom Broom Bugle Carpet Carnelia</p>	<p><i>Callistemon citrinus</i> <i>Buxus microphylla</i> <i>Buxus sempervirens</i> <i>Cytisus</i> spp. <i>Genista pilosa</i> <i>Camellia japonica</i></p>	<p>Japonica Moonlight Vancouver Gold Debutanta Mathotiana Supreme Chansonette</p>
<p>Candy Tuft Carex, Variegated Cedar, Red</p>	<p><i>Camelia sasangua</i> <i>Ilberis</i> spp. <i>Carex</i> <i>Juniperus virginiana</i></p>	<p>Snow White</p>

Celosia	<i>Celosia</i> spp.	Scarlet Plumosa
Centaura	<i>Centaurea montana</i>	Red Kewpie
Cockscomb, Plumosa	<i>Celosia cristata</i>	
Coleus	<i>Coleus blumei</i>	
Columbine	<i>Aquilegia</i> spp.	Moonbloom
Copper leaf	<i>Acalypha wilkesiana</i>	
Coreopsis	<i>Coreopsis</i> spp.	
Cornflower	<i>Centaurea</i> spp.	
Cotoneaster	<i>Ctoneaster apiculatus</i>	
Coyotebrush	<i>Baccharis pelularis</i>	
Cycads	<i>Cycads revolute</i>	
Cypress, Bald	<i>Taxodium distichum</i>	Glauca
Cypress, Italian	<i>Cupressus sempervirens</i>	Gracilis
Cypress, Japanese False	<i>Chamaecyparis obtuse</i>	
Cupress, Leyland	<i>Cupressocyparis leylandii</i>	King Alfred
Daffodil	<i>Narcissus</i> spp.	Aztec Gold
Daylilly	<i>Hemerocallis</i> spp.	Bright Yellow (Hybrid)
		Single Gold (Evergreen)
		Wilson's Yellow
		Magic Fountain
Delphinium	<i>Delphinium</i> spp.	
Dianthus (Sweet William)	<i>Dianthanthus</i> spp.	
Dogwood	<i>Cornus florida</i>	
Dogwood, American	<i>Cornus sericea</i>	Flavarimaea
Douglas Fir	<i>Psuedotsuga menziesil</i>	
Dusty Miller	<i>Senecio cineraria</i>	
Elm, Drake	<i>Ulmus parvifolia</i>	
Euonymus	<i>Euonymus fotunei</i>	Argenteo-variegata
		Auereo-marginata
		Colorata
		Emerald Gaiety
		Emerald 'N Gold
		Gold Edge
		Gold Princess
		Silver King
		Tricolor
		Vegetus
Fan Palm, European	<i>Chamaerops humilis</i>	
Fan, Plam, Mexican	<i>Washingtonia robusta</i>	
Fern (various)	<i>Asparagus</i> spp.	
Fescue	<i>Festuca glauca</i>	Rainbow
Fetterbush	<i>Leucothoe fontiana</i>	Nitidia
Ficus	<i>Ficus retusa</i>	
Fir, Fraser	<i>Abies fraseri</i>	
Forsythia	<i>Forsythia X intermeida</i>	Arnold Dwarf
		Bronxensis Dwarf
		Lynwood Gold
		Meadowlark
		Weeping
Fuchsia	<i>Spring Glory</i>	
Galium	<i>Fuchsia</i> spp.	
Gardenia	<i>Galium ordoratum</i>	Mystery
	<i>Gardenia jasminoides</i>	Radicans
Geranium	<i>Pelargonium X hortorum</i>	

<p>Gum Hawthorn</p>	<p><i>Eucalyptus citriodora</i> <i>Crataegus spp.</i></p>	<p>Cockspur White Crimson Cloud Enchantress Jack Evans Washington White Mediterranean Pink</p>
<p>Heather, Twisted Hemlock, Canada Hibiscus</p>	<p><i>Erica cinerea</i> <i>Tsuga canadensis</i> <i>Hibiscus spp.</i></p>	<p>Blue Bird Brilliant Hula Girl Blue Boy Blue Girl Burfodii China Girl Compacta Forsteri Hellerie Japanese Northern Beauty Needlepoint Nellie R. Stevens Savanah</p>
<p>Holly</p>	<p><i>Ilex spp.</i> <i>Ilex X meserveae</i></p>	<p>Blue Bird Brilliant Hula Girl Blue Boy Blue Girl Burfodii China Girl Compacta Forsteri Hellerie Japanese Northern Beauty Needlepoint Nellie R. Stevens Savanah</p>
<p>Holly, Chinese Holly, Japanese Holly, Yaupon Honeysuckle</p>	<p><i>Ilex X attenuata</i> <i>Ilex cornuta</i> <i>Ilex crenata</i> <i>Ilex vomitoria</i> <i>Lonicera japonica</i></p>	<p>Clavey's Dwarf Hilliana Tatarian Canadian White Zebelli Red Hosta Albo Marginata</p>
<p>Hosta</p>	<p><i>Hosta spp.</i></p>	<p>Dwarf Blue Wedgewood Bulgaria Thorndale</p>
<p>Ice Plant Impatiens Iris</p>	<p><i>Hosta lancifolia</i> <i>Carpobrotus edulis</i> <i>Impatiens spp.</i> <i>Iris spp.</i></p>	<p>Dwarf Blue Wedgewood Bulgaria Thorndale</p>
<p>Ivy, English</p>	<p><i>Hedera helix</i></p>	<p>Arcadia</p>
<p>Jasmine, Asian Juniper</p>	<p><i>Trachelospermum asiaticum</i> <i>Juniperus spp.</i></p>	<p>Armstrong Bar Harbor Blue Chip Blue Pacific Blue Rug Blue Star Broadmoor Buffalo Calgary Carpet Emerald Sea Emerald Spreader Endora Compacta Fruitlandi Green Gold Tip Hetzi Hughes</p>
<p></p>	<p><i>Juniperus horizontalis</i></p>	<p>Armstrong Bar Harbor Blue Chip Blue Pacific Blue Rug Blue Star Broadmoor Buffalo Calgary Carpet Emerald Sea Emerald Spreader Endora Compacta Fruitlandi Green Gold Tip Hetzi Hughes</p>

		Manhattan Blue Parsoni Pfitzeriana Plumosa Prince of Wales Procumbens Dwarf San Jose Sargent Blue Sargent Green Scandia Scopulorum Moonglow Scopulorum Skyrocket Spartan Tamariscifolia Weberi Youngstown Yukon Belle
King Palm Laurel, Australian Laurel, Mountain Leucothoe Ligustrum, Japanese Lily, African	<i>Juniperus chinensis</i> <i>Juniperus Sabina</i> <i>Archontophoenix cunninghamiana</i> <i>Pittosporum tobira</i> <i>Kalmia latifolia</i> <i>Leucothoe fontanesiana</i> <i>Ligustrum japonicum</i> <i>Agapanthus africanus</i>	Albus Peter Pan
Lily, African Blue Lily of the Valley Lilyturf	<i>Pieris japonica</i> <i>Liriope muscari</i>	Mt. Fire Evergreen Giant Lilac Beauty Majestic Monroe White Variegata
Liriope, Green Maple, Japanese Maple, Norway Maple, Red Maple, Silver Maple, Sugar Marigold	<i>Liriope spicata</i> <i>Acer japonicum</i> <i>Acer platanoides</i> <i>Acer rubrum</i> <i>Acer saccharinum</i> <i>Acer saccharum</i> <i>Tagetes patula</i>	Honeycomb Variegata Wheeler's Dwarf Golden Snowflake Double White
Mock Orange*	<i>Philadelphus spp.</i>	Sunnyside Faurei Langer Muskogee Standard Pink
Mondo Grass Moss Rose Myrtle, Crape	<i>Ophiopogon japonicus</i> <i>Portulaca grandiflora</i> <i>Lagerstroemia indica</i>	Compacta Nana
Myrtle, Wax Nandina	<i>Myrica cerifera</i> <i>Nandina domestica</i>	
Narcissus Oak, Laurel Oak, Pin Oak, Red Oak, Southern	<i>Narcissus spp.</i> <i>Quercus laurifolia</i> <i>Quercus palustris</i> <i>Quercus rubra</i> <i>Quercus virginiana</i>	

Oak, Willow Oleander	<i>Quercus phellos</i> <i>Nerium oleander</i>	Hardy Red Petite Pink Sister Agnes Whirligig
Osteospermum Pachysandra Palm, Bangalow Pampas Grass Pansy Paper Flower Peach*	<i>Osteospermum fruticosum</i> <i>Pachysandra terminalis</i> <i>Cortaderia selloana</i> <i>Viola x wittrockiana</i> <i>Bougainvillea glabra</i> <i>Prunus persica</i>	Barbara Karst
Periwinkle, Dwarf Petunia Photinia, Red Tip Pieris Pine, Australian Pine, Japanese Black Pine, Loblolly Pine, Longleaf Pine, Mugho Pine, Scotch Pine, Slash Pine, Swiss Mt. Pine, Virginia Pine, White Pineapple, Guava Pittosporum, Japan Potentilla	<i>Vinca minor</i> <i>Petunia X hybrida</i> <i>Photinia X fraseri</i> <i>Pieris japonica</i> <i>Pinus nigra</i> <i>Pinus thunbergiana</i> <i>Pinus taeda</i> <i>Pinus palustris</i> <i>Pinus mugho</i> <i>Pinus sylvestris</i> <i>Pinus elliotii</i> <i>Pinus mugo</i> <i>Pinus virginiana</i> <i>Pinus strobes</i> <i>Feijoa fellowiana</i>	Picoti
Privet	<i>Potentilla nepalensis</i> <i>Potentilla fruticosa</i> <i>Ligustrum japonicum</i>	Abbotswood Golden Vicary Regal Texanum Wax Yellow Tipped
Privet, Glossy Pyracantha	<i>Ligustrum lucidum</i> <i>Pyracantha koidzumii</i>	Gnome Lalandei Victory
Queen Palm Quince, Japanese* Rhododendron	<i>Arecastrum rammanzoffianum</i> <i>Rhododendron</i> spp.	Album Cunningham White Fashion Hardy PJM Purple Gem Silvery Pink
Rhododendron, Carolina Rhododendron, Catawba Rockcress Rhodia Max (Rosebay) Rose Rosemary* Rosemary, Bog Salvia Sedum	<i>Rhododendron carolinianum</i> <i>Rhododendron catawbiense</i> <i>Arabis</i> spp. <i>Rhododendron maximum</i> <i>Rosa banksiae</i> <i>Rosmarinus officinalis</i> <i>Andromeda polifolia</i> <i>Salvia farinacea</i> <i>Sedum spurium</i>	Snowcap Luta Nana Rhea Dragon Blood Red

<p>Snapdragon Sourwood Spiraea Spiraea</p> <p>Spiraea, Garland Spruce, Black Hills Spruce, Colorado Blue Spruce, Norway Spruce, White Sweetflag, Grassyy-Le Sweetgum Sycamore Tree Fern (Tiki Fern) Trumpet Flower, Evening Tulip Verbana, Shrub Vervain Viburnum</p> <p>Vinca (Periwinkle) Windmill Palm Xylosma Yarrow Yaupon Yew</p>	<p><i>Antirrhinum</i> spp. <i>Oxydendrum arboretum</i> <i>Astilbe X arendsii</i> <i>Spirea</i> spp.</p> <p><i>Spirea</i> spp. <i>Picea glauca</i> <i>Picea pungens</i> <i>Picea abies</i> <i>Picea glauca</i></p> <p><i>Liquidambar styraciflua</i> <i>Plantanus occidentalis</i> <i>Asparagus virgatus</i> <i>Gelsemium sempervirens</i> <i>Tulipa</i> spp. <i>Lantana sellowiana</i> <i>Verbena</i> spp. <i>Viburnum</i> spp.</p> <p><i>Vinca minor</i> <i>Trachycarpus fortunei</i> <i>Xylosma congestum</i> <i>Achillea</i> spp. <i>Ilex vomitoria</i> <i>Taxus cuspidate</i> <i>Taxus X media</i></p>	<p>Red Carpet Yellow</p> <p>Fanail Anthony Waterer Red Dolchica Froebeli Pink Goldenflame Red Snowmound White Van Houtte White</p> <p>Conica</p> <p>Apeldoorn</p> <p>St. Paul American Cranberry Bush Arrowood Common Snowball European Cranberry Bush Linden Mohican Wright</p> <p>Dwarf Denisformis</p>
<p>* Ornamental species only. Do not use on food producing trees and ornamentals.</p>		

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container in a locked storage area out of reach of children and domestic animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

(Alternate wording for household use:
If empty: Do not reuse this container. Place in trash or offer for recycling if available.
If partly filled: Call your local solid waste agency for disposal instructions. Never place any unused product down any indoor or outdoor drain or toilet.

CONDITIONS OF SALE AND WARRANTY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying and using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The DIRECTIONS FOR USE of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Lawn and turf injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of Control Solutions, Inc. or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

Control Solutions, Inc. warrants that this product conforms to the chemical description on the label, and is reasonably fit for the purposes set forth in the DIRECTIONS FOR USE, when it is used in accordance with such directions, subject to the inherent risks mentioned above.

TO THE EXTENT PERMITTED BY LAW, CONTROL SOLUTIONS, INC. NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESSED OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. THIS WARRANTY EXTENDS TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNING, OR CAUTIONS. BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. CONTROL SOLUTIONS, INC. and the seller offer this product, and the Buyer and User accept it, subject to the foregoing CONDITIONS OF SALE AND WARRANTY.