

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460



OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

October 8, 2014

Ms. Jane Miller
Agent for Control Solutions, Inc.
c/o Biologic Consulting, Inc.
115 Obtuse Hill Road
Brookfield, CT 06804

Subject: Label Amendment – Adding weed species and specific ornamental plants
Product Name: Quali-Pro Prodoxaben 3.7 SC
EPA Registration Number: 53883-325
Application Date: May 27, 2014
Decision Number: 492296

Dear Ms. Miller:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). If you have any questions, please contact Beth Benbow by phone at 703-347-8072, or via email at Benbow.bethany@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Kathryn V. Montague". The signature is written in a cursive style with a long, sweeping horizontal line extending from the end of the name.

Kathryn V. Montague, Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Attachment

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by " / " denotes and/or options.

Label amendment 09 29 2014

QUALI-PRO[®] PRODOXABEN 3.7 SC

A SELECTIVE COMBINATION PREEMERGENCE HERBICIDE FOR THE CONTROL OF GRASS AND BROADLEAF WEEDS IN ESTABLISHED TURFGRASSES (EXCLUDING GOLF COURSE PUTTING GREENS), LAWNS AND ORNAMENTALS, CONIFER AND HARDWOOD SEEDLING NURSERIES, NON-CROP AREAS INCLUDING RIGHTS-OF WAY FOR TRANSPORTATION SYSTEMS AND UTILITIES (INCLUDING ROADWAYS, ROADSIDES, RAILWAYS AND EQUIPMENT YARDS), FACILITIES INCLUDING SUBSTATIONS, TANK FARMS, PUMPING STATIONS, PARKING AND STORAGE AREAS, AND UNGRAZED FENCE ROWS, AND CHRISTMAS TREE FARMS.

ACTIVE INGREDIENTS:

	% BY WT
Prodiamine*	23.3%
Isoxaben**	15.5%
OTHER INGREDIENTS:	<u>61.2%</u>
TOTAL:	100.0%

* CAS No. 29091-21-2

** CAS No. 82558-50-7

Contains 2.2 pounds prodiamine per gallon and 1.5 pounds isoxaben per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

[See attached labeling for directions for use and additional precautionary statements.]

Manufactured for:


Control Solutions, Inc.
5903 Genoa Red Bluff
Pasadena, TX 77507

EPA Reg. No. 53883-325

EPA Est. No. xxx-xx-xxx

Letter(s) in lot number correspond(s) to superscript in EPA Est. No.

Net Contents: xxxx gal (xxxx L)



NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

FIRST AID	
IF SWALLOWED:	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious or convulsing person.
IF IN EYES:	Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF INHALED:	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
IF ON SKIN OR CLOTHING:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact X-XXX-XXX-XXX for emergency medical treatment information.	

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

May be harmful if swallowed, absorbed through skin or inhaled. Avoid breathing vapor. Avoid contact with skin, eyes, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

WPS USES

Applicators, mixers, loaders, and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) - agricultural plant uses are covered - must wear:

- Long-sleeved shirt and long pants
- Chemical Resistant gloves.
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided with all of the PPE specified above for applicators and other handlers, and

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- After handling this product immediately remove PPE, wash yourself thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limit of solubility, this product is not toxic to fish. However, at concentrations substantially above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Drift may result in reduced germination or emergence of non-target plants adjacent to treated area. Do not contaminate water when disposing of equipment wash water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170.

This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves such as butyl rubber \geq 14 mils, or natural rubber \geq 14 mils, or neoprene rubber \geq 14 mils or, nitrile rubber \geq 14 mils
- Shoes plus socks

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

NON-AGRICULTURAL USE REQUIREMENTS

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forest or nurseries.

Applicators and handlers should wear:

- Long-sleeved shirt
- Long pants
- Chemical-resistant gloves
- Shoes with socks
- Protective eyewear.

Thoroughly wash face and hands with soap and water after exposure and before eating or smoking. Launder exposed clothing before reuse. Do not allow reentry to treated areas until sprays have dried.

PRODUCT INFORMATION

Formulation and Mode of Action

Quali-Pro[®] Prodoxaben 3.7 SC is a liquid herbicide containing the active ingredient, prodiamine, a member of the dinitroaniline family of herbicides which inhibits mitotic processes in plants, and isoxaben, which has a cellulose biosynthesis inhibitor (CBI) mode of action. The actives control susceptible weeds by preventing growth and development of newly germinated seeds. Each gallon contains 3.7 pounds active ingredient (2.2 pounds prodiamine per gallon plus 1.5 pounds isoxaben per gallon). Quali-Pro[®] Prodoxaben 3.7 SC is used for pre-emergent control of grasses and broadleaf weeds. Weed control is most effective when Quali-Pro[®] Prodoxaben 3.7 SC is activated by at least 0.5 inches of rainfall or irrigation.

Uses and Application

Apply Quali-Pro[®] Prodoxaben 3.7 SC for selective pre-emergence control of many grass and broadleaf weeds in the following areas:

- Established turfgrasses (excluding golf course putting greens)
- Lawns
- Container, field-grown and landscape ornamentals
- Conifer and hardwood seedling nurseries
- Non-crop areas including managed rights-of-way for transportation systems and utilities (such as roadways, roadsides, railways, and equipment yards)
- Non-crop facilities including substations, tank farms, pumping stations, parking and storage areas
- Non-grazed fence rows
- Christmas tree farms

Apply Quali-Pro[®] Prodoxaben 3.7 SC in a minimum of 20 gallons/acre (0.5 gallons /1,000 sq ft) [of carrier (water and or fluid fertilizer)] using a calibrated, low-pressure sprayer with 50-mesh or coarser screens. A broadcast boom or handheld wand designed for herbicide or insecticide application will provide the best results. Select nozzle pressure and flow rate to provide complete coverage.

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

New Plantings, Replanting and Rotational Plantings

Rotate nursery, landscape, or non-cropped land areas treated with Quali-Pro® Prodoxaben 3.7 SC only to ornamental species listed on this label for 1 (one) year following application unless the following test has shown species safety:

- Before planting a species not listed on this label, sow several test strips of an indicator plant such as wheat, sorghum or corn into the treated area. If the indicator plants germinate and grow normally to a height of 12 inches with normal root development, it is safe to plant.

In areas disturbed by new plantings or replanting of labeled species, it may be necessary to retreat exposed soil to maintain satisfactory weed control, but do not apply more than 87 fl. oz. of product per acre per year.

MIXING QUALI-PRO® PRODOXABEN 3.7 SC

When Used Alone

Mix Quali-Pro® Prodoxaben 3.7 SC thoroughly in the spray tank to ensure uniform application. Steps to follow when mixing Quali-Pro® Prodoxaben 3.7 SC include:

1. Filling the spray tank 1/4 full with clean water only;
2. Starting the agitation system and checking to ensure it is working properly;
3. Maintaining vigorous agitation in the spray tank before and during the application to ensure a well-mixed spray suspension;
4. Adding a spray colorant with Quali-Pro® Prodoxaben 3.7 SC to mark treated areas and improve application accuracy by minimizing swath skips and overlaps; and
5. After use, thoroughly clean the sprayer by flushing the system with water containing a detergent.

Refer to the **PESTICIDE DISPOSAL** section of this label for waste disposal. Do not allow spray suspension to dry in the tank. [If Quali-Pro® Prodoxaben 3.7 SC was mixed with fertilizer instead of water in the spray tank, the fertilizer aids in re-suspension of Quali-Pro® Prodoxaben 3.7 SC if agitation is disrupted.] Use the entire tank before stopping agitation.

When Used in Tank Mixtures

Quali-Pro® Prodoxaben 3.7 SC can be applied in tank-mixtures with certain other EPA-registered herbicides to provide a broader spectrum of weed control and/or to control emerged weeds. Refer to the specific directions for use for tank-mix partner[s], and refer to the label[s] of the individual tank-mix partner[s] for use rate, application timing, weeds controlled, and specific precautions and/or restrictions. Tank mixes are permitted only in states where the tank-mix partner[s] are registered for the application site and the turf and ornamental species listed. When using Quali-Pro® Prodoxaben 3.7 SC in a tank mixture with other pesticides, follow restrictions and precautions on the labels of the products used.

When mixing Quali-Pro® Prodoxaben 3.7 SC with other components (carrier and partner pesticide products), allow products to completely dissolve between steps and be sure to maintain agitation throughout mixing and application. Add other components to the spray tank in the following order:

1. Products packaged in water-soluble bags. Agitate the tank mixture. Allow the water-soluble bags to completely dissolve and the products to disperse before adding any other tank-mix partners.
2. Water-dispersible granules (WDG or WG formulations) and wettable powders (WP formulations). Add wettable powders to the tank as agitation continues. Allow the product to disperse completely before other products are added.
3. Spray adjuvants and spray markers. Read the adjuvant's label first and use only those adjuvants approved for application to turf and ornamentals.

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

4. Quali-Pro® Prodoxaben 3.7 SC, other flowable liquids (FL) or suspension concentrates (SC) must be added next.
5. Add emulsifiable concentrates (EC) last.

Compatibility Testing

Before mixing Quali-Pro® Prodoxaben 3.7 SC with other pesticides in the spray tank, test for compatibility by mixing all components (carrier and pesticide products) in an appropriate container in proportionate quantities. For example, 1 quart would be 1/100 the volume of a 25 gallons/acre spray rate. [At 21 fl. oz./acre, the Quali-Pro® Prodoxaben 3.7 SC rate would be approximately 1 teaspoon to a quart of water (see **Table 1**).]

[Table 1: Amount of tank mix component to add to one quart of spray carrier to test various tank mixes. Proportions are representative of a larger volume of 25 gallons/acre]

Component Formulations	Rate Per		Level Teaspoons
	Acre	1,000 sq ft	
Quali-Pro Prodoxaben 3.7 SC	21.0 fl. Oz.	0.5 fl. Oz.	1.0
[Dry Tank-Mix Partners	1.0 lb	0.4 oz	1.5]
[Liquid Tank-Mix Partners	16 fl. Oz.	0.4 fl. Oz.	0.5]

If components do not ball-up or form flakes, sludge, gels, oily films, or layers, then the mixture is compatible. Let the mixture stand for 15 minutes. Incompatibility will usually occur within 5 minutes after mixing. If components are not compatible, use a compatibility agent and rerun the test to determine if the mixture is suitable. If the components are still not compatible, do not tank mix.

Follow all applicable directions, restrictions, and precautions on the labels of EPA-registered tank-mix partners.

PRODUCT USE RESTRICTIONS

- Do not graze or feed livestock forage cut from areas treated with Quali-Pro® Prodoxaben 3.7 SC.
- Do not apply Quali-Pro® Prodoxaben 3.7 SC to plants that will be consumed for food use.
- Do not blend Quali-Pro® Prodoxaben 3.7 SC onto dry fertilizer or any other granular material.
- Do not apply to golf course putting greens.
- Do not exceed a dosage of 2.5 lb. active ingredient / acre (87 fl.oz. / acre) (1.5 lb. Prodiamine and 1.0 lb. Isoxaben) per year on any use site and do not repeat an application sooner than 60 days after a previous application.
- **Chemigation:** Do not apply this product through any type of irrigation system. Do not apply aurally.

State Specific Restrictions:

- The state of Arizona has not approved this product for use on agricultural commodities. If grown for commercial production, use on the following sites/crops is prohibited: container-grown field ornamentals, field-grown ornamentals, ornamental bulbs, and Christmas tree/conifer plantations.
- Ornamental bulbs grown for commercial production is restricted in California.
- Not for sale, use or distribution in the state of New York.

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

Treatment of Turfgrass or Ornamental Species Not Listed on the Label for Quali-Pro® Prodoxaben 3.7 SC:

Users who wish to use Quali-Pro® Prodoxaben 3.7 SC are limited to non-food, ornamental plant species located strictly in labeled use sites. Although this label contains a large number of ornamental species, it is not possible to include all of the ornamental plants that may be encountered in a nursery or landscape setting[s]. On a plant species not listed on this label the user may determine the suitability of such use by treating a small area or number of these plants at a specified rate. Prior to treatment of larger areas, the treated area or plants should be observed for any sign of herbicidal injury during the 30 to 60 days of typical growing conditions.

Quali-Pro® Prodoxaben 3.7 SC use rates and A.I. load simplified:

- 1 fl. oz. per 1,000 sq.ft. or 43.5 fl. oz. per acre is equivalent to 0.75 lbs. of Prodiamine and 0.5 lb. of Isoxaben per acre.
- 2.0 fl. oz. per 1,000 sq.ft. or 87 fl. oz. per acre is equivalent to 1.5 lbs. of Prodiamine and 1.0 lb. of Isoxaben per acre.

WEEDS CONTROLLED OR SUPPRESSED IN ESTABLISHED TURF AND ORNAMENTALS

When used as directed in this label for established turf and container, field grown and landscape ornamental plantings including Christmas tree farms, Quali-Pro® Prodoxaben 3.7 SC will control or suppress the grass and broadleaf weeds listed in **Table 2**.

Table 2. Weeds controlled or suppressed by Quali-Pro® Prodoxaben 3.7 SC [in Established Turf, Container, Field-grown and Landscape Ornamental Plantings Including Christmas Tree Farms]

Common Name	Scientific Name	Rates		
		C=Control, S=Suppression		
		1oz	1.5oz	2oz
Broadleaf Weeds / Dicots				
American Burnweed	<i>Erechtites hieraciifolia</i>	[C ¹	C ¹	C ¹
Annual Bursage	<i>Ambrosia acanthicarpa</i>	C	C	C
Annual Sowthistle	<i>Sonchus oleraceus</i>	C	C	C
Black Medic	<i>Medicago lupulina</i>	C	C	C
Black Mustard	<i>Brassica nigra</i>	C	C	C
Black Nightshade	<i>Solanum nigrum</i>	C	C	C
Blackleaved Fleabane	<i>Conyza bonariensis</i>	C	C	C
Bracted Plantain	<i>Plantago aristata</i>	C	C	C
Bristly Oxtongue	<i>Picris echioides</i>	C	C	C
Broadleaf Plantain	<i>Plantago major</i>	C	C	C
Buckhorn Plantain	<i>Plantago lanceolata</i>	C	C	C
Burning Nettle	<i>Urtica urens</i>	C	C	C
Burningbush	<i>Kochia scoparia</i>	C	C	C

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

Burningbush	<i>Bassia scoparia</i>	C	C	C
Buttercup Oxalis	<i>Oxalis pes-caprae</i>	C	C	C
California Burclover	<i>Medicago polymorpha</i>	C	C	C
Chamberbitter	<i>Phyllanthus urinaria</i>	C	C	C
Canadian Horseweed	<i>Conyza canadensis</i>	C	C	C
Carolina Geranium	<i>Geranium carolinianum</i>	C	C	C
Carpetweed	<i>Mollugo verticillata</i>	C	C	C
Coast Fiddleneck	<i>Amsinckia menziesii var. intermedia</i>	C	C	C
Common Chickweed ¹	<i>Stellaria media</i>	C	C	C
Common Groundsel	<i>Senecio vulgaris</i>	C	C	C
Common Lambsquarters	<i>Chenopodium album</i>	C	C	C
Common Mallow	<i>Malva neglecta</i>	C	C	C
Common Pokeweed	<i>Phytolacca americana</i>	C	C	C
Common Purslane	<i>Portulaca oleracea</i>	C	C	C
Common Ragweed	<i>Ambrosia artemisiifolia</i>	C	C	C
Common Vetch	<i>Vicia sativa</i>	C	C	C
Creeping Woodsorrel	<i>Oxalis corniculata</i>	C	C	C
Curly Dock	<i>Rumex crispus</i>	C	C	C
Dandelion	<i>Taraxacum officinale</i>	C	C	C
Dogfennel	<i>Eupatorium capillifolium</i>	C	C	C
Dwarf Fleabane	<i>Conyza ramosissima</i>	C	C	C
Evening Primrose	<i>Oenothera spp.</i>	S ²	S ²	S ²
False Daisy	<i>Eclipta prostrata</i>	C	C	C
Field Bindweed	<i>Convolvulus arvensis</i>	C	C	C
Field Burweed	<i>Soliva sessilis</i>	C	C	C
Florida Betony	<i>Stachys flordans</i>	C	C	C
Florida Pusley	<i>Richardia scabra</i>	C	C	C
Florida Tasselflower	<i>Emilia fosbergii</i>	C	C	C
Fringed Redmaids	<i>Calandrinia ciliata</i>	C	C	C
Green Tansymustard	<i>Descurainia pinnata spp. brachycarpa</i>	C	C	C
Hairy Bittercress	<i>Cardamine hirsuta</i>	C	C	C
Hairy Crabweed	<i>Fatoua vilosa</i>	C	C	C
Hairy Galinsoga	<i>Galinsoga quadriradiata</i>	C	C	C
Heath Aster	<i>Symphotrichum ericoides</i>	C	C	C
Henbit ¹	<i>Lamium amplexicaule</i>	S	S	S
Honeyvine Milkweed	<i>Cynanchum laeve</i>	C	C	C
Hyssop Spurge	<i>Chamaesyce hyssopifolia</i>	C	C	C

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

Indian Mustard	<i>Brassica juncea</i>	C	C	C
Ivyleaf Morning-Glory	<i>Ipomoea hederacea</i>	C	C	C
Jimsonweed	<i>Datura stramonium</i>	C	C	C
Knotweed ¹	<i>Polygonum spp.</i>	C	C	C
Ladysthumb	<i>Polygonum persicaria</i>	C	C	C
Lanceleaf Groundcherry	<i>Physalis angulata</i>	C	C	C
Little Mallow	<i>Malva parviflora</i>	S	S	S
Little Western Bittercress	<i>Cardamine oligosperma</i>	C	C	C
London Rocket	<i>Sisymbrium irio</i>	C	C	C
Long-stalked Phyllanthus	<i>Phyllanthus tenellus</i>	C	C	C
Marsh Parsley	<i>Cyclosporum leptophyllum</i>	C	C	C
Mexican Fireplant	<i>Poinsettia heterophylla</i>	C	C	C
Mousear Chickweed	<i>Cerastium fontanum spp. vulgare</i>	C	C	C
Mousear Chickweed	<i>Cerastium vulgatum</i>	C	C	C
Nettle-Leaved Goosefoot	<i>Chenopodium murale</i>	C	C	C
Northern Willowherb	<i>Epilobium ciliatum</i>	C	C	C
Panicle Willoweed	<i>Epilobium brachycarpum</i>	C	C	C
Pennsylvania Smartweed	<i>Polygonum pennsylvanicum</i>	C	C	C
Pennywort	<i>Hydrocotyle spp.</i>	C	C	C
Persian Speedwell	<i>Veronica persica</i>	C	C	C
Petty Spurge	<i>Euphorbia peplus</i>	C	C	C
Pigweed	<i>Amaranthus spp.</i>	C	C	C
Pillpod sandmat	<i>Chamaesyce hirta</i>	C	C	C
Pineappleweed	<i>Matricaria discoidea</i>	C	C	C
Prickly Lettuce	<i>Lactuca serriola</i>	S	S	S
Prickly Sida	<i>Sida spinosa</i>	C	C	C
Prostrate Knotweed	<i>Polygonum aviculare</i>	C	C	C
Prostrate Pigweed	<i>Amaranthus blitum</i>	S	S	S
Prostrate Spurge	<i>Chamaesyce humistrata</i>	C	C	C
Purple Cudweed	<i>Gnaphalium purpureum</i>	C	C	C
Purslane Speedwell	<i>Veronica peregrina</i>	C	C	C
Rattail Fescue	<i>Vulpia myuros</i>	C	C	C
Redroot Pigweed	<i>Amaranthus retroflexus</i>	C	C	C
Redstem Filaree	<i>Erodium cicutarium</i>	C	C	C
Russian Thistle	<i>Salsola tragus</i>	C	C	C
Santa Maria Feverfew	<i>Parthenium hysterophorus</i>	C	C	C
Scarlet Pimpernel	<i>Anagallis arvensis</i>	C	C	C

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

Shepherdspurse ¹	<i>Capsella bursa-pastoris</i>	C	C	C
Silversheath Knotweed	<i>Polygonum argyrocoleon</i>	C	C	C
Slender Aster	<i>Symphotrichum divaricatum</i>	C	C	C
Slender Plantain	<i>Plantago heterophylla</i>	C	C	C
Southern Waterbuttons	<i>Cotula australis</i>	C	C	C
Spanish needles	<i>Bidens alba</i>	C	C	C
Spiny Sowthistle	<i>Sonchus asper</i>	C	C	C
Spotted Spurge	<i>Chamaesyce maculata</i>	C	C	C
Spotted Spurge	<i>Euphorbia maculata</i>	C	C	C
Stinking Chamomile	<i>Anthemis cotula</i>	C	C	C
Sunflower	<i>Helianthus spp.</i>	C	C	C
Swinecress	<i>Coronopus didymus</i>	C	C	C
Tall Morning-Glory	<i>Ipomoea purpurea</i>	C	C	C
Telegraphplant	<i>Heterotheca grandiflora</i>	C	C	C
Thistle Musk	<i>Carduus nutans</i>	C	C	C
Thymeleaf Speedwell	<i>Veronica serpyllifolia</i>	C	C	C
Turkey Mullein	<i>Croton setigerus</i>	C	C	C
Venice Mallow	<i>Hibiscus trionum</i>	C	C	C
Virginia Pepperweed	<i>Lepidium virginicum</i>	C	C	C
Virginia Winged Rockcress	<i>Sibara virginica</i>	C	C	C
White Clover	<i>Trifolium repens</i>	C	C	C
White Eclipta	<i>Eclipta alba</i>	C	C	C
Whitestem Filaree	<i>Erodium moschatum</i>	C	C	C
Wild Carrot	<i>Daucus carota</i>	C	C	C
Wild Mustard	<i>Sinapis arvensis</i>	C	C	C
Wild Radish	<i>Raphanus raphanistrum</i>	C	C	C
Yellow Sweetclover	<i>Melilotus officinalis</i>	C	C	C
Yellow Woodsorrel	<i>Oxalis stricta</i>	S	S	S]
Grass Weeds and Monocots				
Annual Bluegrass	<i>Poa annua</i>	C	C	C
Baldwin's Flatsedge	<i>Cyperus croceus</i>	C	C	C
Barnyardgrass	<i>Echinochloa crus-galli</i>	C	C	C
Broadleaf Signalgrass	<i>Brachiaria platyphylla</i>	C	C	C
Browntop Panicum	<i>Panicum fasciculatum</i>	C	C	C
Crowfootgrass	<i>Dactyloctenium aegypticum</i>	C	C	C
Fall Panicum	<i>Panicum dichotomiflorum</i>	C	C	C
Giant Foxtail	<i>Setaria faberi</i>	C	C	C

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

Goosegrass ³	<i>Eleusine indica</i>	C	C	C
Green Foxtail	<i>Setaria viridis</i>	C	C	C
Itchgrass	<i>Rottboellia cochinchinensis</i>	C	C	C
Japanese Stiltgrass	<i>Microstegium viminium</i>	C	C	C
Johnsongrass	<i>Sorghum halepense</i>	S	S	S
Jungle Rice	<i>Echinochloa colonum</i>	C	C	C
Large Crabgrass	<i>Digitaria sanguinalis</i>	C	C	C
Lovegrass	<i>Eragrostis spp.</i>	C	C	C
Nakedstem Dewflower	<i>Murdannia nudiflora</i>	C	C	C
Red Sorrell	<i>Rumex acetosella</i>	C	C	C
Rescuegrass ²	<i>Bromus catharticus</i>	C	C	C
Rice Flatsedge	<i>Cyperus iria</i>	C	C	C
Smooth Crabgrass	<i>Digitaria ischaemum</i>	C	C	C
Sprangletop	<i>Leptochloa spp.</i>	C	C	C
Texas Panicum	<i>Panicum texanum</i>	C	C	C
Wandering Jew ^{2†}	<i>Tradescantia zebrina</i>	C	C	C
Witchgrass	<i>Panicum capillare</i>	C	C	C
Woolly Cupgrass	<i>Eriochloa villosa</i>	C	C	C
Yellow Foxtail	<i>Setaria lutescens</i>	C	C	C

† Not for use in California

c, 1 Controlled at the indicated rate. To control this weed, apply Quali-Pro[®] Prodoxaben 3.7 SC in late summer, fall, or winter before weeds germinate.

s, 2 Suppression only at the indicated rate.

3 In many areas a single application of 43.5-87.0 fl.oz./acre of Quali-Pro[®] Prodoxaben 3.7 SC will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, weed control will be most effective by making an initial application of 43.5 fl.oz./acre followed by a second application 60-90 days later.

Note to PM: This product can control or suppress the list of weeds in Table 2 at a range of acceptable use rates. The final artwork label will only have one use rate for control or suppression for each listed weed indicated by a “C” or an “S” respectively.

SPECIFIC USE DIRECTIONS - ESTABLISHED TURF

Product Information

Quali-Pro[®] Prodoxaben 3.7 SC is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turfgrasses including golf courses (excluding putting greens), lawns and sod nurseries. The maximum amount of Quali-Pro[®] Prodoxaben 3.7 SC that can be applied per year is given for each turfgrass species in **Table 4** of this label.

USE RESTRICTIONS - Turfgrass: Golf Courses (excluding golf course putting greens), Lawns, and Sod Nurseries

- Do not apply Quali-Pro[®] Prodoxaben 3.7 SC to areas where dichondra, colonial bentgrass,

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.

- Do not cut (harvest) treated sod before 90 days after application. To avoid turfgrass injury, do not apply to newly set sod until the sod has rooted and exposed edges have filled in.
- **Do not apply Quali-Pro® Prodoxaben 3.7 SC to golf course putting greens.**
- Do not apply Quali-Pro® Prodoxaben 3.7 SC to turf grass grown for seed.
- Do not apply Quali-Pro® Prodoxaben 3.7 SC to turf stressed by conditions such as drought, low fertility, or pest damage.

USE PRECAUTIONS - Turfgrass: Golf Courses (excluding golf course putting greens), Lawns, and Sod Nurseries

- Disturbing the herbicide barrier with cultural practices such as disking can result in reduced weed control.
- If the depth of the creeping bentgrass root system becomes shallow and root tips contact proflaminate-treated soil, new root formation can be inhibited. Mowing height can affect the depth of a plants root system. To avoid this do not apply Quali-Pro® Prodoxaben 3.7 SC to creeping bentgrass less than 0.5 inches in height.

For optimum weed control, activate Quali-Pro® Prodoxaben 3.7 SC with at least 0.5 inches of rainfall or irrigation before weed seeds germinate and within 14 days following application. See the map below for approximate crabgrass seed germination dates.

Crabgrass Seed Germination Dates



Timing and Method of Application to Turfgrass

Apply Quali-Pro® Prodoxaben 3.7 SC as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made before target weeds germinate because Quali-Pro® Prodoxaben 3.7 SC will not control weeds that have already emerged. Apply Quali-Pro® Prodoxaben 3.7 SC in a minimum of 20 gallons/acre (0.5 gallons/1,000 sq ft) of carrier (water and/or fluid fertilizer) using a calibrated, low-pressure sprayer with 50-mesh or coarser screens. A broadcast boom or handheld wand designed for herbicide or insecticide application will provide the best results. Select nozzle pressure and gallonage to provide complete coverage.

The amount of Quali-Pro® Prodoxaben 3.7 SC to apply depends upon the length of residual weed control desired (the higher the application rate, the longer the control), the turf species, and the maximum amount which can be applied to the turf species per calendar year. Specific information is provided in **Table 3** and **Table 4**.

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

Table 3. Length of Crabgrass Control* With Quali-Pro® Prodoxaben 3.7 SC

Note: Length of control varies by region. This is an average:

0.5 fl. oz. per 1,000 sq.ft. or 21.75 fl. oz./acre - up to 3 months control of crabgrass.

0.75 fl. oz. per 1,000 sq.ft or 32.6 fl. oz./acre - up to 4 months control of crabgrass.

1.0 Fl. oz. per 1,000 sq.ft or 43.5 fl.oz./acre - up to 6 months control of crabgrass.

1.3 fl. oz. per 1,000 sq.ft. or 56.5 fl. oz./acre- up to 8 months control of crabgrass.

Dosage

Quali-Pro® Prodoxaben 3.7 SC can be safely applied to the turfgrass species listed in **Table 4**. Do not exceed the highest rate listed for each species in a calendar year as injury can occur.

Table 4. Maximum Application Rate of Quali-Pro® Prodoxaben 3.7 SC per Calendar Year by Turf Species

Turf Species	Fl. Oz. Product/Acre ¹	fl. Oz. Product/1,000 sq ft ¹
Bermudagrass ² Bahigrass Centipedegrass Kikuyugrass Seashore Paspalum St. Augustinegrass ³ Tall Fescue (including turf-type) Zoysiagrass	38-87	0.87-2.0
Buffalograss Kentucky Bluegrass Perennial Ryegrass	18-54	0.41-1.2
Fine Fescue	18-43	0.41-1.0
Creeping Bentgrass (0.5 inch or more in height ⁴)	18-38	0.41-0.87

¹ Quali-Pro® Prodoxaben 3.7 SC can be applied more than once per year as long as the total amount applied is not greater than the maximum application rate for each turf species. All applications must be made before weed seeds germinate.

² Quali-Pro® Prodoxaben 3.7 SC can be used on newly sprigged or plugged bermudagrass at rates not to exceed 30.7 fl. oz./acre (0.7 oz./1,000 sq ft). Newly sprigged or plugged bermudagrass stolon rooting can be temporarily inhibited.

³ Use an initial rate of 29-58 fl. oz./acre per application.

⁴ To avoid grass injury, do not apply Quali-Pro® Prodoxaben 3.7 SC to creeping bentgrass mowed at less than 0.5 inch in height.

When to Apply Quali-Pro® Prodoxaben 3.7 SC after Overseeding Turf

Injury to desirable seedlings is likely if Quali-Pro® Prodoxaben 3.7 SC is applied before the secondary roots of seedlings are in the second inch of soil, not thatch plus soil. To reduce the potential to injure overseeded turf, wait 60 days after seeding or until after the second mowing, whichever is longer, before applying Quali-Pro® Prodoxaben 3.7 SC.

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

When to Overseed After Application - All States

Quali-Pro® Prodoxaben 3.7 SC will inhibit the development of turfgrass species overseeded too soon after application. Follow rates and intervals in **Table 5** for best overseeding/reseeding results.

Table 5. Waiting Interval before Overseeding Perennial Ryegrass Following Application of Quali-Pro® Prodoxaben 3.7 SC

Amount of Quali-Pro® Prodoxaben 3.7 SC FL. Oz. Product/Acre	Interval (Months Before Overseeding)*		
	North	Transition	South
29	4	4	4
38	5	4	4
43	6	5	5
47	---	6	6
56	---	7	7
65	---	---	9
76	---	---	10
87	---	---	12

* In AZ, CA, NV, and TX the overseeding interval can be shorter in established bermudagrass that has been overseeded with perennial ryegrass. See the next section for additional information.

Poa Annu Control in Established Bermudagrass Overseeded with Perennial Ryegrass (AZ, CA, NV, and TX only): Use on golf courses (excluding golf course putting greens), lawns, and sod nurseries when overseeding with perennial rye grass. (Minimum seeding rate of 350 lbs/ace).

Table 6. Amount of Quali-Pro® Prodoxaben 3.7 SC to Apply to Bermudagrass Overseeded with Perennial Ryegrass

Amount to Apply	When to Apply	Expected Control	Use Precautions
21.7- 38 Fl.Oz./acre *	6-8 weeks before ryegrass overseeding Second application: 4-8 weeks after overseeding or when perennial ryegrass roots are in the second inch of soil	1 application for 70% or greater control of <i>Poa annua</i> Second application can enhance control	<ul style="list-style-type: none"> • Some seedling mortality and temporary reduction in root growth of new seedlings can occur. • To reduce the potential for seedling mortality, maintain a moist seedbed with light, frequent irrigation. • Make no more than 2 applications per year for this use, and do not exceed a total of 48 fl. oz./acre (1.38 pounds active ingredient/acre) per year. • Do not make a second application if any injury to the ryegrass is observed after the

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

			first application. <ul style="list-style-type: none">• Do not make a second application unless the product was first applied before overseeding.
--	--	--	--

* The amount of Quali-Pro® Prodoxaben 3.7 SC to apply depends upon: the length of residual control desired (the higher the application rate, the longer the control). **Note:** The higher the rate, the greater the potential for seedling mortality.

CONTAINER, FIELD-GROWN AND LANDSCAPE ORNAMENTAL PLANTINGS INCLUDING CHRISTMAS TREE FARMS

Product Information

Apply Quali-Pro® Prodoxaben 3.7 SC to soil surfaces for preemergence and limited post-emergence control of many grass and broadleaf weeds around ornamental shrubs and trees, established perennial vegetation and wildflower plantings and Christmas tree farms. [See **Table 2** for a list of weeds controlled.]

USE RESTRICTIONS - Container, Field-grown and Landscape Ornamental Plantings and Christmas Tree Farms

- Do not apply Quali-Pro® Prodoxaben 3.7 SC to seedbeds, or seedling transplant beds. Only apply to established plantings.
- Do not apply to unrooted liners or cuttings that have been planted in pots for the first time.
- Do not apply to pots less than six inches wide
- Do not apply to ground covers until they are established and well rooted
- Do not apply to bedding plants or areas where bedding plants will be planted or transplanted within one year after application.
- Do not apply Quali-Pro Prodoxaben 3.7 SC to plants inside enclosed structures.
- Do not cover plants treated with Quali-Pro Prodoxaben 3.7 SC until 21 days have passed after application.
- Do not apply Quali-Pro Prodoxaben 3.7 SC to newly transplanted ornamentals, nursery stock, or ornamental bulbs. Delay application until soil has settled and roots have formed.
- Do not apply after emergence of ornamental bulbs.

USE PRECAUTIONS - Container, Field-grown and Landscape Ornamental Plantings and Christmas Tree Farms

- Disturbing the herbicide barrier with cultural practices such as disturbing containers, removal of emerged weeds, disking, harrowing, and deep raking can result in reduced weed control.
- Applications of Quali-Pro Prodoxaben 3.7 SC over the top of plants with newly forming buds may cause injury. Possible plant injury may be avoided by application as a directed spray to the soil surface beneath ornamental plants.

Timing and Method of Application

Quali-Pro® Prodoxaben 3.7 SC is primarily a preemergence herbicide and will not control most grass or broadleaf weeds post-germination. It works best when applied to soil that is free of clods, weeds and debris and is activated by at least 0.5 inches of irrigation or rainfall or shallow incorporation to a depth of 1 to 2 inches. Weed control is improved if Quali-Pro® Prodoxaben 3.7 SC is activated in the soil before weed seeds germinate and within 14 days after application. Apply Quali-Pro® Prodoxaben 3.7 SC to newly-transplanted and established ornamentals as an

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

over-the-top spray.

Post-Emergence Control of Yellow Woodsorrel (*Oxalis*) and Bittercress:

Quali-Pro® Prodoxaben 3.7 SC will control or suppress Hairy Bittercress, *Cardamine hirsuta*, and Yellow Woodsorrel, *Oxalis stricta*, seedlings from germination up through the 1st true leaf stage. Post emergent seedling mortality reaches its maximum over a period of 7-28 days depending on local conditions. Warmer conditions typically result in faster results.

Plant Sensitivity

Direct application of Quali-Pro® Prodoxaben 3.7 SC to rapidly growing tissue or buds can injure plants. In the spring when buds are rapidly growing and expanding, over-the-top application of Quali-Pro® Prodoxaben 3.7 SC can injure new growth. However, these effects are temporary. To reduce the possibility of injury, wait to apply Quali-Pro® Prodoxaben 3.7 SC over the top of newly emerged vegetation until it has hardened off, unless your experience indicates that the plant in question will not be injured by the over-the-top application. After application, immediately apply overhead irrigation to the foliage to rinse Quali-Pro® Prodoxaben 3.7 SC from plant surfaces onto soil (wetting the foliage of plants before application can improve the rinsing process).

Table 7. Amount of Quali-Pro® Prodoxaben 3.7 SC to Apply to Container, Field Grown and Landscape Ornamental Plantings and Christmas Trees Farms

Amount to Apply	When to Apply	Comments/Instructions
38-87 fl. oz./acre or 0.9-2.0 fl. oz./1,000 sq ft	In fall or spring before weeds germinate or after weeds are removed.	<ul style="list-style-type: none">• Use the higher amount for longer control.• Quali-Pro® Prodoxaben 3.7 SC can be applied more than once per year as long as the total amount of product applied does not exceed 87 fl. oz./acre (2.5 lbs. active ingredient/acre) per year.

Table 8. Equivalent Measurements for Quali-Pro® Prodoxaben 3.7 SC

Fl. Oz./Acre	Fl. oz./1,000 sq ft	Approximate Equivalent - Tablespoons/1,000 sq ft
38	0.9	1.75
43.5	1.0	2.0
87	2.0	4.0

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

Table 9. Application Sites and Instructions for Quali-Pro® Prodoxaben 3.7 SC Use on Container, Field Grown and Landscape Ornamental Plantings and Christmas Tree Farms

Site	Application Instructions
Newly-Transplanted Container or Field Nursery Stock	<ul style="list-style-type: none">• Do not apply until soil has settled around transplants and roots have formed.• Water transplants thoroughly before application.• [Apply after cuttings form roots and are established.]• [To avoid inhibition of the tissue union,] apply before budding/grafting or after buds/grafts have taken, to avoid inhibition of the tissue union.• Do not apply to pots less than six inches wide
Established Container, Field Nursery Stock, or Landscape Plants	<ul style="list-style-type: none">• Apply at any time as an over-the-top, or directed spray.
Landscape (or Ornamental) Plantings	<ul style="list-style-type: none">• Apply as an over-the-top, or as a directed spray.• Delay applications to newly transplanted ornamentals until soil has settled around transplants.
Bare Ground Application for Container Placement	<ul style="list-style-type: none">• Apply to soil (including mulch, gravel, wood chips, or other permeable base) upon which containerized ornamentals are placed.• After Quali-Pro® Prodoxaben 3.7 SC is applied, perform shallow cultivation or hand weeding only, to avoid disturbing the herbicide barrier.
In Shadehouses and Uncovered Polyhouses	<ul style="list-style-type: none">• After Quali-Pro® Prodoxaben 3.7 SC is applied, houses must remain uncovered for at least 21 days. In addition, plants must receive 2 irrigations totaling at least 0.5 inches of water before covering.
[Ornamental Bulbs (Grown for Commercial Production is prohibited in California and Arizona).	<ul style="list-style-type: none">• Quali-Pro® Prodoxaben 3.7 SC can be applied to bulbs. Apply before bulbs emerge. Do not apply after emergence.]

[Tank Mixtures for Use on Container, Field Grown and Landscape Ornamentals and Christmas Tree Farms

Tank mix Quali-Pro® Prodoxaben 3.7 SC with other registered herbicides listed on this label to provide a broader spectrum of weed control or to control emerged weeds. Tank mixes with Quali-Pro® Prodoxaben 3.7 SC are for use only in states where the tank-mix partner(s), application site and intended use pattern are registered.

Follow the label directions of the tank-mix partner(s) for application rates, timing, weeds controlled, listed ornamentals, and specific use precautions and/or restrictions. Before combining a tank-mix partner in the spray tank, test for compatibility as described in the **Compatibility Testing** section of this label.]

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

[Table 10. Tank-Mix Partners for Quali-Pro® Prodoxaben 3.7 SC Use on Container, Field Grown and Landscape Ornamental Plantings and Christmas Tree Farms]

Product	Precautions/Instructions
Galigan® 2E, (use on conifers only)	<ul style="list-style-type: none"> Mix with Quali-Pro® Prodoxaben 3.7 SC for postemergence control of certain broadleaf weeds including malva and filaree.
Princep®, MANA Parallel	<ul style="list-style-type: none"> See product labels for weed spectrum and listed ornamentals.
Glyphogan® (or other glyphosate-based products) and Finale®	<ul style="list-style-type: none"> These nonselective tank mix herbicides control many emerged annual broadleaves and grasses. Take extreme care to prevent tank mixtures with these partner products from contacting the foliage and stems of turfgrass, trees, shrubs, or other desirable vegetation because desirable vegetation could be severely injured or killed. Apply these tank mixtures as a directed spray and use a shield to prevent spray from contacting foliage of desirable plants. Following instructions on the tank-mix partner's label, delay irrigation of the treated area to allow time for the herbicide to be absorbed by weed foliage.

Ornamental Plant Sensitivity

Quali-Pro® Prodoxaben 3.7 SC will not likely harm the trees, shrubs, vines and flowers listed below in **Table 11**. Quali-Pro® Prodoxaben 3.7 SC can be applied over the top of the listed species. The species that are sensitive to Quali-Pro® Prodoxaben 3.7 SC when grown in containers are indicated.

When plants are under stress (such as heat, drought, or frost damage), some cultivars of listed plants may be sensitive to Quali-Pro® Prodoxaben 3.7 SC.

Table 11. THE PLANTS LISTED IN THIS TABLE ARE NOT LIKELY TO BE INJURED BY QUALI-PRO® PRODOXABEN 3.7 SC AT ANY OF THE LISTED APPLICATION RATES

Scientific name	Common name
<i>Abelia grandiflora</i>	Edward Goucher Glossy Abelia
<i>Abelia grandiflora</i>	Glossy Abelia
<i>Abelia grandiflora</i> †	Abelia: Sherwood
<i>Abies balsamea</i>	Balsam Fir
<i>Abies concolor</i>	White / Concolor Fir
<i>Abies spp.</i>	Fir species** (Balsam, Fraser, Noble, etc.)
<i>Abutilon hybridum</i>	Flowering Maple
<i>Acacia abyssinica</i>	Abyssinica Acacia
<i>Acacia redolens</i>	Prostrate Acacia
<i>Acacia stenophyll</i>	Shoestring Acacia
<i>Acalypha wilkesiana</i>	Copper Plant
<i>Acer ginnala</i>	Amur Maple

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Acer ginnala</i>	Flame Maple
<i>Acer palmatum</i>	Japanese Maple
<i>Acer palmatum</i>	Coral Bark Japanese Maple
<i>Acer palmatum</i>	Dwarf Japanese Maple
<i>Acer platanoides</i>	Norway Maple
<i>Acer saccharinum</i>	Silver Maple
<i>Achillea filipendulina</i>	Moonshine /fern-leaf yarrow
<i>Achillea millefolium</i>	Common Yarrow
<i>Achillea millefolium</i>	Paprika Yarrow
<i>Achillea millefolium</i>	Achillea 'Saucy Seduction'
<i>Achillea spp</i>	Yarrow
<i>Achillea spp. †</i>	Yarrow: King Edward
<i>Achillea tomentosa</i>	Wooly Yarrow
<i>Acoelorrhaphe wrightii</i>	Everglades Palm
<i>Actinidia chinensis*</i>	Kiwi*
<i>Adonidia merrilli</i>	Christmas Palm
<i>Agapanthus africanus</i>	Lily-of-the-Nile (African Lily)
<i>Agapanthus africanus</i>	Queen Anne Lily of the Nile
<i>Agapanthus orientalis†</i>	
<i>Agave americana</i>	Century Plant
<i>Akebia quintata†</i>	Five-Leaf or Chocolate Vine
<i>Albizia julibrissin</i>	Mimosa; Silk Tree
<i>Allium cernuum†</i>	Lady's Leek, Nodding Onion
<i>Alsophils australis</i>	Australian Tree Fern
<i>Amelanchier alnifolia</i>	Saskatoon Serviceberry
<i>Andromeda polifolia</i>	Bog Rosemary
<i>Anemone hybrida†</i>	Japanese Anemone
<i>Anisodonte hypomandarum</i>	Cape Mallow
<i>Aptenia cordifolia</i>	Red Apple Aptenia
<i>Aquilegia</i>	Dragon Fly Columbine
<i>Aquilegia</i>	Red & Gold Columbine
<i>Aquilegia spp. †</i>	Columbine
<i>Araucaria heterophylla</i>	Norfolk Island Pine
<i>Archontophoenix cunninghamiana</i>	King Palm
<i>Arctostaphylos densiflora</i>	Vine Hill Manzanita
<i>Arctostaphylos uva-ursi</i>	Bearberry
<i>Arctotheca calendula</i>	Cape Weed
<i>Ardisia japonica</i>	Chirimen Marlberry
<i>Argyranthemum frutescens</i>	Butterfly Argyranthemum
<i>Armeria maritima</i>	Bloodstone Sea Thrift

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Artemesia lactiflora</i>	White Mugwort
<i>Artemisia spp</i>	Silver Mound Wormwood
<i>Artemisia spp. †</i>	Wormwood; Silver Mound, Castle
<i>Asparagus</i>	
<i>Asparagus densiflorus</i>	Foxtail Fern
<i>Asparagus retrofractus</i>	Ming Asparagus Fern
<i>Asparagus variegata</i>	
<i>Aspidistra elatior</i>	Cast Iron Plant
<i>Aster</i>	Bonny Blue Aster
<i>Aster</i>	Purple Dome Aster
<i>Aster</i>	Aster
<i>Aster novae-angliae</i>	New England Aster
<i>Aster novi-belgii</i>	New York Aster
<i>Aster novi-belgii</i>	Persian Rose Dwarf Aster
<i>Aster spp. †</i>	Aster: Bonny Blue, Purple Dome
<i>Athyrium filix-femina†</i>	Lady Fern
<i>Athyrium nipponimcum</i>	Japanese Painted Fern
<i>Aucuba japonica</i>	Japanese Aucuba
<i>Aucuba japonica</i>	Goldstrike Aucuba, Goldstrike Japanese Laurel
<i>Baccharis pulularis</i>	Coyotebush
<i>Bauhinia galpinii</i>	Red Bauhinia
<i>Begonia semperflorens</i>	Ambassador White Begonia
<i>Begonia spp</i>	Fibrous Begonia
<i>Begonia spp. †</i>	Fibrous Begonia: Hardy Grandis
<i>Berberis gladwynensis</i>	William Penn Barberry
<i>Berberis gladwynensis</i>	Barberry
<i>Berberis julianae</i>	Wintergreen Barberry
<i>Berberis mentorensis</i>	Mentor Barberry
<i>Berberis thunbergii</i>	Japanese Barberry
<i>Berberis thunbergii</i>	Japanese Barberry 'Concord'
<i>Berberis thunbergii</i>	Crimson Pygmy Barberry
<i>Berberis thunbergii</i>	Golden Japanese Barberry
<i>Berberis thunbergii</i>	Rosy Glow Barberry
<i>Berberis thunbergii</i>	Redleaf Japanese Barberry
<i>Berberis thunbergii</i>	Cherry Bomb Barberry
<i>Berberis verruculosa</i>	Warty Barberry
<i>Bergenia cordifolia</i>	Heartleaf Bergenia
<i>Bergenia cordifolia†</i>	
<i>Betula nigra</i>	River Birch
<i>Betula papyrifera</i>	Paper Birch

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Betula pendula</i>	European White Birch
<i>Bidens ferulifolia</i>	Peters Gold Bidens
<i>Boltonia asteroides</i> †	Snowbank
<i>Bougainvillea spp</i>	Barbara Karst Bougainvillea
<i>Bougainvillea spp</i>	California Gold Bougainvillea
<i>Bougainvillea spp</i>	Pink Pixie Bougainvillea
<i>Bougainvillea spp</i>	Scarlet Ohara Bougainvillea
<i>Bougainvillea spp</i>	Temple Fire Bougainvillea
<i>Bougainvillea spp</i>	Texas Dawn Bougainvillea
<i>Bougainvillea spp</i>	Purple Queen Bougainvillea
<i>Bougainvillea spp</i>	Rosenka Bougainvillea
<i>Bougainvillea spp.</i> †	Bougainvillea
<i>Brachychiton populneus</i>	Bottle Tree
<i>Brachycome</i>	Swan River daisy-New amethyst
<i>Bucida buceras</i>	Black Olive
<i>Buddleia davidii</i>	Butterfly Bush
<i>Buddleia davidii</i> †	Butterfly-Bush 'Royal Red'
<i>Butia odorata</i>	Pindo Palm
<i>Buxus 'Green Velvet'</i>	Boxwood 'Green Velvet'
<i>Buxus microphylla</i>	Japanese Boxwood
<i>Buxus microphylla</i>	Green Beauty Boxwood
<i>Buxus microphylla</i>	Japanese Littleleaf Boxwood
<i>Buxus microphylla</i>	Korean Boxwood
<i>Buxus sempervirens</i>	Common Boxwood
<i>Callistemon citrinus</i>	Lemon Bottlebrush
<i>Callistemon citrinus</i>	Little John Bottlebrush
<i>Callistemon citrinus</i> †	Crimson Bottlebrush
<i>Callistemon viminalis</i>	Weeping Bottlebrush
<i>Callistepheus cinensis</i>	China Aster
<i>Calluna vulgaris</i>	Scotch Heather
<i>Calluna vulgaris</i>	Spring Torch Scotch Heather
<i>Camellia japonica</i>	Japanese Camellia
<i>Campanula carpatica</i>	Bellflower; Tussock flower
<i>Campanula carpatica</i> †	Tussock Bellflower; (White Clips)
<i>Campis X tagliabuana</i> †	Trumpet Creeper, Trumpet flower; Madame Galen
<i>Carex</i>	Variegated Carex
<i>Carex albula</i>	Frosty Curls Sedge
<i>Carpobrotus edulis</i>	Hottentot Fig (Ice Plant)
<i>Carpobrotus edulis</i>	Largeleaf Ice Plant
<i>Caryopteris clandonensis</i>	Blue Mist Bluebeard

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by "/" denotes and/or options.

Label amendment 09 29 2014

<i>Caryopteris clandonensis</i>	Dark Knight Bluebeard
<i>Cassia artemisioides</i>	Feathery Cassia
<i>Cassia eremophila</i>	Senna
<i>Catharanthus roseus</i>	Madagascar Periwinkle
<i>Ceanothus rigidus</i>	Wild Lilac
<i>Ceanothus spp</i>	Wild Lilac
<i>Ceanothus x pallidus</i>	Lilac 'Marie Bleu'
<i>Cephalotaxus harringtonia</i>	Japanese Plum Yew
<i>Cerastium tomentosum</i>	Snow in Summer
<i>Ceratonia siliqua</i>	Carob
<i>Ceratostigma plumbaginoides</i> †	
<i>Ceratostigma willmottianum</i>	Chinese Plumbago
<i>Cercis canadensis</i>	Redbud
<i>Chaenomeles japonica</i>	Orange Flowering Quince
<i>Chamaecyparis obtusa</i>	Kosteri Cypress
<i>Chamaecyparis obtusa</i>	Dwarf Hinoki Cypress
<i>Chamaecyparis obtusa</i>	Torulosa Cypress
<i>Chamaecyparis obtusa</i>	Fernspray Falsecypress
<i>Chamaecyparis obtusa</i>	Slender Hinoki False Cypress
<i>Chamaecyparis pisifera</i>	False Cypress
<i>Chamaecyparis pisifera</i>	Squarrosa Minima False Cypress
<i>Chamaecyparis pisifera</i>	Threadleaf False Cypress
<i>Chamaecyparis pisifera</i>	Sawara False Cypress
<i>Chamaecyparis pisifera</i>	Moss Falsecypress
<i>Chamaedorea cataractarum</i>	Cat Palm
<i>Chamaedorea costaricana</i>	Costa Rican Chamaedorea Palm
<i>Chamaedorea elegans</i>	Parlor Palm
<i>Chamaerops humilis</i>	European Fan Palm, Mediterranean Fan Palm
<i>Chasmanthium latifloium</i>	Northern Sea Oats
<i>Chrysanthemum maximum</i>	Shasa Daisy
<i>Chrysanthemum spp</i>	Chrysanthemum
<i>Citrus spp.*</i>	Citrus species*
<i>Clethra alnifolia</i>	Summersweet
<i>Cleyera japonica</i>	Cleyera, Japanese Cleyera
<i>Clivia miniata</i>	Kafir Lily
<i>Colonema pulchrum</i>	Pink Breath of Heaven
<i>Convolvulus cneorum</i>	Bush Morning Glory
<i>Convolvulus mauritanicus</i>	Ground Morning Glory
<i>Cordyline indivisa</i>	Blue Dracaena
<i>Coreopsis</i>	Early Sunrise Coreopsis

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Coreopsis grandiflora</i>	Coreopsis
<i>Coreopsis spp. †</i>	Coreopsis (Calliopsis): Early Sunrise, Moonbeam
<i>Coreopsis verticillata</i>	Coreopsis
<i>Cornus florida</i>	Cloud Nine Dogwood
<i>Cornus florida</i>	Flowering Dogwood
<i>Cornus kousa</i>	Kousa Dogwood
<i>Cornus alba</i>	Siberian Dogwood
<i>Cornus sericea</i>	Red Osier Dogwood
<i>Cornus sericea</i>	Yellowtwig Dogwood
<i>Cornus stolonifera</i>	American Dogwood
<i>Cortaderia selloana</i>	Pampas Grass
<i>Corylus americanus</i>	Harry Lauder's Walking Stick
<i>Cotinus coggygria</i>	Royal Purple Smoke Tree
<i>Cotinus dammeri</i>	Coral Beauty Smoke Tree
<i>Cotinus dammeri</i>	Eichholz Smoke Tree
<i>Cotoneaster adpressus</i>	Early Cotoneaster
<i>Cotoneaster apiculatus</i>	Cranberry Cotoneaster
<i>Cotoneaster buxifolius</i>	Box Leafed Cotoneaster
<i>Cotoneaster congestus</i>	Pyrenees Cotoneaster
<i>Cotoneaster dammeri</i>	Bearberry Cotoneaster
<i>Cotoneaster himalayan</i>	Himalayan Cotoneaster
<i>Cotoneaster horizontalis</i>	Rock Cotoneaster
<i>Cotoneaster microphyllus</i>	Rockspray Cotoneaster
<i>Crataegus Species</i>	Hawthorns
<i>Crataegus viridis</i>	Green Hawthorn
<i>Crocoshmia spp. †</i>	Lucifer
<i>Cryptomeria japonica</i>	Japanese Cryptomeria
<i>Cupaniopsis anacardioides</i>	Carrot Wood
<i>Cuphea hyssopifolia</i>	False or Mexican Heather
<i>Cupressocyparis leylandii</i>	Emerald Isle Leyland Cypress
<i>Cupressocyparis leylandii</i>	Naylor's Blue Leyland Cypress
<i>Cupressus arizonicus</i>	Arizona Cypress
<i>Cupressus arizonicus</i>	Blue Pyramid Arizona Cypress
<i>Cupressus sempervirens</i>	Italian Cypress
<i>Cupressus sempervirens</i>	Glauca Italian Cypress
<i>Cycas revoluta</i>	Sago Palm
<i>Cyperus albostrigatus</i>	Dwarf Umbrella Grass
<i>Cyrtomium fortunei</i>	Holly Fern
<i>Cytisus praecox</i>	Hollandia-Warminster Broom
<i>Cytisus scoparius</i>	Scotch Broom

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Cytisus spp</i>	Holandia Scotch Broom
<i>Cytisus spp</i>	Allgold Scotch Broom
<i>Dahlia</i>	Dwarf Dahlia Wendy Pink
<i>Dahlia hybrid</i>	Dwarf Dahlia
<i>Dalea greggii</i>	Trailing Indigo Bush
<i>Daphne cneorum</i>	Rose Daphne
<i>Daphne odora</i>	Fragrant Daphne
<i>Delosperma alba</i>	White Trailing Ice Plant
<i>Delosperma cooperi</i>	Ice Plant
<i>Delosperma cooperi</i>	Cooperi Pink
<i>Delosperma nubigenum</i>	Hardy Ice Plant
<i>Delphinium spp. †</i>	Larkspur; Blue Elf
<i>Descampsia caespitosa</i>	descampsia
<i>Deutzia crenata</i>	Nakiana-Dwarf Deutzia
<i>Deutzia gracilis</i>	Slender Gracilis
<i>Dianthus deltoides</i>	Zing Maiden Pinks
<i>Dianthus gratianopolitanus</i>	Firewitch Cheddar Pink
<i>Dianthus gratianopolitanus</i>	Crimson treasure Cheddar Pink
<i>Dianthus gratianopolitanus †</i>	Cheddar Pink
<i>Dianthus plumaris</i>	Cottage Pink
<i>Dietes vegeta</i>	Fortnight Lily
<i>Dodona viscosa</i>	Hopseed Bush
<i>Drosanthemum floribundum</i>	Trailing rosea iceplant
<i>Drosanthemum hispidum</i>	Ice Plant
<i>Dryopteris australis</i>	Dixie Wood Fern
<i>Dryopteris erythrosora</i>	Autumn Fern
<i>Dryopteris marginalis</i>	Marginal Wood Fern
<i>Dymondia margaritae</i>	Diamond Marguerite
<i>Dyopsis lutescens</i>	Areca Palm
<i>Elaeagnus ebbengei</i>	Gild Edge Elaeagnus
<i>Elaeagnus pungens</i>	Silverberry
<i>Elaeagnus pungens</i>	Fruitland Silver-Berry
<i>Enkianthus companulatus</i>	Red Veined Enkianthus
<i>Ensete ventricosum</i>	Absynnian Banana
<i>Equsetum scirpoides</i>	Dwarf Horsetail
<i>Erianthus ravennae</i>	Hardy Pampasgrass
<i>Erica cinerea</i>	Purple Bell Heather
<i>Erica darleyensana</i>	Mediterranean Pink Heather
<i>Erica vagans</i>	Cornish Heather
<i>Erigeron speciosum</i>	Darkest of Al Fleabane

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by "/" denotes and/or options.

Label amendment 09 29 2014

<i>Escalonia spp</i>	escallonia
<i>Eucalyptus camaldulensis</i>	Red Gum Eucalyptus
<i>Eucalyptus cinerea</i>	Silver Dollar Eucalyptus
<i>Eucalyptus microtheca</i>	Coolibah Tree
<i>Eucalyptus sideroxylon</i>	Red ironbark Eucalyptus
<i>Eugenia myritifolia</i>	Teenie Genie Brushcherry
<i>Eugenia myritifolia</i>	Dwarf Brush Cherry
<i>Euonymus kiatschovicus</i>	Mahattan Euonymus
<i>Euonymus fortunei</i>	Wintercreeper 'Blondy'
<i>Euonymus fortunei</i>	Wintercreeper
<i>Euonymus fortunei</i>	Canadale Gold Euonymus
<i>Euonymus fortunei</i>	Emerald & Gold Euonymus
<i>Euonymus fortunei</i>	Sunspot Euonymus
<i>Euonymus fortunei vegetus</i>	Bigleaf Wintercreeper
<i>Euonymus japonica</i>	Japanese Spindle Tree (Evergreen Euonymus) "gold spot"
<i>Euonymus japonica</i>	Japanese Spindle Tree
<i>Euonymus japonicus</i>	Silver King Euonymus
<i>Euonymus japonicus</i>	Chollipo Euonymus
<i>Euonymus japonicus</i>	Gold Spot Euonymus
<i>Euonymus japonicus</i>	Silver Princess Euonymus
<i>Euonymus japonicus</i>	Variegated Euonymus
<i>Euonymus kiatschovicus</i>	Spreading Euonymus
<i>Euryops pectinatus</i>	Dwarf Euryops
<i>Eustoma grandiflorum</i>	Pink lisianthus
<i>Evolvulus nuttallianus</i>	Blue Daze Evolvulus
<i>Fagus sylvatica</i>	European Beach
<i>Fatshedra japonica</i>	Japanese Aralia
<i>Fatsia japonica</i>	Japanese Aralia
<i>Felicia amelloides</i>	Blue Marguerite
<i>Festuca ovina</i>	Blue Fescue
<i>Ficus benjamina</i>	Indian Laurel Fig
<i>Ficus benjamina</i>	Mini Indian Laurel Fig
<i>Forsythia intermedia</i>	Border Forsythia
<i>Forsythia</i>	Spring Glory Forsythia
<i>Forsythia suspensa</i>	Weeping Forsythia
<i>Forsythia viridissima</i>	Greenstem Forsythia
<i>Fraxinus udhei</i>	Shamel Ash
<i>Fuchsia</i>	Santa Clause Fuchsia
<i>Gardenia jasminoides</i>	Gardenia, Cape-Jasmine

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Gardenia jasminoides</i>	Gardenia 'Mystery'
<i>Gardenia jasminoides</i>	August Beauty Gardenia
<i>Gardenia jasminoides</i>	Miniature Gardenia
<i>Gardenia radicans</i>	Dwarf Gardenia
<i>Gaultheria procumbens</i>	Wintergreen
<i>Gaultheria shallon</i>	Salal/Lemon Leaf
<i>Gaura</i> spp. †	Guara
<i>Gazania</i> spp	Gazania
<i>Gelsemium sempervirens</i>	Carolina jessamine
<i>Genista pilosa</i>	Woadwaxen
<i>Gentiana dahurica</i> †	Gentian
<i>Geranium cinereum</i> †	Cranesbill
<i>Geranium cinerium</i>	Ballerina Cranesbill
<i>Geranium sanguineum</i>	Bloody Cranesbill
<i>Geranium subcaulescens</i>	Black Eyed Magenta Cranesbill
<i>Geum quellyon</i>	Geum
<i>Geum</i> spp	avens
<i>Ginko biloba</i>	Ginko
<i>Gleditsia triacanthos</i>	Thornless Honey Locust
<i>Gleditsia triacanthos</i>	Shademaster Honeylocust
<i>Gypsophila paniculata</i>	Babys Breath
<i>Gypsophila repens</i> †	Baby's Breath
<i>Hakonechloa macroaureolo</i>	Golden hakonechloa
<i>Hamamelis virginia</i>	Common Witch Hazel
<i>Hamelia patens</i>	Firebush
<i>Hardenbergia violacea</i>	Lilac Vine
<i>Hebe buxifolia</i>	Boxleaf hebe
<i>Hedera canariensis</i>	Algerian Ivy
<i>Hedera helix</i>	English Ivy
<i>Helianthemum</i> spp. †	Sunrose
<i>Helichrysum petiolare</i>	White Licorice helichrysum
<i>Heliotropium fragrans</i>	Common Heliotrope
<i>Hemerocallis</i> spp	Daylily
<i>Hemerocallis</i> spp	Aztec Gold Daylily
<i>Hemerocallis</i> spp	Stella De Oro Daylily
<i>Hemerocallis</i> spp	Tender Love Daylily
<i>Hemerocallis</i> spp. †	Daylily: Aztec Gold, Stella De Oro, Tender Love, Mini Pearl
<i>Hesperalo parvifolia</i>	Red Yucca
<i>Heteromeles arbutiflora</i>	Toyon

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by "/" denotes and/or options.

Label amendment 09 29 2014

<i>Heucherella</i> spp	Coral Bell
<i>Heucherella</i> spp	Bridget Bloom
<i>Heucherella</i> spp. †	Coral Bell
<i>Hibiscus rosa-sinensis</i>	Ross Estey Hibiscus
<i>Hibiscus syriacus</i>	Red Bird Rose of Sharon
<i>Hibiscus syriacus</i>	Red Heart Rose of Sharon
<i>Hibiscus syriacus</i>	Woodbridge Rose of Sharon
<i>Hibiscus Rosa-sinensis</i> **	Chinese Hibiscus**
<i>Hibiscus</i> spp	Disco Bell White Mallow
<i>Hibiscus</i> **	Rose of Sharon**
<i>Hosta</i>	Francee Plantain Lily
<i>Hosta</i>	Patriot Plantain Lily
<i>Hosta lancifolia</i>	Albo-Marginata Hosta
<i>Hosta lancifolia</i>	Narrow Leafed Plantain Lily
<i>Hosta fortunei</i>	Plantain Lily
<i>Hosta plantaginea</i> †	Hosta, Plantain Lily (Fragrant)
<i>Hosta sieboldiana</i> †	Hosta, 'Searsucker'
<i>Hosta</i> x 'Goldstandard'	Hosta 'Goldstandard'
<i>Houttuynia cordata</i>	Chameleon houttuynia
<i>Houttuynia cordata</i>	Variegated Houttuynia
<i>Huchera</i>	Bressingham Coral Bells
<i>Huchera micrantha</i>	Coral Bells
<i>Hymenoxys acaulis</i>	angelita daisy
<i>Hypericum</i> spp	St Johnswort
<i>Ilex</i>	Nellie R Stevens Holly
<i>Ilex aquafolium</i>	Balkans English Holly
<i>Ilex aquafolium</i>	Gold Coast English Holly
<i>Ilex aquipernyi</i>	San Jose Holly
<i>Ilex attenuata</i>	Savannah Holly
<i>Ilex cassine</i>	Cassine Holly
<i>Ilex cornuta</i>	Burford Holly
<i>Ilex cornuta</i>	Needlepoint Holly
<i>Ilex cornuta</i>	Dwarf Burford Holly
<i>Ilex cornuta</i> **	Chinese Holly**
<i>Ilex crenata</i>	Japanese Holly
<i>Ilex crenata</i>	Compacta Holly
<i>Ilex crenata</i>	Convexa Holly
<i>Ilex crenata</i>	Dwarf Japanese Holly
<i>Ilex crenata</i>	Green Luster Holly
<i>Ilex crenata</i>	Helleri Holly

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by "/" denotes and/or options.

Label amendment 09 29 2014

<i>Ilex crenata</i>	Hetzi Holly
<i>Ilex crenata</i>	Steeds Japanese Holly
<i>Ilex crenata</i>	Stokes Japanese Holly
<i>Ilex glabra</i>	Nordic Inkberry Holly
<i>Ilex glabra</i> 'Compacta'	Inkberry
<i>Ilex meserveae</i>	Blue Boy Holly
<i>Ilex meserveae</i>	Blue Girl Holly
<i>Ilex meserveae</i>	China Boy Holly
<i>Ilex meserveae</i>	China Girl Holly
<i>Ilex meserveae</i>	Ebony Magic Holly
<i>Ilex opaca</i>	American Holly
<i>Ilex pernyi</i>	Holly
<i>Ilex vomitoria</i>	Yaupon Holly
<i>Ilex vomitoria</i>	Dwarf Yaupon Holly
<i>Ilex vomitoria</i>	Weeping Yaupon Holly
<i>Ilex</i> x	Holly 'Mary Nell'
<i>Illicium annisatum</i>	
<i>Illicium floridanum</i>	Florida Anise Tree
<i>Impatiens walleryana</i>	Lipstick Impatiens
<i>Imperata cylindrical</i>	Japanese Blood Grass
<i>Inula ensifolia</i>	
<i>Ipomea acuminata</i>	Blue Dawn Morning Glory
<i>Iris ensata</i>	Sword-Leaved Iris, Jodlesong
<i>Iris pumila</i>	Yellow dwarf bearded Iris
<i>Iris siberica</i>	Cabernet Siberian Iris
<i>Iris</i> spp.**	Iris species**
<i>Iris virginica</i>	Virginia Iris
<i>Itea virginica</i>	Henry Garnet Sweetspire
<i>Ixora collinea</i>	Ixora
<i>Jasminium nudiflorum</i>	Winter Jasmine
<i>Jasminum nitidum</i>	Angelwing Jasmine
<i>Jasminum polyanthemum</i>	Pink Jasmine
<i>Juglans</i> spp.*	Walnut*
<i>Juniperus chinensis</i>	Chinese Juniper
<i>Juniperus chinensis</i>	Gold Coast Juniper
<i>Juniperus chinensis</i>	Old Gold Juniper
<i>Juniperus chinensis</i>	Pfizer Juniper
<i>Juniperus chinensis</i>	Blue Pfizer Juniper
<i>Juniperus chinensis</i>	Sea Green Juniper
<i>Juniperus conferta</i>	Shore Juniper

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Juniperus conferta</i>	Emerald Sea Shore Juniper
<i>Juniperus davuricae</i> †	Parsoni
<i>Juniperus horizontalis</i>	Creeping Juniper (except “prince of wales” variety)
<i>Juniperus horizontalis</i>	Andorra Juniper
<i>Juniperus horizontalis</i>	Bar Harbor Juniper
<i>Juniperus horizontalis</i>	Blue Chip Juniper
<i>Juniperus horizontalis</i>	Blue Rug Juniper
<i>Juniperus horizontalis</i>	Creeping Juniper
<i>Juniperus horizontalis</i>	Dwarf Andorra Juniper
<i>Juniperus horizontalis</i>	Huntington Blue Juniper
<i>Juniperus horizontalis</i>	Andorra Juniper
<i>Juniperus horizontalis</i>	Blue Carpet Juniper
<i>Juniperus procumbens</i>	Dwarf Japanese Garden Juniper
<i>Juniperus prostrata</i>	Prostrate Juniper
<i>Juniperus sabina</i>	Broadmoor Juniper
<i>Juniperus sabina</i>	Hicks Juniper
<i>Juniperus sabina</i>	Savin Juniper
<i>Juniperus sabina</i>	Tam Juniper
<i>Juniperus scopulorum</i>	Emerald Green Juniper
<i>Juniperus scopulorum</i>	Wichita Blue Juniper
<i>Juniperus squamata</i>	Blue Juniper
<i>Juniperus squamata</i>	Blue Star Juniper
<i>Juniperus squamata</i>	Parson's Juniper
<i>Juniperus torulosa</i>	Hollywood Juniper
<i>Juniperus virginiana</i>	Eastern Red Cedar
<i>Justica brandegeana</i>	Shrimp Plant
<i>Kalmia latifolia</i>	Mountain Laurel
<i>Kniphofia uvaria</i>	Flamenco Red Hot Poker
<i>Lagerstroemia indica</i>	Crape Myrtle
<i>Lagerstroemia indica</i> X <i>fauriei</i>	Tuscarora Crape Myrtle
<i>Lagerstromia indica</i>	Crape Myrtle 'Burgundy Cotton'
<i>Lagerstromia indica</i> and <i>hybrid</i> †	Crape Myrtle
<i>Lampranthus spectabilis</i>	Trailing iceplant
<i>Lantana montevidensis</i> †	Weeping Lantana
<i>Lavender</i> spp. †	Lavender; Munstead
<i>Leontopodium alpinum</i> †	Edelweiss
<i>Leptospermum scoparium</i>	Broom Teatree/Manuka
<i>Leptospermum scoparium</i>	New Zealand Tea Tree
<i>Leptospermum scoparium</i>	Ruby Glow New Zealand Tea Tree
<i>Leucophyllum frutescens</i>	Texas Sage

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Leucophyllum laevigatum</i>	Chihuahan Sage
<i>Leucothoe axillaris</i>	Coast leucothoe
<i>Leucothoe fontanesiana</i>	Drooping Leucothoe
<i>Liatris spicata</i>	Floristan violet gay feather
<i>Ligustrum texanum</i>	Howard's Privet
<i>Ligustrum vulgare</i>	Lodense Common Privet
<i>Ligustrum amurense</i>	Amur Privet
<i>Ligustrum japonicum</i>	Japanese Privet
<i>Ligustrum japonicum</i>	Wax Privet
<i>Ligustrum japonicum</i>	Variegated Wax Privet
<i>Ligustrum lucidum</i>	Glossy Privet
<i>Ligustrum ovalifolium</i>	California Privet
<i>Ligustrum sinense†</i>	Chinese Privet
<i>Ligustrum sinense†</i>	Variegated Chinese Privet
<i>Ligustrum texanum</i>	Wax Leaf Ligustrum
<i>Ligustrum vulgaris</i>	Cheyenne' Ligustrum
<i>Ligustrum x vicaryi</i>	Golden Privet
<i>Lilium spp</i>	Jazz Lily
<i>Lilium spp. †</i>	Lily
<i>Limonium perezii</i>	Status
<i>Liquidambar styraciflua</i>	American Sweetgum
<i>Liriope gigantea</i>	White Lily Turf
<i>Liriope gigantea</i>	Giant Lily Turf
<i>Liriope muscari</i>	Liriope; Big Blue Lillyturf
<i>Liriope muscari</i>	Lilac Beauty Lily Turf
<i>Liriope muscari</i>	Majestic Lily Turf
<i>Liriope muscari</i>	Monroe White Lily Turf
<i>Liriope muscari</i>	Silvery Sunproof Lily Turf
<i>Liriope muscari</i>	Variegated Lily Turf
<i>Liriope spicata</i>	Green/Creeping Lily Turf
<i>Liriope spicata</i>	Silver Dragon Lily Turf
<i>Liriope spicata†</i>	Liriope, creeping
<i>Livistona chinensis</i>	Chinese Fan Palm
<i>Lobelia cardinalis†</i>	Cardinal flower, Indian Pink
<i>Lonicera fragrantissima</i>	Winter Honeysuckle
<i>Lonicera japonica</i>	Japanese Honeysuckle
<i>Lonicera periclymenum</i>	flowering Woodbine
<i>Lonicera periclymenum</i>	Serotina Woodbine
<i>Lonicera sempervirens</i>	Trumpet Honeysuckle
<i>Lonicera tatarica</i>	Tatarian Honeysuckle

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by "/" denotes and/or options.

Label amendment 09 29 2014

<i>Loropetalum chinense</i> †	Loropetalum 'Burgundy'
<i>Loropetalum chinense</i> †	Loropetalum 'Ruby'
<i>Loropetalum chinensis</i>	Burgandy Loropetalum
<i>Loropetalum chinensis</i>	Sizzling Pink Fringe Flower
<i>Loropetalum chinensis</i>	Razzleberri Fringe Flower
<i>Lysimacia mummularia</i>	Moneywort
<i>Lysimacia punctata</i>	Dotted Loosestrife
<i>Lythrum</i> spp. †	Loosestrife; Modern Pink
<i>Magnolia grandiflora</i>	Southern Magnolia
<i>Magnolia soulangeana</i>	Saucer Magnolia
<i>Magnolia</i> spp.**	Magnolia species**
<i>Magnolia stellata</i>	Royal Star Magnolia
<i>Mahonia aquafolium</i>	Dwarf Oregon Grape Holly/Mahonia
<i>Mahonia bealei</i>	Leather Leaf Mahonia
<i>Mahonia repens</i>	Creeping Mahonia
<i>Maleophora luteola</i>	Ice Plant
<i>Malus sargentii</i>	Non-Bearing Crabapple
<i>Malus</i> spp.*	Crabapple*
<i>Mandevilla splendens</i>	Red Riding Hood Mandevilla
<i>Matteuccia struthiopteris</i>	Ostrich Fern
<i>Matthiola incana</i>	Stock
<i>Michelia figo</i>	Banana Shrub
<i>Miscanthus sinensis</i>	Maiden Grass
<i>Miscanthus sinensis</i>	Silberfeder Miscanthus
<i>Miscanthus sinensis</i> **†	Yaku Jima**, Silberfeder**
<i>Moraea iridioides</i>	African Iris
<i>Morus alba</i>	White Mulberry
<i>Musa acuminata</i>	Banana
<i>myoporum parvifolium</i>	Putah Creek
<i>Myrica cerifera</i>	Southern Waxmyrtle
<i>Myrica pennsylvanica</i>	Bayberry
<i>Nandina domestica</i>	Compact Dwarf Heavenly Bamboo
<i>Nandina domestica</i>	Harbour Dwarf Nandina
<i>Nandina domestica</i>	Nandina
<i>Nandina domestica</i>	Dwarf Purple Nandina
<i>Nandina domestica</i>	Woods Dwarf Nandina
<i>Narcissus</i> spp.**	Narcissus species**
<i>Nerium oleander</i>	Hardy Red Oleander
<i>Nerium oleander</i>	Oleander
<i>Nerium</i> spp.	Oleander

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by "/" denotes and/or options.

Label amendment 09 29 2014

<i>Nipponanthemum nipponicum</i> †	Montauk Daisy or Nippon Daisy
<i>Oenothera missouriensis</i> †	Evening Primrose, Ozark Sundrops
<i>Oenothera speciosa</i>	Siskiyou Evening Primrose
<i>Olea europaea</i> *	Olive*
<i>Onoclea sensibilis</i>	Sensitive Fern
<i>Ophiopogon japonicus</i>	Dwarf Mondo Grass
<i>Ophiopogon japonicus</i> **	Mondo Grass**
<i>Origanum libanoticum</i>	Oregano
<i>Osmanthus fortunei</i>	Fortune's Osmanthus
<i>Osmanthus fragrans</i>	Sweet Olive Osmanthus
<i>Osmanthus heterophyllus</i>	Osmanthus, False Holly
<i>Osmanthus heterophyllus</i> †	Osmanthus (False Holly): Gulf Tide
<i>Osmunda cinnamomea</i>	Cinnamon Fern
<i>Osmunda regalis</i>	Royal Fern
<i>Osteospermum fruticosum</i> †	Trailing African Daisy
<i>Oxydendrum arboreum</i>	Sourwood
<i>Pachysandra terminalis</i>	Japanese Spurge
<i>Pachysandra terminalis</i>	Green Sheen Japanese Spurge
<i>Paeonia suffruticosa</i> †	Tree Peony
<i>Parthenocissus quinquefolia</i>	Virginia Creeper
<i>Pelargonium hortorum</i>	Zonal Geranium
<i>Pelargonium peltatum</i>	Ivy Geranium
<i>Pennisetum alopecuroides</i>	Fountain Grass
<i>Pennisetum setaceum</i>	Chrimson Fountaingrass
<i>Pennisetum setaceum</i>	Purple Fountain Grass
<i>Pennisetum setaceum</i> **†	Fountain Grass (Dwarf)**
<i>Penstemon</i>	Apple Blossom Penstemon
<i>Pentas lanceolata</i>	Star Clusters
<i>Perovskia atriplicifolia</i> †	Russian Sage
<i>Persea americana</i> *	Avocado*
<i>Petunia hybrids</i>	Petunias
<i>Phalaris arundinacea</i>	Ribbon Grass
<i>Phlox subulata</i>	Moss Pink
<i>Phoenix roebelenii</i>	Pygmy Date Palm
<i>Phormium tenax</i>	Jack Spratt New Zealand Flax
<i>Photinia fraseri</i>	Red Tip Photinia
<i>Physocarpus opulifolius</i>	Dwarf Ninebark
<i>Physostegia virginiana</i> †	Dragonhead, False; Vivid
<i>Picea pungens</i>	Colorado Blue Spruce
<i>Picea abies</i>	Weeping Norway Spruce

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Picea abies</i>	Spreading Norway Spruce
<i>Picea abies</i>	Norway Spruce
<i>Picea glauca</i>	White Spruce
<i>Picea glauca</i>	Dwarf Alberta Spruce
<i>Picea pungens</i>	Hoopsi Blue Spruce
<i>Picea pungens</i>	Koster Blue Spruce
<i>Picea spp</i>	Spruces
<i>Picea spp.</i> **	Spruce species** (Colorado Blue, Norway, etc.)
<i>Pieris</i>	Forest Flame Pieris
<i>Pieris japonica</i>	Japanese Andromeda; Lily-of-the-Valley Shrub
<i>Pieris japonica</i>	Mountain of Fire Pieris
<i>Pieris japonica</i>	Snowdrift Pieris
<i>Pieris japonica</i>	Temple Bells Pieris
<i>Pieris japonica</i>	Valley Rose Pieris
<i>Pieris japonica</i>	Valley Valentine Pieris
<i>Pinus mugo</i>	Mugo Pine
<i>Pinus aristata</i>	Bristlecone Pine
<i>Pinus brutia</i>	Calabrian Pine
<i>Pinus canariensis</i>	Canary Island Pine
<i>Pinus contorta</i>	Shore Pine, Beach Pine
<i>Pinus eldarica</i>	Eldarica Pine
<i>Pinus elliottii</i>	Slash Pine
<i>Pinus halapensis</i>	Aleppo Pine
<i>Pinus leucodermis</i>	Bosnian Pine
<i>Pinus nigra</i>	Austrian Black Pine
<i>Pinus palustris</i>	Longleaf Pine
<i>Pinus radiata</i>	Monterey Pine
<i>Pinus strobus</i>	Eastern White Pine
<i>Pinus sylvestris</i>	Scotch Pine
<i>Pinus sylvestris</i>	Columnar Scotch Pine
<i>Pinus taeda</i>	Loblolly Pine
<i>Pinus thunbergii</i>	Japanese Black Pine
<i>Pinus virginiana</i>	Virginia Pine
<i>Pistacia spp.*</i>	Pistachio*
<i>Pittosporum rhombifolium</i>	Queensland Pittosporum
<i>Pittosporum tenuifolia</i>	Golf Ball Pittosporum
<i>Pittosporum tobira</i>	Japanese Pittosporum
<i>Pittosporum tobira</i>	Green Pittosporum
<i>Pittosporum tobira</i>	Wheelers Dwarf Pittosporum
<i>Platanus occidentalis</i>	American Sycamore

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by "/" denotes and/or options.

Label amendment 09 29 2014

<i>Platanus racemosa</i>	California Sycamore
<i>Plumbago auriculata</i>	Plumbago
<i>Plumbago capensis</i>	Plumbago
<i>Podocarpus macrophyllus</i>	Yew Pine/Japanese Yew
<i>Podocarpus Species</i>	Podocarpus
<i>Polygala fructicosa</i>	Sweet Pea Shrub
<i>Polystichum acrostichoides</i>	Christmas Fern
<i>Polystichum polyblepharum</i>	Tassel Fern
<i>Populus deltoides</i>	Cottonwood
<i>Potentilla fragiformis</i>	Cinquefoil
<i>Potentilla fruticosa</i>	Cinquefoil
<i>Potentilla fruticosa</i>	Floppy Disc Cinquefoil
<i>Potentilla fruticosa</i>	Gold Drop Cinquefoil
<i>Potentilla fruticosa</i>	Goldfinger Cinquefoil
<i>Potentilla fruticosa</i>	Red Ace Cinquefoil
<i>Potentilla fruticosa</i>	Sunset Cinquefoil
<i>Potentilla fruticosa</i>	Tangerine Cinquefoil
<i>Potentilla spp</i>	Cinquefoil
<i>Potentilla verna</i>	Spring Cinquefoil
<i>Prosopis chiliensis chiliensis</i>	Chilean Mesquite
<i>Prunus laurocerasus</i>	English Laurel
<i>Prunus laurocerasus</i>	Otto Luyken English Laurel
<i>Prunus caroliniana</i>	Carolina Cherry Laurel
<i>Prunus caroliniana</i>	Bright 'n Tight Cherry Laurel
<i>Prunus glandulosa</i>	Dwarf Flowering Almond
<i>Prunus spp.*</i>	Almond, Apricot, Nectarine, Peach, Plum and Prune*
<i>Prunus x yedoensis</i>	Yoshino Cherry
<i>Pseudotsuga menziesii**</i>	Douglas Fir**
<i>Psidium cattleianum</i>	Strawberry Guava
<i>Pyracantha coccinea</i>	Firethorn Scarlet
<i>Pyracantha coccinea</i>	Lalandei Firethorn
<i>Pyracantha fortuneana</i>	Firethorn
<i>Pyracantha fortuneana</i>	Lolendei Monrovia Pyracantha
<i>Pyracantha fortuneana</i>	Monon Pyracantha
<i>Pyracantha fortuneana</i>	Red Elf Pyracantha
<i>Pyracantha fortuneana</i>	Rutgers Hybrid Pyracantha
<i>Pyracantha fortuneana</i>	Santa Cruz Pyracantha
<i>Pyracantha fortuneana</i>	Victory Pyracantha
<i>Pyracantha koidzumii</i>	Pyracantha 'Santa Cruz'
<i>Pyracantha koidzumii</i>	Firethorn

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Pyrus spp</i>	Bradford Pears
<i>Pyrus spp.</i>	Pear spp. including Bradford
<i>Quercus ilicifolia</i>	Bear Oak
<i>Quercus palustris</i>	Pin Oak
<i>Quercus phellos</i>	Willow Oak
<i>Quercus rubra</i>	Red Oak
<i>Quercus shumardii†</i>	Oak, Shumard's Red
<i>Raphiolepis indica</i>	Indian Hawthorne
<i>Raphiolepis umbellata†</i>	Yedda Hawthorne
<i>Ratbida columnifera</i>	Mexican Hat
<i>Ravenea rivularis</i>	Majesty Palm
<i>Rhaphiolepis indica</i>	Charisma-monruce Indian Hawthorn
<i>Rhaphiolepis indica</i>	Enchantress-moneess Indian Hawthorn
<i>Rhaphiolepis indica</i>	Indian Hawthorn
<i>Rhaphiolepis indica</i>	Springtime Indian Hawthorn
<i>Rhaphiolepis indica</i>	Ballerina Indian Hawthorn
<i>Rhaphiolepis ovata</i>	Round-Leaf Indian Hawthorn
<i>Rhododendron</i>	Fashion Azalea
<i>Rhododendron</i>	Wakeiebisu Azalea
<i>Rhododendron</i>	George Tabor Azalea
<i>Rhododendron</i>	Gerard Azalea
<i>Rhododendron</i>	Delaware Valley White Azalea
<i>Rhododendron</i>	Girard Roberta Azalea
<i>Rhododendron</i>	Girard Crimson Azalea
<i>Rhododendron</i>	Golden Flare Exbury Azalea
<i>Rhododendron</i>	Helmut Vogel Azalea
<i>Rhododendron</i>	Hershey Red Azalea
<i>Rhododendron</i>	Hot Shot Azalea
<i>Rhododendron</i>	Hume Azalea
<i>Rhododendron</i>	Inga Azalea
<i>Rhododendron</i>	Irene Koster Azalea
<i>Rhododendron</i>	President Clay Azalea
<i>Rhododendron</i>	Traditional Azalea
<i>Rhododendron forrestii x griersonianum</i>	Elizabeth Rhododendron
<i>Rhododendron calendulaceum</i>	Flame Azalea
<i>Rhododendron calendulaceum</i>	Golden Fire Azalea
<i>Rhododendron calendulaceum</i>	Klondike Azalea
<i>Rhododendron campylocarpum</i>	Butterfly Rhododendron
<i>Rhododendron catawbiense</i>	White Catawba Rhododendron
<i>Rhododendron catawbiense</i>	Catawba Rhododendron

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Rhododendron catawbiense</i>	Lord Roberts Catawba Rhododendron
<i>Rhododendron catawbiense</i>	Rocket Catawba Rhododendron
<i>Rhododendron caucasicum X ponticum</i>	Cunningham White Rhododendron
<i>Rhododendron exbury</i>	Canon's Double Exbury Azalea
<i>Rhododendron exbury</i>	Golden Flare Exbury Azalea
<i>Rhododendron exbury</i>	Klondike Exbury Azalea
<i>Rhododendron forrestii</i>	Gomer Waterer Rhododendron
<i>Rhododendron griffithianum</i>	Jean Marie Rhododendron
<i>Rhododendron hybrid spp</i>	America Rhododendron
<i>Rhododendron hybrid spp</i>	Nova Zembla Rhododendron
<i>Rhododendron hybrid spp</i>	Scintillation Rhododendron
<i>Rhododendron Hybrids</i>	Carror Azalea
<i>Rhododendron impeditum</i>	Rhododendron
<i>Rhododendron indica</i>	Formosa Azalea
<i>Rhododendron indica</i>	Waucabusa Azalea
<i>Rhododendron kaempferi</i>	Blue Danube Azalea
<i>Rhododendron kerume</i>	Coral Bells Azalea
<i>Rhododendron kerume</i>	Hino Crimson Azalea
<i>Rhododendron kerume</i>	Hino Pink Azalea
<i>Rhododendron kerume</i>	Mildred Azalea
<i>Rhododendron kerume</i>	Snow Azalea
<i>Rhododendron maximum</i>	Rose Bay Rhododendron
<i>Rhododendron mucronulatum</i>	Rhododendron
<i>Rhododendron obtusum</i>	Hino Crimson Azalea
<i>Rhododendron ponticum</i>	Chioniodes Rhododendron
<i>Rhododendron ponticum</i>	Daphinoides Rhododendron
<i>Rhododendron racemosum</i>	Dwarf Scarlet Wonder Rhododendron
<i>Rhododendron racemosum</i>	Tribly Rhododendron
<i>Rhododendron racemosum</i>	Unique Rhododendron
<i>Rhododendron racemosum</i>	Vulcan Rhododendron
<i>Rhododendron sassthigiatum X carolinianum</i>	Ramapo Rhododendron
<i>Rhododendron satuski</i>	Gumpo Pink Azalea
<i>Rhododendron satuski</i>	Gumpo White Azalea
<i>Rhododendron satuski</i>	Higasa Azalea
<i>Rhododendron satuski</i>	Reijn Azalea
<i>Rhododendron schlippenbachii</i>	Royal Azalea
<i>Rhododendron spp. †</i>	Rhododendrons, Azaleas (Except Carolina and roseum elegans variety rhododendron)
<i>Rhododendron x 'George Tabor'</i>	Azalea 'George Tabor'

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Rhododendron x obtusum</i>	Kurume Azalea 'Tradition'
<i>Rhus lancea</i>	African Sumac
<i>Rhus typhina</i>	Staghorn Sumac
<i>Rosa banksiae</i>	Lady Bank's Rose
<i>Rosa Flower Carpet</i>	Red Groundcover Rose
<i>Rosa rugosa</i>	Rugosa Rose
<i>Rosa x</i>	Knock Out Rose Radrazz
<i>Rosa x</i>	Knock Out Double Pink
<i>Rosmarinus officinalis</i>	Rosemary
<i>Rudbeckia fulgida</i>	Black-Eyed Susan 'Early Bird'
<i>Rudbeckia fulgida</i>	Blackeyed Susan
<i>Rudbeckia fulgida †</i>	Black-Eyed Susan 'Goldstrum'
<i>Rumohra adiantiformis</i>	Leatherleaf Fern
<i>Ruscus hypophyllum</i>	Butchers Broom
<i>Salix babylonica</i>	Babylon Weeping Willow
<i>Salix matsudana</i>	Corkscrew Willow
<i>Salvia daghestancia</i>	Platinum Sage
<i>Salvia grahamii</i>	Grahams Sage
<i>Santolina virens</i>	Green Lavender Cotton
<i>Santolina virens</i>	Green Santolina
<i>Sasa pygmaea</i>	Dwarf Bamboo
<i>Saxifraga spp. †</i>	Saxifrage; Purple Dome
<i>Scabiosa spp. †</i>	Pincushion flower
<i>Scutellaria resinosa</i>	Skull Cap
<i>Sedum pachyclados</i>	Grey Stonecrop
<i>Sedum spurium</i>	Sedum 'John Creech'
<i>Senecio cineraria</i>	Dusty Miller
<i>Sequoia sempervirens</i>	Coast Redwood
<i>Sequoiadendron giganteum</i>	Giant Sequoia
<i>Skimmia japonica</i>	Japanese skimmia
<i>Skimmia revesiana</i>	Reeves Skimmia
<i>Solanum rantonetii</i>	Paraguay Nightshade
<i>Spiraea bumalda†</i>	Spirea 'Anthony Waterer'
<i>Spiraea cinerea</i>	First Snow Spiraea
<i>Spiraea japonica</i>	Dolchia Spiraea
<i>Spiraea japonica</i>	Japanese Alpine Spiraea
<i>Spiraea japonica</i>	Magic Carpet Spiraea
<i>Spiraea japonica</i>	Neon Flash Spiraea
<i>Spiraea japonica</i>	Shirobana Spiraea
<i>Spiraea vanhouttei</i>	Vanhouttei Spirea

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Spirea japonica</i>	Spirea 'Neon Flash'
<i>Spirea japonica</i>	Spirea 'Shirobana'
<i>Streptosolen jamesonii</i>	Marmalade Bush
<i>Swietenia mahogani</i>	Mahogany
<i>Syagrus romanzoffianum</i>	Queen Palm
<i>Syringa rothomagenesis</i>	Chinese Lilac
<i>Syringa vulgaris</i>	Common Lilac
<i>Syzygium paniculatum</i>	Japanese Boxcherry
<i>Syzygium paniculatum</i>	Australian Brushcherry
<i>Tabebuia caraiba</i>	Yellow Tabebuia
<i>Tagetes patula</i>	Little Hero Marigold
<i>Taxus cuspidata</i>	Japanese Yew
<i>Taxus media</i> †	Yew
<i>Tecomaria capensis</i>	Cape Honeysuckle
<i>Ternstroemia gymnanthera</i>	Japanese Ternstroemia
<i>Teucrium fruiticans</i>	Bush Germander
<i>Teucrium spp</i> †.	Germander
<i>Thalictrum dipterocarpum</i> †	Meadow Rue
<i>Thevetia nerifolia</i>	Yellow Oleander
<i>Thuja occidentalis</i>	American Arborvitae
<i>Thuja occidentalis</i>	Emerald Arborvitae
<i>Thuja occidentalis</i>	Globe Arborvitae
<i>Thuja occidentalis</i>	Little Giant Dwarf Arborvitae
<i>Thuja occidentalis</i>	Dark American Arborvitae
<i>Thuja occidentalis</i>	Pyramidal Arborvitae
<i>Thuja occidentalis</i>	Rheingold Arborvitae
<i>Thuja occidentalis</i>	Techny Arborvitae
<i>Thuja occidentalis</i>	Woodward's Arborvitae
<i>Thuja orientalis</i>	Dwarf Golden Arborvitae
<i>Thuja orientalis</i>	Dwarf Arborvitae
<i>Tibouchina urvilleana</i>	Princess Flower
<i>Trachelospermum asiaticum</i>	Asiatic Jasmine, Star Jasmine
<i>Trachycarpus fortunei</i>	Windmill Palm
<i>Tsuga canadensis</i>	Canada Hemlock, Eastern Hemlock
<i>Tulbaghia violacea</i>	Society Garlic
<i>Tulipa spp</i>	Tulips
<i>Ulmus parvifolia</i>	Chinese Elm
<i>Verbena rigida</i>	Veined Verbena
<i>Veronica spp</i>	Sunny Border, Veronica /Speedwell
<i>Veronica spp.</i> †	Veronica, Speedwell

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by "/" denotes and/or options.

Label amendment 09 29 2014

<i>Viburnum bodnantense</i>	Pink Dawn Viburnum
<i>Viburnum carlesii</i>	Koreanspice Viburnum
<i>Viburnum davidii</i>	David Viburnum
<i>Viburnum japonicum</i>	Japanese Viburnum
<i>Viburnum judd</i>	Judd Viburnum
<i>Viburnum lantana</i>	Wayfaringtree Viburnum
<i>Viburnum odoratissimum</i>	Sweet Viburnum
<i>Viburnum opulus</i> 'Roseum'	Roseum Viburnum
<i>Viburnum opulus sterile</i>	Common Snowball Viburnum
<i>Viburnum plicatum</i>	Japanese Snowball
<i>Viburnum plicatum</i>	Doublefile Viburnum
<i>Viburnum pragense</i>	Prague Viburnum
<i>Viburnum rigidum</i>	Canary Island Viburnum
<i>Viburnum setigerum</i>	Tea Viburnum
<i>Viburnum suspensum</i> †	Arrowwood viburnum
<i>Viburnum tinus</i>	Laurustinus
<i>Viburnum tinus</i>	Spring Bouquet Viburnum
<i>Viburnum trilobium</i>	Cranberry Bush
<i>Viburnum trilobum</i>	American Cranberrybush Viburnum
<i>Viburnum trilobum</i>	Dwarf American Cranberrybush Viburnum
<i>Viburnum wrightii</i>	Leatherleaf Viburnum
<i>Vinca major</i>	Vinca
<i>Vinca major</i>	Bigleaf Periwinkle
<i>Vinca minor</i>	Dwarf Periwinkle
<i>Vinca spp</i>	Periwinkle
<i>Vitis spp.*</i>	Grape*
<i>Washingtonia robusta</i>	Mexican Fan Palm
<i>Washingtonia robusta</i>	California Fan Palm
<i>Weigela florida</i>	Old Fashioned Weigela
<i>Weigelia florida</i>	Bristol Ruby Weigelia
<i>Weigelia florida</i>	Java Red Weigelia
<i>Weigelia florida</i>	Minuet Weigelia
<i>Xylosma congestum</i>	Xylosma
<i>Xylosma senticosa</i>	Shiny Xylosma
<i>Yucca aloifolia</i>	Spanish Bayonet
<i>Yucca filamentosa</i>	Yucca, Adam's Needle
<i>Zamia furfuracea</i>	Cardboard Palm

† Not for use in California

*Do not use on food producing trees, vines, or plants

**Not for use on container grown plants.

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

Table 12. THE PLANTS LISTED IN THIS TABLE ARE NOT LIKELY TO BE INJURED BY OF QUALI-PRO® PRODOXABEN 3.7 SC AT THE 1 OZ/1000 SQ FT (43.5 OZ/ACRE) RATE

Scientific name	Common name
<i>Abelia grandiflora</i>	Edward Goucher Glossy Abelia
<i>Abelia grandiflora</i>	Glossy Abelia
<i>Abelia grandiflora</i> †	Abelia: Sherwood
<i>Abies balsamea</i>	Balsam Fir
<i>Abies concolor</i>	White / Concolor Fir
<i>Abies spp.</i>	Fir species** (Balsam, Fraser, Noble, etc.)
<i>Abutilon hybridum</i>	Flowering Maple
<i>Acacia abyssinica</i>	Abyssinica Acacia
<i>Acacia redolens</i>	Prostrate Acacia
<i>Acacia stenophyll</i>	Shoestring Acacia
<i>Acalypha wilkesiana</i>	Copper Plant
<i>Acer ginnala</i>	Amur Maple
<i>Acer ginnala</i>	Flame Maple
<i>Acer palmatum</i>	Japanese Maple
<i>Acer palmatum</i>	Coral Bark Japanese Maple
<i>Acer palmatum</i>	Dwarf Japanese Maple
<i>Acer platanoides</i>	Norway Maple
<i>Acer saccharinum</i>	Silver Maple
<i>Achillea filipendulina</i>	Moonshine /fern-leaf yarrow
<i>Achillea millefolium</i>	Common Yarrow
<i>Achillea millefolium</i>	Paprika Yarrow
<i>Achillea millefolium</i>	Achillea 'Saucy Seduction'
<i>Achillea spp</i>	Yarrow
<i>Achillea spp.</i> †	Yarrow: King Edward
<i>Achillea tomentosa</i>	Wooly Yarrow
<i>Acoelorrhaphe wrightii</i>	Everglades Palm
<i>Actinidia chinensis</i> *	Kiwi*
<i>Adonidia merrilli</i>	Christmas Palm
<i>Agapanthus africanus</i>	Lily-of-the-Nile (African Lily)
<i>Agapanthus africanus</i>	Queen Anne Lily of the Nile
<i>Agapanthus orientalis</i> †	
<i>Agave americana</i>	Century Plant
<i>Akebia quintata</i> †	Five-Leaf or Chocolate Vine
<i>Albizia julibrissin</i>	Mimosa; Silk Tree
<i>Allium cernuum</i> †	Lady's Leek, Nodding Onion
<i>Alsophils australis</i>	Australian Tree Fern

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Amelanchier alnifolia</i>	Saskatoon Serviceberry
<i>Andromeda polifolia</i>	Bog Rosemary
<i>Anemone hybrida</i> †	Japanese Anemone
<i>Anisodonte hypomandarum</i>	Cape Mallow
<i>Aptenia cordifolia</i>	Red Apple Aptenia
<i>Aquilegia</i>	Dragon Fly Columbine
<i>Aquilegia</i>	Red & Gold Columbine
<i>Aquilegia spp.</i> †	Columbine
<i>Araucaria heterophylla</i>	Norfolk Island Pine
<i>Archontophoenix cunninghamiana</i>	King Palm
<i>Arctostaphylos densiflora</i>	Vine Hill Manzanita
<i>Arctostaphylos uva-ursi</i>	Bearberry
<i>Arctotheca calendula</i>	Cape Weed
<i>Ardisia japonica</i>	Chirimen Marlberry
<i>Argyranthemum frutescens</i>	Butterfly Argyranthemum
<i>Armeria maritima</i>	Bloodstone Sea Thrift
<i>Artemesia lactiflora</i>	White Mugwort
<i>Artemisia spp</i>	Silver Mound Wormwood
<i>Artemisia spp.</i> †	Wormwood; Silver Mound, Castle
<i>Asparagus</i>	
<i>Asparagus densiflorus</i>	Foxtail Fern
<i>Asparagus retrofractus</i>	Ming Asparagus Fern
<i>Asparagus variegata</i>	
<i>Aspidistra elatior</i>	Cast Iron Plant
<i>Aster</i>	Bonny Blue Aster
<i>Aster</i>	Purple Dome Aster
<i>Aster</i>	Aster
<i>Aster novae-angliae</i>	New England Aster
<i>Aster novi-belgii</i>	New York Aster
<i>Aster novi-belgii</i>	Persian Rose Dwarf Aster
<i>Aster spp.</i> †	Aster: Bonny Blue, Purple Dome
<i>Athyrium filix-femina</i> †	Lady Fern
<i>Athyrium nipponimcum</i>	Japanese Painted Fern
<i>Aucuba japonica</i>	Japanese Aucuba
<i>Aucuba japonica</i>	Goldstrike Aucuba, Goldstrike Japanese Laurel
<i>Baccharis pulularis</i>	Coyotebush
<i>Bauhinia galpinii</i>	Red Bauhinia
<i>Begonia semperflorens</i>	Ambassador White Begonia
<i>Begonia spp</i>	Fibrous Begonia
<i>Begonia spp.</i> †	Fibrous Begonia: Hardy Grandis

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Berberis gladwynensis</i>	William Penn Barberry
<i>Berberis gladwynensis</i>	Barberry
<i>Berberis julianae</i>	Wintergreen Barberry
<i>Berberis mentorensis</i>	Mentor Barberry
<i>Berberis thunbergii</i>	Japanese Barberry
<i>Berberis thunbergii</i>	Japanese Barberry 'Concord'
<i>Berberis thunbergii</i>	Crimson Pygmy Barberry
<i>Berberis thunbergii</i>	Golden Japanese Barberry
<i>Berberis thunbergii</i>	Rosy Glow Barberry
<i>Berberis thunbergii</i>	Redleaf Japanese Barberry
<i>Berberis thunbergii</i>	Cherry Bomb Barberry
<i>Berberis verruculosa</i>	Warty Barberry
<i>Bergenia cordifolia</i>	Heartleaf Bergenia
<i>Bergenia cordifolia</i> †	
<i>Betula nigra</i>	River Birch
<i>Betula papyrifera</i>	Paper Birch
<i>Betula pendula</i>	European White Birch
<i>Bidens ferulifolia</i>	Peters Gold Bidens
<i>Boltonia asteroides</i> †	Snowbank
<i>Bougainvillea spp</i>	Barbara Karst Bougainvillea
<i>Bougainvillea spp</i>	California Gold Bougainvillea
<i>Bougainvillea spp</i>	Pink Pixie Bougainvillea
<i>Bougainvillea spp</i>	Scarlet Ohara Bougainvillea
<i>Bougainvillea spp</i>	Temple Fire Bougainvillea
<i>Bougainvillea spp</i>	Texas Dawn Bougainvillea
<i>Bougainvillea spp</i>	Purple Queen Bougainvillea
<i>Bougainvillea spp</i>	Rosenka Bougainvillea
<i>Bougainvillea spp.</i> †	Bougainvillea
<i>Brachychiton populneus</i>	Bottle Tree
<i>Brachycome</i>	Swan River daisy-New amethyst
<i>Bucida buceras</i>	Black Olive
<i>Buddleia davidii</i>	Butterfly Bush
<i>Buddleia davidii</i> †	Butterfly-Bush 'Royal Red'
<i>Butia odorata</i>	Pindo Palm
<i>Buxus 'Green Velvet'</i>	Boxwood 'Green Velvet'
<i>Buxus microphylla</i>	Japanese Boxwood
<i>Buxus microphylla</i>	Green Beauty Boxwood
<i>Buxus microphylla</i>	Japanese Littleleaf Boxwood
<i>Buxus microphylla</i>	Korean Boxwood
<i>Buxus sempervirens</i>	Common Boxwood

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Callistemon citrinus</i>	Lemon Bottlebrush
<i>Callistemon citrinus</i>	Little John Bottlebrush
<i>Callistemon citrinus</i> †	Crimson Bottlebrush
<i>Callistemon viminalis</i>	Weeping Bottlebrush
<i>Callistepheus cinensis</i>	China Aster
<i>Calluna vulgaris</i>	Scotch Heather
<i>Calluna vulgaris</i>	Spring Torch Scotch Heather
<i>Camellia japonica</i>	Japanese Camellia
<i>Campanula carpatica</i>	Bellflower; Tussock flower
<i>Campanula carpatica</i> †	Tussock Bellflower; (White Clips)
<i>Campis X tagliabuana</i> †	Trumpet Creeper, Trumpet flower; Madame Galen
<i>Carex</i>	Variegated Carex
<i>Carex albula</i>	Frosty Curls Sedge
<i>Carpobrotus edulis</i>	Hottentot Fig (Ice Plant)
<i>Carpobrotus edulis</i>	Largeleaf Ice Plant
<i>Caryopteris clandonensis</i>	Blue Mist Bluebeard
<i>Caryopteris clandonensis</i>	Dark Knight Bluebeard
<i>Cassia artemisioides</i>	Feathery Cassia
<i>Cassia eremophila</i>	Senna
<i>Catharanthus roseus</i>	Madagascar Periwinkle
<i>Ceanothus rigidus</i>	Wild Lilac
<i>Ceanothus spp</i>	Wild Lilac
<i>Ceanothus x pallidus</i>	Lilac 'Marie Bleu'
<i>Cephalotaxus harringtonia</i>	Japanese Plum Yew
<i>Cerastium tomentosum</i>	Snow in Summer
<i>Ceratonia siliqua</i>	Carob
<i>Ceratostigma plumbaginoides</i> †	
<i>Ceratostigma willmottianum</i>	Chinese Plumbago
<i>Cercis canadensis</i>	Redbud
<i>Chaenomeles japonica</i>	Orange Flowering Quince
<i>Chamaecyparis obtusa</i>	Kosteri Cypress
<i>Chamaecyparis obtusa</i>	Dwarf Hinoki Cypress
<i>Chamaecyparis obtusa</i>	Torulosa Cypress
<i>Chamaecyparis obtusa</i>	Fernspray Falsecypress
<i>Chamaecyparis obtusa</i>	Slender Hinoki False Cypress
<i>Chamaecyparis pisifera</i>	False Cypress
<i>Chamaecyparis pisifera</i>	Squarrosa Minima False Cypress
<i>Chamaecyparis pisifera</i>	Threadleaf False Cypress
<i>Chamaecyparis pisifera</i>	Sawara False Cypress
<i>Chamaecyparis pisifera</i>	Moss Falsecypress

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by "/" denotes and/or options.

Label amendment 09 29 2014

<i>Chamaedorea cataractarum</i>	Cat Palm
<i>Chamaedorea costaricana</i>	Costa Rican Chamaedorea Palm
<i>Chamaedorea elegans</i>	Parlor Palm
<i>Chamaerops humilis</i>	European Fan Palm, Mediterranean Fan Palm
<i>Chasmanthium latifloium</i>	Northern Sea Oats
<i>Chrysanthemum maximum</i>	Shasa Daisy
<i>Chrysanthemum spp</i>	Chrysanthemum
<i>Citrus spp.</i> *	Citrus species*
<i>Clethra alnifolia</i>	Summersweet
<i>Cleyera japonica</i>	Cleyera, Japanese Cleyera
<i>Clivia miniata</i>	Kafir Lily
<i>Colonema pulchrum</i>	Pink Breath of Heaven
<i>Convolvulus cneorum</i>	Bush Morning Glory
<i>Convolvulus mauritanicus</i>	Ground Morning Glory
<i>Cordyline indivisa</i>	Blue Dracaena
<i>Coreopsis</i>	Early Sunrise Coreopsis
<i>Coreopsis grandiflora</i>	Coreopsis
<i>Coreopsis spp.</i> †	Coreopsis (Calliopsis): Early Sunrise, Moonbeam
<i>Coreopsis verticillata</i>	Coreopsis
<i>Cornus florida</i>	Cloud Nine Dogwood
<i>Cornus florida</i>	Flowering Dogwood
<i>Cornus kousa</i>	Kousa Dogwood
<i>Cornus alba</i>	Siberian Dogwood
<i>Cornus sericea</i>	Red Osier Dogwood
<i>Cornus sericea</i>	Yellowtwig Dogwood
<i>Cornus stolonifera</i>	American Dogwood
<i>Cortaderia selloana</i>	Pampas Grass
<i>Corylus americanus</i>	Harry Lauder's Walking Stick
<i>Cotinus coggygria</i>	Royal Purple Smoke Tree
<i>Cotinus dammeri</i>	Coral Beauty Smoke Tree
<i>Cotinus dammeri</i>	Eichholz Smoke Tree
<i>Cotoneaster adpressus</i>	Early Cotoneaster
<i>Cotoneaster apiculatus</i>	Cranberry Cotoneaster
<i>Cotoneaster buxifolius</i>	Box Leafed Cotoneaster
<i>Cotoneaster congestus</i>	Pyrenees Cotoneaster
<i>Cotoneaster dammeri</i>	Bearberry Cotoneaster
<i>Cotoneaster himalayan</i>	Himalayan Cotoneaster
<i>Cotoneaster horizontalis</i>	Rock Cotoneaster
<i>Cotoneaster microphyllus</i>	Rockspray Cotoneaster
<i>Crataegus Species</i>	Hawthorns

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Crataegus viridis</i>	Green Hawthorn
<i>Crocoshmia spp. †</i>	Lucifer
<i>Cryptomeria japonica</i>	Japanese Cryptomeria
<i>Cupaniopsis anacardioides</i>	Carrot Wood
<i>Cuphea hyssopifolia</i>	False or Mexican Heather
<i>Cupressocyparis leylandii</i>	Emerald Isle Leyland Cypress
<i>Cupressocyparis leylandii</i>	Naylor's Blue Leyland Cypress
<i>Cupressus arizonicus</i>	Arizona Cypress
<i>Cupressus arizonicus</i>	Blue Pyramid Arizona Cypress
<i>Cupressus sempervirens</i>	Italian Cypress
<i>Cupressus sempervirens</i>	Glauca Italian Cypress
<i>Cycas revoluta</i>	Sago Palm
<i>Cyperus albobstriatus</i>	Dwarf Umbrella Grass
<i>Cyrtomium fortunei</i>	Holly Fern
<i>Cytisus praecox</i>	Hollandia-Warminister Broom
<i>Cytisus scoparius</i>	Scotch Broom
<i>Cytisus spp</i>	Holandia Scotch Broom
<i>Cytisus spp</i>	Allgold Scotch Broom
<i>Dahlia</i>	Dwarf Dahlia Wendy Pink
<i>Dahlia hybrid</i>	Dwarf Dahlia
<i>Dalea greggii</i>	Trailing Indigo Bush
<i>Daphne cneorum</i>	Rose Daphne
<i>Daphne odora</i>	Fragrant Daphne
<i>Delosperma alba</i>	White Trailing Ice Plant
<i>Delosperma cooperi</i>	Ice Plant
<i>Delosperma cooperi</i>	Cooperi Pink
<i>Delosperma nubigenum</i>	Hardy Ice Plant
<i>Delphinium spp. †</i>	Larkspur; Blue Elf
<i>Descampsia caespitosa</i>	descampsia
<i>Deutzia crenata</i>	Nakiana-Dwarf Deutzia
<i>Deutzia gracilis</i>	Slender Gracilis
<i>Dianthus deltooides</i>	Zing Maiden Pinks
<i>Dianthus gratianopolitanus</i>	Firewitch Cheddar Pink
<i>Dianthus gratianopolitanus</i>	Crimson treasure Cheddar Pink
<i>Dianthus gratianopolitanus†</i>	Cheddar Pink
<i>Dianthus plumaris</i>	Cottage Pink
<i>Dietes vegeta</i>	Fortnight Lily
<i>Dodona viscosa</i>	Hopseed Bush
<i>Drosanthemum floribundum</i>	Trailing rosea iceplant
<i>Drosanthemum hispidum</i>	Ice Plant

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by "/" denotes and/or options.

Label amendment 09 29 2014

<i>Dryopteris australis</i>	Dixie Wood Fern
<i>Dryopteris erythrosora</i>	Autumn Fern
<i>Dryopteris marginalis</i>	Marginal Wood Fern
<i>Dymondia margaritae</i>	Diamond Marguerite
<i>Dyopsis lutescens</i>	Areca Palm
<i>Elaeagnus ebbengei</i>	Gild Edge Elaeagnus
<i>Elaeagnus pungens</i>	Silverberry
<i>Elaeagnus pungens</i>	Fruitland Silver-Berry
<i>Enkianthus companulatus</i>	Red Veined Enkianthus
<i>Ensete ventricosum</i>	Absynnian Banana
<i>Equisetum scirpoides</i>	Dwarf Horsetail
<i>Erianthus ravennae</i>	Hardy Pampasgrass
<i>Erica cinerea</i>	Purple Bell Heather
<i>Erica darleyensana</i>	Mediterranean Pink Heather
<i>Erica vagans</i>	Cornish Heather
<i>Erigeron speciosum</i>	Darkest of Al Fleabane
<i>Escalonia spp</i>	escallonia
<i>Eucalyptus camaldulensis</i>	Red Gum Eucalyptus
<i>Eucalyptus cinerea</i>	Silver Dollar Eucalyptus
<i>Eucalyptus microtheca</i>	Coolibah Tree
<i>Eucalyptus sideroxylon</i>	Red ironbark Eucalyptus
<i>Eugenia myritifolia</i>	Teenie Genie Brushcherry
<i>Eugenia myritifolia</i>	Dwarf Brush Cherry
<i>Euonymus kiatschovicus</i>	Mahattan Euonymus
<i>Euonymus fortunei</i>	Wintercreeper 'Blondy'
<i>Euonymus fortunei</i>	Wintercreeper
<i>Euonymus fortunei</i>	Canadale Gold Euonymus
<i>Euonymus fortunei</i>	Emerald & Gold Euonymus
<i>Euonymus fortunei</i>	Sunspot Euonymus
<i>Euonymus fortunei vegetus</i>	Bigleaf Wintercreeper
<i>Euonymus japonica</i>	Japanese Spindle Tree (Evergreen Euonymus) "gold spot"
<i>Euonymus japonica</i>	Japanese Spindle Tree
<i>Euonymus japonicus</i>	Silver King Euonymus
<i>Euonymus japonicus</i>	Chollipo Euonymus
<i>Euonymus japonicus</i>	Gold Spot Euonymus
<i>Euonymus japonicus</i>	Silver Princess Euonymus
<i>Euonymus japonicus</i>	Variegated Euonymus
<i>Euonymus kiatschovicus</i>	Spreading Euonymus
<i>Euryops pectinatus</i>	Dwarf Euryops

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Eustoma grandiflorum</i>	Pink lisianthus
<i>Evolvulus nuttallianus</i>	Blue Daze Evolvulus
<i>Fagus sylvatica</i>	European Beach
<i>Fatshedra japonica</i>	Japanese Aralia
<i>Fatsia japonica</i>	Japanese Aralia
<i>Felicia amelloides</i>	Blue Marguerite
<i>Festuca ovina</i>	Blue Fescue
<i>Ficus benjamina</i>	Indian Laurel Fig
<i>Ficus benjamina</i>	Mini Indian Laurel Fig
<i>Forshthia intermedia</i>	Border Forsythia
<i>Forsythia</i>	Spring Glory Forsythia
<i>Forsythia suspensa</i> †	Weeping Forsythia
<i>Forsythia viridissima</i>	Greenstem Forsythia
<i>Fraxinus udhei</i>	Shamel Ash
<i>Fuchsia</i>	Santa Clause Fuchsia
<i>Gardenia jasminoides</i>	Gardenia, Cape-Jasmine
<i>Gardenia jasminoides</i>	Gardenia 'Mystery'
<i>Gardenia jasminoides</i>	August Beauty Gardenia
<i>Gardenia jasminoides</i>	Miniature Gardenia
<i>Gardenia radicans</i>	Dwarf Gardenia
<i>Gaultheria procumbens</i>	Wintergreen
<i>Gaultheria shallon</i>	Salal/Lemon Leaf
<i>Gaura spp.</i> †	Guara
<i>Gazania spp</i>	Gazania
<i>Gelsemium sempervirens</i>	Carolina jessamine
<i>Genista pilosa</i>	Woadwaxen
<i>Gentiana dahurica</i> †	Gentian
<i>Geranium cinereum</i> †	Cranesbill
<i>Geranium cinerium</i>	Ballerina Cranesbill
<i>Geranium sanguineum</i>	Bloody Cranesbill
<i>Geranium subcaulescens</i>	Black Eyed Magenta Cranesbill
<i>Geum quellyon</i>	Geum
<i>Geum spp</i>	avens
<i>Ginko biloba</i>	Ginko
<i>Gleditsia triacanthos</i>	Thornless Honey Locust
<i>Gleditsia triacanthos</i>	Shademaster Honeylocust
<i>Gypsophila paniculata</i>	Babys Breath
<i>Gypsophila repens</i> †	Baby's Breath
<i>Hakonechloa macroaureolo</i>	Golden hakonechloa
<i>Hamamelis virginia</i>	Common Witch Hazel

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by "/" denotes and/or options.

Label amendment 09 29 2014

<i>Hamelia patens</i>	Firebush
<i>Hardenbergia violacea</i>	Lilac Vine
<i>Hebe buxifolia</i>	Boxleaf hebe
<i>Hedera canariensis</i>	Algerian Ivy
<i>Hedera helix</i>	English Ivy
<i>Helianthemum spp. †</i>	Sunrose
<i>Helichrysum petiolare</i>	White Licorice helichrysum
<i>Heliotropium fragrans</i>	Common Heliotrope
<i>Hemerocallis spp</i>	Daylily
<i>Hemerocallis spp</i>	Aztec Gold Daylily
<i>Hemerocallis spp</i>	Stella De Oro Daylily
<i>Hemerocallis spp</i>	Tender Love Daylily
<i>Hemerocallis spp. †</i>	Daylily: Aztec Gold, Stella De Oro, Tender Love, Mini Pearl
<i>Hesperalo parvifolia</i>	Red Yucca
<i>Heteromeles arbutiflora</i>	Toyon
<i>Heucherella spp</i>	Coral Bell
<i>Heucherella spp</i>	Bridget Bloom
<i>Heucherella spp. †</i>	Coral Bell
<i>Hibiscus rosa-sinensis</i>	Ross Estey Hibiscus
<i>Hibiscus syriacus</i>	Red Bird Rose of Sharon
<i>Hibiscus syriacus</i>	Red Heart Rose of Sharon
<i>Hibiscus syriacus</i>	Woodbridge Rose of Sharon
<i>Hibiscus Rosa-sinensis**</i>	Chinese Hibiscus**
<i>Hibiscus spp</i>	Disco Bell White Mallow
<i>Hibiscus**</i>	Rose of Sharon**
<i>Hosta</i>	Francee Plantain Lily
<i>Hosta</i>	Patriot Plantain Lily
<i>Hosta lancifolia</i>	Albo-Marginata Hosta
<i>Hosta lancifolia</i>	Narrow Leafed Plantain Lily
<i>Hosta fortunei</i>	Plantain Lily
<i>Hosta plantaginea†</i>	Hosta, Plantain Lily (Fragrant)
<i>Hosta sieboldiana†</i>	Hosta, 'Searsucker'
<i>Hosta x 'Goldstandard'</i>	Hosta 'Goldstandard'
<i>Houttuynia cordata</i>	Chameleon houttuynia
<i>Houttuynia cordata</i>	Variegated Houttuynia
<i>Huchera</i>	Bressingham Coral Bells
<i>Huchera micrantha</i>	Coral Bells
<i>Hymenoxys acaulis</i>	angelita daisy
<i>Hypericum spp</i>	St Johnswort

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by "/" denotes and/or options.

Label amendment 09 29 2014

<i>Ilex</i>	Nellie R Stevens Holly
<i>Ilex aquafolium</i>	Balkans English Holly
<i>Ilex aquafolium</i>	Gold Coast English Holly
<i>Ilex aquipernyi</i>	San Jose Holly
<i>Ilex attenuata</i>	Savannah Holly
<i>Ilex cassine</i>	Cassine Holly
<i>Ilex cornuta</i>	Burford Holly
<i>Ilex cornuta</i>	Needlepoint Holly
<i>Ilex cornuta</i>	Dwarf Burford Holly
<i>Ilex cornuta</i> **	Chinese Holly**
<i>Ilex crenata</i>	Japanese Holly
<i>Ilex crenata</i>	Compacta Holly
<i>Ilex crenata</i>	Convexa Holly
<i>Ilex crenata</i>	Dwarf Japanese Holly
<i>Ilex crenata</i>	Green Luster Holly
<i>Ilex crenata</i>	Helleri Holly
<i>Ilex crenata</i>	Hetzi Holly
<i>Ilex crenata</i>	Steeds Japanese Holly
<i>Ilex crenata</i>	Stokes Japanese Holly
<i>Ilex glabra</i>	Nordic Inkberry Holly
<i>Ilex glabra</i> 'Compacta'	Inkberry
<i>Ilex meserveae</i>	Blue Boy Holly
<i>Ilex meserveae</i>	Blue Girl Holly
<i>Ilex meserveae</i>	China Boy Holly
<i>Ilex meserveae</i>	China Girl Holly
<i>Ilex meserveae</i>	Ebony Magic Holly
<i>Ilex opaca</i>	American Holly
<i>Ilex pernyi</i>	Holly
<i>Ilex vomitoria</i>	Yaupon Holly
<i>Ilex vomitoria</i>	Dwarf Yaupon Holly
<i>Ilex vomitoria</i>	Weeping Yaupon Holly
<i>Ilex</i> x	Holly 'Mary Nell'
<i>Illicium annisatum</i>	
<i>Illicium floridanum</i>	Florida Anise Tree
<i>Impatiens walleryana</i>	Lipstick Impatiens
<i>Imperata cylindrical</i>	Japanese Blood Grass
<i>Inula ensifolia</i>	
<i>Ipomea acuminata</i>	Blue Dawn Morning Glory
<i>Iris ensata</i>	Sword-Leaved Iris, Jodlesong
<i>Iris pumila</i>	Yellow dwarf bearded Iris

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Iris siberica</i>	Cabernet Siberian Iris
<i>Iris spp.**</i>	Iris species**
<i>Iris virginica</i>	Virginia Iris
<i>Itea virginica</i>	Henry Garnet Sweetspire
<i>Ixora collinea</i>	Ixora
<i>Jasminium nudiflorum</i>	Winter Jasmine
<i>Jasminum nitidum</i>	Angelwing Jasmine
<i>Jasminum polyanthum</i>	Pink Jasmine
<i>Juglans spp.*</i>	Walnut*
<i>Juniperus chinensis</i>	Chinese Juniper
<i>Juniperus chinensis</i>	Gold Coast Juniper
<i>Juniperus chinensis</i>	Old Gold Juniper
<i>Juniperus chinensis</i>	Pfitzer Juniper
<i>Juniperus chinensis</i>	Blue Pfitzer Juniper
<i>Juniperus chinensis</i>	Sea Green Juniper
<i>Juniperus conferta</i>	Shore Juniper
<i>Juniperus conferta</i>	Emerald Sea Shore Juniper
<i>Juniperus davuricae†</i>	Parsoni
<i>Juniperus horizontalis</i>	Creeping Juniper (except “prince of wales” variety)
<i>Juniperus horizontalis</i>	Andorra Juniper
<i>Juniperus horizontalis</i>	Bar Harbor Juniper
<i>Juniperus horizontalis</i>	Blue Chip Juniper
<i>Juniperus horizontalis</i>	Blue Rug Juniper
<i>Juniperus horizontalis</i>	Creeping Juniper
<i>Juniperus horizontalis</i>	Dwarf Andorra Juniper
<i>Juniperus horizontalis</i>	Huntington Blue Juniper
<i>Juniperus horizontalis</i>	Andorra Juniper
<i>Juniperus horizontalis</i>	Blue Carpet Juniper
<i>Juniperus procumbens</i>	Dwarf Japanese Garden Juniper
<i>Juniperus prostrata</i>	Prostrate Juniper
<i>Juniperus sabina</i>	Broadmoor Juniper
<i>Juniperus sabina</i>	Hicks Juniper
<i>Juniperus sabina</i>	Savin Juniper
<i>Juniperus sabina</i>	Tam Juniper
<i>Juniperus scopulorum</i>	Emerald Green Juniper
<i>Juniperus scopulorum</i>	Wichita Blue Juniper
<i>Juniperus squamata</i>	Blue Juniper
<i>Juniperus squamata</i>	Blue Star Juniper
<i>Juniperus squamata</i>	Parson's Juniper
<i>Juniperus torulosa</i>	Hollywood Juniper

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Juniperus virginiana</i>	Eastern Red Cedar
<i>Justica brandegeana</i>	Shrimp Plant
<i>Kalmia latifolia</i>	Mountain Laurel
<i>Kniphofia uvaria</i>	Flamenco Red Hot Poker
<i>Lagerstroemia indica</i>	Crape Myrtle
<i>Lagerstroemia indica X fauriei</i>	Tuscarora Crape Myrtle
<i>Lagerstromia indica</i>	Crape Myrtle 'Burgundy Cotton'
<i>Lagerstromia indica and hybridst</i>	Crape Myrtle
<i>Lampranthus spectabilis</i>	Trailing iceplant
<i>Lantana montevidensis†</i>	Weeping Lantana
<i>Lavender spp. †</i>	Lavender; Munstead
<i>Leontopodium alpinum†</i>	Edelweiss
<i>Leptospermum scoparium</i>	Broom Teatree/Manuka
<i>Leptospermum scoparium</i>	New Zealand Tea Tree
<i>Leptospermum scoparium</i>	Ruby Glow New Zealand Tea Tree
<i>Leucophyllum frutescens</i>	Texas Sage
<i>Leucophyllum laevigatum</i>	Chihuahan Sage
<i>Leucothoe axillaris</i>	Coast leucothoe
<i>Leucothoe fontanesiana</i>	Drooping Leucothoe
<i>Liatris spicata</i>	Floristan violet gay feather
<i>Ligustrum texanum</i>	Howard's Privet
<i>Ligustrum vulgare</i>	Lodense Common Privet
<i>Ligustrum amurense</i>	Amur Privet
<i>Ligustrum japonicum</i>	Japanese Privet
<i>Ligustrum japonicum</i>	Wax Privet
<i>Ligustrum japonicum</i>	Variegated Wax Privet
<i>Ligustrum lucidum</i>	Glossy Privet
<i>Ligustrum ovalifolium</i>	California Privet
<i>Ligustrum sinense†</i>	Chinese Privet
<i>Ligustrum sinense†</i>	Variegated Chinese Privet
<i>Ligustrum texanum</i>	Wax Leaf Ligustrum
<i>Ligustrum vulgaris</i>	Cheyenne' Ligustrum
<i>Ligustrum x vicaryi</i>	Golden Privet
<i>Lilium spp</i>	Jazz Lily
<i>Lilium spp. †</i>	Lily
<i>Limonium perezii</i>	Status
<i>Liquidambar styraciflua</i>	American Sweetgum
<i>Liriope gigantea</i>	White Lily Turf
<i>Liriope gigantea</i>	Giant Lily Turf
<i>Liriope muscari</i>	Liriope; Big Blue Lillyturf

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Liriope muscari</i>	Lilac Beauty Lily Turf
<i>Liriope muscari</i>	Majestic Lily Turf
<i>Liriope muscari</i>	Monroe White Lily Turf
<i>Liriope muscari</i>	Silvery Sunproof Lily Turf
<i>Liriope muscari</i>	Variegated Lily Turf
<i>Liriope spicata</i>	Green/Creeping Lily Turf
<i>Liriope spicata</i>	Silver Dragon Lily Turf
<i>Liriope spicata</i> †	Liriope, creeping
<i>Livistona chinensis</i>	Chinese Fan Palm
<i>Lobelia cardinalis</i> †	Cardinal flower, Indian Pink
<i>Lonicera fragrantissima</i>	Winter Honeysuckle
<i>Lonicera japonica</i>	Japanese Honeysuckle
<i>Lonicera periclymenum</i>	flowering Woodbine
<i>Lonicera periclymenum</i>	Serotina Woodbine
<i>Lonicera sempervirens</i>	Trumpet Honeysuckle
<i>Lonicera tatarica</i>	Tatarian Honeysuckle
<i>Loropetalum chinense</i> †	Loropetalum 'Burgundy'
<i>Loropetalum chinense</i> †	Loropetalum 'Ruby'
<i>Loropetalum chinensis</i>	Burgandy Loropetalum
<i>Loropetalum chinensis</i>	Sizzling Pink Fringe Flower
<i>Loropetalum chinensis</i>	Razzleberri Fringe Flower
<i>Lysimachia mummularia</i>	Moneywort
<i>Lysimachia punctata</i>	Dotted Loosestrife
<i>Lythrum spp.</i> †	Loosestrife; Modern Pink
<i>Magnolia grandiflora</i>	Southern Magnolia
<i>Magnolia soulangeana</i>	Saucer Magnolia
<i>Magnolia spp.</i> **	Magnolia species**
<i>Magnolia stellata</i>	Royal Star Magnolia
<i>Mahonia aquafolium</i>	Dwarf Oregon Grape Holly/Mahonia
<i>Mahonia bealei</i>	Leather Leaf Mahonia
<i>Mahonia repens</i>	Creeping Mahonia
<i>Maleophora luteola</i>	Ice Plant
<i>Malus sargentii</i>	Non-Bearing Crabapple
<i>Malus spp.</i> *	Crabapple*
<i>Mandevilla splendens</i>	Red Riding Hood Mandevilla
<i>Matteuccia struthiopteris</i>	Ostrich Fern
<i>Matthiola incana</i>	Stock
<i>Michelia figo</i>	Banana Shrub
<i>Miscanthus sinensis</i>	Maiden Grass
<i>Miscanthus sinensis</i>	Silberfeder Miscanthus

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by "/" denotes and/or options.

Label amendment 09 29 2014

<i>Miscanthus sinensis</i> **†	Yaku Jima**, Silberfeder**
<i>Moraea iridioides</i>	African Iris
<i>Morus alba</i>	White Mulberry
<i>Musa acuminata</i>	Banana
<i>myoporum parvifolium</i>	Putah Creek
<i>Myrica cerifera</i>	Southern Waxmyrtle
<i>Myrica pennsylvanica</i>	Bayberry
<i>Nandina domestica</i>	Compact Dwarf Heavenly Bamboo
<i>Nandina domestica</i>	Harbour Dwarf Nandina
<i>Nandina domestica</i>	Nandina
<i>Nandina domestica</i>	Dwarf Purple Nandina
<i>Nandina domestica</i>	Woods Dwarf Nandina
<i>Narcissus spp.</i> **	Narcissus species**
<i>Nerium oleander</i>	Hardy Red Oleander
<i>Nerium oleander</i>	Oleander
<i>Nerium spp.</i>	Oleander
<i>Nipponanthemum nipponicum</i> †	Montauk Daisy or Nippon Daisy
<i>Oenothera missourensis</i> †	Evening Primrose, Ozark Sundrops
<i>Oenothera speciosa</i>	Siskiyou Evening Primrose
<i>Olea europaea</i> *	Olive*
<i>Onoclea sensibilis</i>	Sensitive Fern
<i>Ophiopogon japonicus</i>	Dwarf Mondo Grass
<i>Ophiopogon japonicus</i> **	Mondo Grass**
<i>Origanum libanoticum</i>	Oregano
<i>Osmanthus fortunei</i>	Fortune's Osmanthus
<i>Osmanthus fragrans</i>	Sweet Olive Osmanthus
<i>Osmanthus heterophyllus</i>	Osmanthus, False Holly
<i>Osmanthus heterophyllus</i> †	Osmanthus (False Holly): Gulf Tide
<i>Osmunda cinnamomea</i>	Cinnamon Fern
<i>Osmunda regalis</i>	Royal Fern
<i>Osteospermum fruticosum</i> †	Trailing African Daisy
<i>Oxydendrum arboreum</i>	Sourwood
<i>Pachysandra terminalis</i>	Japanese Spurge
<i>Pachysandra terminalis</i>	Green Sheen Japanese Spurge
<i>Paeonia suffruticosa</i> †	Tree Peony
<i>Parthenocissus quinquefolia</i>	Virginia Creeper
<i>Pelargonium hortorum</i>	Zonal Geranium
<i>Pelargonium peltatum</i>	Ivy Geranium
<i>Pennisetum alopecuroides</i>	Fountain Grass
<i>Pennisetum setaceum</i>	Chrimson Fountaingrass

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Pennisetum setaceum</i>	Purple Fountain Grass
<i>Pennisetum setaceum</i> **†	Fountain Grass (Dwarf)**
<i>Penstemon</i>	Apple Blossom Penstemon
<i>Pentas lanceolata</i>	Star Clusters
<i>Perovskia atriplicifolia</i> †	Russian Sage
<i>Persea americana</i> *	Avocado*
<i>Petunia hybrids</i>	Petunias
<i>Phalaris arundinacea</i>	Ribbon Grass
<i>Phlox subulata</i>	Moss Pink
<i>Phoenix roebelenii</i>	Pygmy Date Palm
<i>Phormium tenax</i>	Jack Spratt New Zealand Flax
<i>Photinia fraseri</i>	Red Tip Photinia
<i>Physocarpus opulifolius</i>	Dwarf Ninebark
<i>Physostegia virginiana</i> †	Dragonhead, False; Vivid
<i>Picea pungens</i>	Colorado Blue Spruce
<i>Picea abies</i>	Weeping Norway Spruce
<i>Picea abies</i>	Spreading Norway Spruce
<i>Picea abies</i>	Norway Spruce
<i>Picea glauca</i>	White Spruce
<i>Picea glauca</i>	Dwarf Alberta Spruce
<i>Picea pungens</i>	Hoopsi Blue Spruce
<i>Picea pungens</i>	Koster Blue Spruce
<i>Picea spp</i>	Spruces
<i>Picea spp.</i> **	Spruce species** (Colorado Blue, Norway, etc.)
<i>Pieris</i>	Forest Flame Pieris
<i>Pieris japonica</i>	Japanese Andromeda; Lily-of-the-Valley Shrub
<i>Pieris japonica</i>	Mountain of Fire Pieris
<i>Pieris japonica</i>	Snowdrift Pieris
<i>Pieris japonica</i>	Temple Bells Pieris
<i>Pieris japonica</i>	Valley Rose Pieris
<i>Pieris japonica</i>	Valley Valentine Pieris
<i>Pinus mugo</i>	Mugo Pine
<i>Pinus aristata</i>	Bristlecone Pine
<i>Pinus brutia</i>	Calabrian Pine
<i>Pinus canariensis</i>	Canary Island Pine
<i>Pinus contorta</i>	Shore Pine, Beach Pine
<i>Pinus eldarica</i>	Eldarica Pine
<i>Pinus elliotii</i>	Slash Pine
<i>Pinus halapensis</i>	Aleppo Pine
<i>Pinus leucodermis</i>	Bosnian Pine

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Pinus nigra</i>	Austrian Black Pine
<i>Pinus palustris</i>	Longleaf Pine
<i>Pinus radiata</i>	Monterey Pine
<i>Pinus strobus</i>	Eastern White Pine
<i>Pinus sylvestris</i>	Scotch Pine
<i>Pinus sylvestris</i>	Columnar Scotch Pine
<i>Pinus taeda</i>	Loblolly Pine
<i>Pinus thunbergii</i>	Japanese Black Pine
<i>Pinus virginiana</i>	Virginia Pine
<i>Pistacia spp.*</i>	Pistachio*
<i>Pittosporum rhombifolium</i>	Queensland Pittosporum
<i>Pittosporum tenuifolia</i>	Golf Ball Pittosporum
<i>Pittosporum tobira</i>	Japanese Pittosporum
<i>Pittosporum tobira</i>	Green Pittosporum
<i>Pittosporum tobira</i>	Wheeler's Dwarf Pittosporum
<i>Platanus occidentalis</i>	American Sycamore
<i>Platanus racemosa</i>	California Sycamore
<i>Plumbago auriculata</i>	Plumbago
<i>Plumbago capensis</i>	Plumbago
<i>Podocarpus macrophyllus</i>	Yew Pine/Japanese Yew
<i>Podocarpus Species</i>	Podocarpus
<i>Polygala fructicosa</i>	Sweet Pea Shrub
<i>Polystichum acrostichoides</i>	Christmas Fern
<i>Polystichum polyblepharum</i>	Tassel Fern
<i>Populus deltoides</i>	Cottonwood
<i>Potentilla fragiformis</i>	Cinquefoil
<i>Potentilla fruticosa</i>	Cinquefoil
<i>Potentilla fruticosa</i>	Floppy Disc Cinquefoil
<i>Potentilla fruticosa</i>	Gold Drop Cinquefoil
<i>Potentilla fruticosa</i>	Goldfinger Cinquefoil
<i>Potentilla fruticosa</i>	Red Ace Cinquefoil
<i>Potentilla fruticosa</i>	Sunset Cinquefoil
<i>Potentilla fruticosa</i>	Tangerine Cinquefoil
<i>Potentilla spp</i>	Cinquefoil
<i>Potentilla verna</i>	Spring Cinquefoil
<i>Prosopis chiliensis chiliensis</i>	Chilean Mesquite
<i>Prunus laurocerasus</i>	English Laurel
<i>Prunus laurocerasus</i>	Otto Luyken English Laurel
<i>Prunus caroliniana</i>	Carolina Cherry Laurel
<i>Prunus caroliniana</i>	Bright 'n Tight Cherry Laurel

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by "/" denotes and/or options.

Label amendment 09 29 2014

<i>Prunus glandulosa</i>	Dwarf Flowering Almond
<i>Prunus spp.*</i>	Almond, Apricot, Nectarine, Peach, Plum and Prune*
<i>Prunus x yedoensis</i>	Yoshino Cherry
<i>Pseudotsuga menziesii**</i>	Douglas Fir**
<i>Psidium cattleianum</i>	Strawberry Guava
<i>Pyracantha coccinea</i>	Firethorn Scarlet
<i>Pyracantha coccinea</i>	Lalandei Firethorn
<i>Pyracantha fortuneana</i>	Firethorn
<i>Pyracantha fortuneana</i>	Lolendei Monrovia Pyracantha
<i>Pyracantha fortuneana</i>	Monon Pyracantha
<i>Pyracantha fortuneana</i>	Red Elf Pyracantha
<i>Pyracantha fortuneana</i>	Rutgers Hybrid Pyracantha
<i>Pyracantha fortuneana</i>	Santa Cruz Pyracantha
<i>Pyracantha fortuneana</i>	Victory Pyracantha
<i>Pyracantha koidzumii</i>	Pyracantha 'Santa Cruz'
<i>Pyracantha koidzumii</i>	Firethorn
<i>Pyrus spp</i>	Bradford Pears
<i>Pyrus spp.</i>	Pear spp. including Bradford
<i>Quercus ilicifolia</i>	Bear Oak
<i>Quercus palustris</i>	Pin Oak
<i>Quercus phellos</i>	Willow Oak
<i>Quercus rubra</i>	Red Oak
<i>Quercus shumardii†</i>	Oak, Shumard's Red
<i>Raphiolepis indica</i>	Indian Hawthorne
<i>Raphiolepis umbellata†</i>	Yedda Hawthorne
<i>Ratbida columnifera</i>	Mexican Hat
<i>Ravenea rivularis</i>	Majesty Palm
<i>Rhaphiolepis indica</i>	Charisma-monruce Indian Hawthorn
<i>Rhaphiolepis indica</i>	Enchantress-moneess Indian Hawthorn
<i>Rhaphiolepis indica</i>	Indian Hawthorn
<i>Rhaphiolepis indica</i>	Springtime Indian Hawthorn
<i>Rhaphiolepis indica</i>	Ballerina Indian Hawthorn
<i>Rhaphiolepis ovata</i>	Round-Leaf Indian Hawthorn
<i>Rhododendron</i>	Fashion Azalea
<i>Rhododendron</i>	Wakeiebisu Azalea
<i>Rhododendron</i>	George Tabor Azalea
<i>Rhododendron</i>	Gerard Azalea
<i>Rhododendron</i>	Delaware Valley White Azalea
<i>Rhododendron</i>	Girard Roberta Azalea
<i>Rhododendron</i>	Girard Crimson Azalea

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Rhododendron</i>	Golden Flare Exbury Azalea
<i>Rhododendron</i>	Helmut Vogel Azalea
<i>Rhododendron</i>	Hershey Red Azalea
<i>Rhododendron</i>	Hot Shot Azalea
<i>Rhododendron</i>	Hume Azalea
<i>Rhododendron</i>	Inga Azalea
<i>Rhododendron</i>	Irene Koster Azalea
<i>Rhododendron</i>	President Clay Azalea
<i>Rhododendron</i>	Traditional Azalea
<i>Rhododendron forrestii</i> x <i>griersonianum</i>	Elizabeth Rhododendron
<i>Rhododendron calendulaceum</i>	Flame Azalea
<i>Rhododendron calendulaceum</i>	Golden Fire Azalea
<i>Rhododendron calendulaceum</i>	Klondike Azalea
<i>Rhododendron campylocarpum</i>	Butterfly Rhododendron
<i>Rhododendron catawbiense</i>	White Catawba Rhododendron
<i>Rhododendron catawbiense</i>	Catawba Rhododendron
<i>Rhododendron catawbiense</i>	Lord Roberts Catawba Rhododendron
<i>Rhododendron catawbiense</i>	Rocket Catawba Rhododendron
<i>Rhododendron caucasicum</i> X <i>ponticum</i>	Cunningham White Rhododendron
<i>Rhododendron exbury</i>	Canon's Double Exbury Azalea
<i>Rhododendron exbury</i>	Golden Flare Exbury Azalea
<i>Rhododendron exbury</i>	Klondike Exbury Azalea
<i>Rhododendron forrestii</i>	Gomer Waterer Rhododendron
<i>Rhododendron griffithianum</i>	Jean Marie Rhododendron
<i>Rhododendron hybrid spp</i>	America Rhododendron
<i>Rhododendron hybrid spp</i>	Nova Zembla Rhododendron
<i>Rhododendron hybrid spp</i>	Scintillation Rhododendron
<i>Rhododendron Hybrids</i>	Carror Azalea
<i>Rhododendron impeditum</i>	Rhododendron
<i>Rhododendron indica</i>	Formosa Azalea
<i>Rhododendron indica</i>	Waucabusa Azalea
<i>Rhododendron kaempferi</i>	Blue Danube Azalea
<i>Rhododendron kerume</i>	Coral Bells Azalea
<i>Rhododendron kerume</i>	Hino Crimson Azalea
<i>Rhododendron kerume</i>	Hino Pink Azalea
<i>Rhododendron kerume</i>	Mildred Azalea
<i>Rhododendron kerume</i>	Snow Azalea
<i>Rhododendron maximum</i>	Rose Bay Rhododendron
<i>Rhododendron mucronulatum</i>	Rhododendron
<i>Rhododendron obtusum</i>	Hino Crimson Azalea

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Rhododendron ponticum</i>	Chioniodes Rhododendron
<i>Rhododendron ponticum</i>	Daphinoides Rhododendron
<i>Rhododendron racemosum</i>	Dwarf Scarlet Wonder Rhododendron
<i>Rhododendron racemosum</i>	Tribly Rhododendron
<i>Rhododendron racemosum</i>	Unique Rhododendron
<i>Rhododendron racemosum</i>	Vulcan Rhododendron
<i>Rhododendron sassthigiatum X carolinianum</i>	Ramapo Rhododendron
<i>Rhododendron satuski</i>	Gumpo Pink Azalea
<i>Rhododendron satuski</i>	Gumpo White Azalea
<i>Rhododendron satuski</i>	Higasa Azalea
<i>Rhododendron satuski</i>	Reijn Azalea
<i>Rhododendron schlippenbachii</i>	Royal Azalea
<i>Rhododendron spp. †</i>	Rhododendrons, Azaleas (Except Carolina and roseum elegans variety rhododendron)
<i>Rhododendron x 'George Tabor'</i>	Azalea 'George Tabor'
<i>Rhododendron x obtusum</i>	Kurume Azalea 'Tradition'
<i>Rhus lancea</i>	African Sumac
<i>Rhus typhina</i>	Staghorn Sumac
<i>Rosa banksiae</i>	Lady Bank's Rose
<i>Rosa Flower Carpet</i>	Red Groundcover Rose
<i>Rosa rugosa</i>	Rugosa Rose
<i>Rosa x</i>	Knock Out Rose Radrazz
<i>Rosa x</i>	Knock Out Double Pink
<i>Rosmarinus officinalis</i>	Rosemary
<i>Rudbeckia fulgida</i>	Black-Eyed Susan 'Early Bird'
<i>Rudbeckia fulgida</i>	Blackeyed Susan
<i>Rudbeckia fulgida †</i>	Black-Eyed Susan 'Goldstrum'
<i>Rumohra adiantiformis</i>	Leatherleaf Fern
<i>Ruscus hypophyllum</i>	Butchers Broom
<i>Salix babylonica</i>	Babylon Weeping Willow
<i>Salix matsudana</i>	Corkscrew Willow
<i>Salvia daghestancia</i>	Platinum Sage
<i>Salvia grahamii</i>	Grahams Sage
<i>Santolina virens</i>	Green Lavender Cotton
<i>Santolina virens</i>	Green Santolina
<i>Sasa pygmaea</i>	Dwarf Bamboo
<i>Saxifraga spp. †</i>	Saxifrage; Purple Dome
<i>Scabiosa spp. †</i>	Pincushion flower
<i>Scutellaria resinosa</i>	Skull Cap
<i>Sedum pachyclados</i>	Grey Stonecrop

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Sedum spurium</i>	Sedum 'John Creech'
<i>Senecio cineraria</i>	Dusty Miller
<i>Sequoia sempervirens</i>	Coast Redwood
<i>Sequoiadendron giganteum</i>	Giant Sequoia
<i>Skimmia japonica</i>	Japanese skimmia
<i>Skimmia revesiana</i>	Reeves Skimmia
<i>Solanum rantonetii</i>	Paraguay Nightshade
<i>Spiraea bumalda</i> †	Spiraea 'Anthony Waterer'
<i>Spiraea cinerea</i>	First Snow Spiraea
<i>Spiraea japonica</i>	Dolchia Spiraea
<i>Spiraea japonica</i>	Japanese Alpine Spiraea
<i>Spiraea japonica</i>	Magic Carpet Spiraea
<i>Spiraea japonica</i>	Neon Flash Spiraea
<i>Spiraea japonica</i>	Shirobana Spiraea
<i>Spiraea vanhouttei</i>	Vanhouttei Spiraea
<i>Spiraea japonica</i>	Spiraea 'Neon Flash'
<i>Spiraea japonica</i>	Spiraea 'Shirobana'
<i>Streptosolen jamesonii</i>	Marmalade Bush
<i>Swietenia mahogani</i>	Mahogany
<i>Syagrus romanzoffianum</i>	Queen Palm
<i>Syringa rothomagensis</i>	Chinese Lilac
<i>Syringa vulgaris</i>	Common Lilac
<i>Syzygium paniculatum</i>	Japanese Boxcherry
<i>Syzygium paniculatum</i>	Australian Brushcherry
<i>Tabebuia caraiba</i>	Yellow Tabebuia
<i>Tagetes patula</i>	Little Hero Marigold
<i>Taxus cuspidata</i>	Japanese Yew
<i>Taxus media</i> †	Yew
<i>Tecomaria capensis</i>	Cape Honeysuckle
<i>Ternstroemia gymnanthera</i>	Japanese Ternstroemia
<i>Teucrium fruiticans</i>	Bush Germander
<i>Teucrium spp</i> †.	Germander
<i>Thalictrum dipterocarpum</i> †	Meadow Rue
<i>Thevetia nerifolia</i>	Yellow Oleander
<i>Thuja occidentalis</i>	American Arborvitae
<i>Thuja occidentalis</i>	Emerald Arborvitae
<i>Thuja occidentalis</i>	Globe Arborvitae
<i>Thuja occidentalis</i>	Little Giant Dwarf Arborvitae
<i>Thuja occidentalis</i>	Dark American Arborvitae
<i>Thuja occidentalis</i>	Pyramidal Arborvitae

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Thuja occidentalis</i>	Rheingold Arborvitae
<i>Thuja occidentalis</i>	Techny Arborvitae
<i>Thuja occidentalis</i>	Woodward's Arborvitae
<i>Thuja orientalis</i>	Dwarf Golden Arborvitae
<i>Thuja orientalis</i>	Dwarf Arborvitae
<i>Tibouchina urvilleana</i>	Princess Flower
<i>Trachelospermum asiaticum</i>	Asiatic Jasmine, Star Jasmine
<i>Trachycarpus fortunei</i>	Windmill Palm
<i>Tsuga canadensis</i>	Canada Hemlock, Eastern Hemlock
<i>Tulbaghia violacea</i>	Society Garlic
<i>Tulipa spp</i>	Tulips
<i>Ulmus parvifolia</i>	Chinese Elm
<i>Verbena rigida</i>	Veined Verbena
<i>Veronica spp</i>	Sunny Border, Veronica /Speedwell
<i>Veronica spp. †</i>	Veronica, Speedwell
<i>Viburnum bodnantense</i>	Pink Dawn Viburnum
<i>Viburnum carlesii</i>	Koreanspice Viburnum
<i>Viburnum davidii</i>	David Viburnum
<i>Viburnum japonicum</i>	Japanese Viburnum
<i>Viburnum judd</i>	Judd Viburnum
<i>Viburnum lantana</i>	Wayfaringtree Viburnum
<i>Viburnum odoratissimum</i>	Sweet Viburnum
<i>Viburnum opulus 'Roseum'</i>	Roseum Viburnum
<i>Viburnum opulus sterile</i>	Common Snowball Viburnum
<i>Viburnum plicatum</i>	Japanese Snowball
<i>Viburnum plicatum</i>	Doublefile Viburnum
<i>Viburnum pragense</i>	Prague Viburnum
<i>Viburnum rigidum</i>	Canary Island Viburnum
<i>Viburnum setigerum</i>	Tea Viburnum
<i>Viburnum suspensum†</i>	Arrowwood viburnum
<i>Viburnum tinus</i>	Laurustinus
<i>Viburnum tinus</i>	Spring Bouquet Viburnum
<i>Viburnum trilobium</i>	Cranberry Bush
<i>Viburnum trilobum</i>	American Cranberrybush Viburnum
<i>Viburnum trilobum</i>	Dwarf American Cranberrybush Viburnum
<i>Viburnum wrightii</i>	Leatherleaf Viburnum
<i>Vinca major</i>	Vinca
<i>Vinca major</i>	Bigleaf Periwinkle
<i>Vinca minor</i>	Dwarf Periwinkle
<i>Vinca spp</i>	Periwinkle

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

<i>Vitis</i> spp.*	Grape*
<i>Washingtonia robusta</i>	Mexican Fan Palm
<i>Washingtonia robusta</i>	California Fan Palm
<i>Weigelia florida</i>	Old Fashioned Weigela
<i>Weigelia florida</i>	Bristol Ruby Weigelia
<i>Weigelia florida</i>	Java Red Weigelia
<i>Weigelia florida</i>	Minuet Weigelia
<i>Xylosma congestum</i>	Xylosma
<i>Xylosma senticoso</i>	Shiny Xylosma
<i>Yucca aloifolia</i>	Spanish Bayonet
<i>Yucca filamentosa</i>	Yucca, Adam's Needle
<i>Zamia furfuracea</i>	Cardboard Palm

† Not for use in California

*Do not use on food producing trees, vines, or plants

**Not for use on container grown plants.

Table 13. Do not use this product for weed control in the following ornamental plant species:

Scientific Name	Common Name
<i>Ajuga</i> spp.	Bugleweed or ajuga
<i>Brassica</i> spp.	mustard
<i>Echinacea purpurea</i>	purple coneflower
<i>Euonymus alatus 'Compacta'</i>	dwarf burning bush
<i>Euphorbia</i> spp.	spurge
<i>Hydrangea</i> spp.	hydrangea
<i>Iberis</i> spp.	candytuft
<i>Juniperus horizontalis 'Prince of Wales'</i>	Prince of Wales juniper
<i>Melaleuca quinquenervia</i>	cajeput tree
<i>Rhododendron caroliniaum</i>	Carolina rhododendron
<i>Rhododendron catawbiense 'Roseum elegans'</i>	roseum elegans rhododendron
<i>Sedum</i> spp.	stonecrop
<i>Yucca recurvifolia</i>	green yucca

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

VEGETATION MANAGEMENT

Product Information

Apply Quali-Pro[®] Prodoxaben 3.7 SC to soil surfaces for preemergence control of many grass and broadleaf weeds in non-crop areas including ornamentals (does not include container or field grown ornamentals) or surrounding non-crop areas such as managed rights-of-way for transportation systems and utilities (including roadways, roadsides, railways, and equipment yards) and facilities including substations, tank farms, pumping stations, parking and storage areas, and non-grazed fence rows. The best weed control occurs when the product is activated in the soil by 0.5 inch of irrigation or rainfall before weed seeds germinate and within 14 days after application.

Timing and Method of Application

Quali-Pro[®] Prodoxaben 3.7 SC is a preemergence herbicide only and will not control emerged grass or broadleaf weeds. It works best when applied to soil that is free of clods, weeds and debris and is activated by at least 0.5 inches of irrigation or rainfall or shallow incorporation to a depth of 1 to 2 inches. Weed control is improved if Quali-Pro[®] Prodoxaben 3.7 SC is activated in the soil before weed seeds germinate and within 14 days after application. Apply Quali-Pro[®] Prodoxaben 3.7 SC to newly-transplanted and established ornamentals as a broadcast or over-the-top spray.

Plant Sensitivity

Direct application of Quali-Pro[®] Prodoxaben 3.7 SC to rapidly growing tissue or buds can injure desirable plants. In the spring when buds are rapidly growing and expanding, over-the-top application of Quali-Pro[®] Prodoxaben 3.7 SC can injure new growth of desirable plants, however, these effects are temporary. To reduce the possibility of injury at this time, wait to apply Quali-Pro[®] Prodoxaben 3.7 SC over the top of newly emerged vegetation until it has hardened off, unless your experience indicates that the ornamental plant will not be injured by the over-the-top application. After application, immediately apply overhead irrigation to the foliage to wash Quali-Pro[®] Prodoxaben 3.7 SC from plant surfaces onto soil (watering the foliage of plants before application can improve the washing process

Tank Mixtures for Use in Vegetation Management

Tank mix Quali-Pro[®] Prodoxaben 3.7 SC with other registered herbicides listed on this label to provide a broader spectrum of weed control or to control emerged weeds. Tank mixes with Quali-Pro[®] Prodoxaben 3.7 SC are for use only in states where the tank-mix partner(s), application site and intended use pattern are registered.

Follow the label directions of the tank-mix partner(s) for application rates, timing, weeds controlled, listed ornamentals, and specific use precautions and/or restrictions. Before combining a tank-mix partner in the spray tank, test for compatibility as described on this label.

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

Table 14. Tank-Mix Partners for Quali-Pro® Prodoxaben 3.7 SC Use in Vegetation Management

Product	Precautions/Instructions
Galigan 2E®, (use on conifers only)	<ul style="list-style-type: none"> Mix with Quali-Pro® Prodoxaben 3.7 SC for postemergence control of certain broadleaf weeds including malva and filaree.
Quali-Pro Isoxaben 75 WG, Princep®, MANA Parallel	<ul style="list-style-type: none"> See product labels for weed spectrum and listed ornamentals.
Glyphogan® (or other glyphosate-based products) and Finale®	<ul style="list-style-type: none"> These nonselective tank mix herbicides control many emerged annual broadleaves and grasses. Take extreme care to prevent tank mixtures with these partner products from contacting the foliage and stems of turfgrass, trees, shrubs, or other desirable vegetation because desirable vegetation might be severely injured or killed. Apply these tank mixtures as a directed spray and use a shield to prevent spray from contacting foliage of desirable plants. Following instructions on the tank-mix partner's label, delay irrigation of the treated area to allow time for the herbicide to be absorbed by weed foliage.

Sensitivity of Ornamental Species - Vegetation Management

Quali-Pro® Prodoxaben 3.7 SC will not likely harm the trees, shrubs, vines and flowers listed above in **Table 11**. Quali-Pro® Prodoxaben 3.7 SC can be applied over the top of the listed species.

When plants are under stress (such as heat, drought, or frost damage), some cultivars of listed plants may be sensitive to Quali-Pro® Prodoxaben 3.7 SC.

VEGETATION MANAGEMENT - NON-CROP AREAS

Product Information

Apply Quali-Pro® Prodoxaben 3.7 SC to soil surfaces for preemergence control of many grass and broadleaf weeds in non-crop areas including ornamentals (does not include container or field grown ornamentals) and established perennials on or surrounding non-crop areas as directed in **Table 14**. The best weed control occurs when the product is activated in the soil by 0.5 inches of irrigation or rainfall before weed seeds germinate and within 14 days after application.

Table 15. Amount of Quali-Pro® Prodoxaben 3.7 SC to Apply to Non-Crop Areas for Control of Weeds Listed in Table 2.

Site	Application Rate	Timing	Comments/Instructions
Non-crop areas, including ornamentals, on or surrounding managed rights-of-way for transportation, systems and utilities (including roadways, roadsides, railways, and	38 – 87 fl. oz./acre or 0.9- 2.0 fl. oz./1,000 sq ft	Before weed seeds germinate	Use higher rate range for longer control. More than one application per year is permitted, but do

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

equipment yards); Facilities including substations, tank farms, pumping stations, parking and storage areas, and non-grazed fence rows.			not apply more than 87 fl. oz./acre (2.5 lbs. active/acre) per year. Sequential applications must be separated by a minimum of 60 days.
--	--	--	--

Tank Mixtures for Use in Non-Crop Areas

Tank mix Quali-Pro® Prodoxaben 3.7 SC with other registered herbicides listed on this label to provide a broader spectrum of weed control or to control emerged weeds. Tank mixes with Quali-Pro® Prodoxaben 3.7 SC are for use only in states where the tank-mix partner(s), application site and intended use pattern are registered.

Follow the label directions of the tank-mix partner(s) for application rates, timing, weeds controlled, listed ornamentals, and specific use precautions and/or restrictions. Before combining a tank-mix partner in the spray tank, test for compatibility as described on this label. See the **Compatibility Testing** section of this label.

For non-crop areas, tank mix Quali-Pro® Prodoxaben 3.7 SC with Glyphogan® (or other glyphosate-based products), Parazone 3SL® (or other paraquat-based products), Diuron 4L (or other diuron-based products), Oryzalin 4AS (or other oryzalin based products), Reward®, Predict®, Princep®, Vanquish®, Finale®, Garlon®, Krovar® I and II, Oust®, Arsenal®, Spike® and Telar®.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal or cleaning of equipment.

PESTICIDE STORAGE: Store in original container in dry area. Do not store in a manner where cross-contamination with other pesticides, fertilizers, food or feed could occur. If spilled during storage or handling, absorb with sand or other inert material and dispose of absorbent in accordance with the Pesticide Disposal instructions listed below.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable Container [(five gallons or less)]: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. If recycling is not available, puncture and dispose of in a sanitary landfill or by incineration or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by “/” denotes and/or options.

Label amendment 09 29 2014

LIMITATION OF WARRANTY AND LIABILITY

Read the entire direction for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following CONDITIONS, DISCLAIMER OF WARRANTIES, and LIMITATIONS OF LIABILITY.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Control Solutions, Inc. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, Control Solutions, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Control Solutions, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Control Solutions, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Control Solutions, Inc election, the replacement of product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Control Solutions, Inc.'s election, the replacement of product.

Quali-Pro® is a registered trademark of Makhteshim Agan of North America, Inc.

[If you have questions regarding the use of this product, call TOLL FREE x-xxx-xxx-xxxx]

[Gemini® is a registered trademark of Everris International B.V.]
