



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

August 22, 2024

Haley Wade
Lead Regulatory Manager
Control Solutions, Inc.
5903 Genoa-Red Bluff
Pasadena, TX 77507-1041

Subject: Label Amendment - Registration Review Mitigation for Oxadiazon
Product Name: QUALI-PRO OXADIAZON 2G
EPA Registration Number: 53883-372
Application Date: June 3, 2022
Decision Number: 585154

Dear Haley Wade:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Oxadiazon Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable. As part of the registration review, EPA identified the need for this product to be classified as Restricted Use. The labeling approved today incorporates the necessary changes for classification as Restricted Use, therefore the classification is effective on the date of this letter.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. Due to the classification of this product as Restricted Use, in accordance with 40 CFR 152.167, after 120 days, any of this product you sell or distribute must:

- 1) Bear an approved amended label containing the terms of the Restricted Use classification;
- 2) Bear a sticker containing the product name, EPA registration number, and any terms of restricted use; **or**
- 3) Be accompanied by supplemental labeling bearing the product name, EPA registration numbers, and any terms of restricted use.

After 270 days, any of this product you sell or distribute must bear the approved amended labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Theodore Varns by phone at 202-566-2241, or via email at varns.theodore@epa.gov.

Sincerely,

A handwritten signature in dark ink, appearing to read "Kevin Costello", is positioned above the typed name and title.

Kevin Costello, Branch Chief
Risk Management and Implementation Branch 2
Pesticide Re-Evaluation Division
Office of Pesticide Programs

ENCLOSURE: Stamped label

RESTRICTED USE PESTICIDE

Due to potential human health risks of concern (cancer) from chronic dietary or dermal exposure when used in accordance with label directions.

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

OXADIAZON GROUP 14 HERBICIDE

Quali-Pro® Oxadiazon 2G

For sale to and use by professional applicators only. Not for sale to or use by homeowners/consumers. Not for use on home lawns.

Quali-Pro Oxadiazon 2G is a selective, preemergence herbicide for control of annual grasses and broadleaf weeds in turf and woody ornamental shrubs, vines, and trees.

ACTIVE INGREDIENT:

Oxadiazon [2-tert-butyl-4-(2,4 dichloro-5-isopropoxyphenyl)-2-1, 3,4-oxadiazoline-5-one]:.....2.0%

OTHER INGREDIENTS:.....98.0%

TOTAL:100.0%

[Optional Text – Guaranteed Fertilizer Analysis]

EPA Reg. No. 53883-372

EPA Est. No.

Manufactured for:

Control Solutions, Inc.
5903 Genoa Red Bluff Rd.
Pasadena, TX 77507

Net Weight: ____lbs.

KEEP OUT OF REACH OF CHILDREN

WARNING – AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail).

FIRST AID	
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall 1-866-897-8050 for emergency medical treatment information.	

ACCEPTED

Aug 22, 2024

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 53883-372

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING**

Causes substantial but temporary eye injury. DO NOT get in eyes, on skin, or on clothing. Wear protective eyewear such as goggles, face shield, or safety glasses.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Waterproof gloves

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

Runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

This pesticide is toxic to oysters and shrimp.

This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water.

This product is classified as having high potential for reaching surface water via runoff for several months or more after application.

A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of oxadiazon from runoff water and sediment.

NON-TARGET ORGANISM ADVISORY: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize drift.

DIRECTIONS FOR USE

Restricted Use Pesticide

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval, and notification to workers (as applicable). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the worker protection standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Shoes plus socks, and
- Waterproof gloves

WEED RESISTANCE MANAGEMENT

For resistance management, Oxadiazon 2G is a Group 14 herbicide. Any weed population may contain or develop plants naturally resistant to Oxadiazon 2G and other Group 14 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of Oxadiazon 2G or other Group 14 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.

- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact your local Control Solutions Inc. distributor or sales representative.

PRODUCT INFORMATION

Quali-Pro® Oxadiazon 2G is a selective, pre-emergent herbicide for control of annual grasses and many broadleaf weeds in commercial turf and on a wide variety of woody ornamental shrubs, vines, and trees. Quali-Pro Oxadiazon 2G controls weeds by killing the young weed seedlings as they come in contact with the herbicide during germination.

After application, user must thoroughly irrigate the site as soon as possible on the same day of application.

State Specific Restrictions:

- The state of Arizona has not approved this product for use on agricultural sites. Do not use this product on uses considered by the Arizona statutes to be agricultural uses.

For all use sites:

- The minimum retreatment interval (RTI) between applications is 60 days.
- Total applications of oxadiazon of all physical forms must not exceed 6.0 lbs a.i./A/Year if this product is used on the same site with liquid formulations.

For oxadiazon use on turf:

- A maximum single application rate of 3.0 lbs a.i./A, and a maximum annual application rate of 6.0 lbs a.i./A.
- For applications on non-agricultural turf (e.g., golf courses, parks, athletic fields) that follow a liquid application, Quali-Pro® Oxadiazon 2G must be applied 60 days or later after the liquid application. After applying this product, no liquid applications of oxadiazon are allowed within the same calendar year on the same area of turf.
- For applications to non-agricultural turf (e.g., golf courses, parks, athletic fields), this product may not be applied prior to any liquid applications of oxadiazon within the same calendar year on the same area of turf.
- On golf course turf only: In areas of heavy weed infestation only, the maximum single application rate is 4.0 pounds of active ingredient per acre, and the maximum annual application rate is 8.0 pounds of active ingredient per acre. No liquid applications may be applied on areas of turf treated at this maximum application rate in the same year.
- On golf course turf only: total applications of oxadiazon of all physical forms are limited to 30% of the total managed turf area of a golf course per year. Applications may be made to any turf surface type, including fairways, tees, greens, roughs, non-playable turf surfaces, or other managed turf areas. This 30% limit may be applied to the same or different turf areas from year to year.
- On golf course turf and sod farms: a minimum 10-foot-wide untreated vegetative buffer must be maintained between the treated area and any surface water features.

For ornamental landscaping, trees, shrubs, and nurseries:

- A maximum single application rate of 3.0 lbs a.i./A, and a maximum annual application rate of 6.0 lbs a.i./A.

Do not apply more than 300 lbs. Quali-Pro Oxadiazon 2G per acre per year (equivalent to 6 lbs. a.i./A/year). In areas of heavy weed infestation on golf course turf only, the maximum application rate is 400 lbs. Quali-Pro Oxadiazon 2G per acre per year (equivalent to 8 lbs. a.i./A/year).

COMMERCIAL TURF

For use in established perennial Bluegrass, Bermudagrass, perennial Ryegrass, St. Augustinegrass, Tall Fescue, and Zoysia turf; such as fairways, parks, golf courses, and lawns. Quali-Pro Oxadiazon 2G controls Crabgrass (*digitaria spp.*), Annual Bluegrass (*Poa annua*), Goosegrass (*Eleusine indica*); and it also controls Florida Pusley (*Richardia scabra*), Oxalis (*Oxalis spp.*), Stinging Nettle (*Urtica dioica*), Carpetgrass (*Axonopus offinis*), and Pigweed (*Amaranthus spp.*).

Apply to dry foliage only with a properly calibrated granular applicator at recommended rate in early spring before weed seed germination. If rain is not expected shortly after application, irrigation to remove herbicide granules from foliage will increase effectiveness.

For Crabgrass, Goosegrass and Crowfoot control, apply 100-150 pounds Quali-Pro Oxadiazon 2G per acre (2 to 3 pounds active ingredient) or 2 ¼ to 3 3/8 pounds per 1000 sq. ft. Late applications are not recommended and may not be fully effective.

For Annual Bluegrass control apply 150 pounds/acre Quali-Pro Oxadiazon 2G (3 pounds active ingredient) or 3 3/8 pounds/1000 sq. ft. in early spring or late summer prior to seed germination.

For Pigweed, Florida Pusley, Oxalis, Stinging Nettle and Carpetgrass control, apply 150 pounds/acre Quali-Pro Oxadiazon 2G (3 pounds active ingredient) or 3 3/8 pounds per 1000 sq. ft. in early spring prior to weed seed germination.

For St. Augustinegrass turf, do not exceed the rate of 150 pounds/acre (3 pounds active ingredient) of Quali-Pro Oxadiazon 2G.

Discoloration of Bermudagrass and St. Augustinegrass may result from use of this herbicide, but this discoloration is temporary and normally outgrown within 2 to 3 weeks.

Red Fescue and Bentgrass are susceptible to Quali-Pro Oxadiazon 2G and may be severely damaged. Avoid use of this product on turf containing Red Fescue or Bentgrass except where they are undesirable grasses. Do not apply to newly seeded areas. Seeding into treated areas should be delayed until 4 months after treatment. Do not apply to Dichondra or Centipedegrass. Quali-Pro Oxadiazon 2G should not be used on putting greens or tees because of varying cultural practices used.

WOODY ORNAMENTALS SHRUBS, VINES, AND TREES

APPLICATION:

Apply Quali-Pro Oxadiazon 2G anytime during the year, prior to weed seed germination, for weed control. Quali-Pro Oxadiazon 2G can be applied to both newly transplanted and established ornamentals. Application can be made to actively growing or dormant ornamentals. Rainfall or overhead irrigation after application of Quali-Pro Oxadiazon 2G will improve weed control activity. Quali-Pro Oxadiazon 2G should not be mixed into the soil. Cultivating or disturbing the soil may reduce the weed control activity of Quali-Pro Oxadiazon 2G. Do not apply to wet foliage or under conditions in which granules will collect on leaves.

AMOUNT TO USE:

Ground Application- Apply Quali-Pro Oxadiazon 2G broadcast to ornamentals at 100 to 150 pounds of product per acre (or 2 ¼ to 3 3/8 pounds per 1000 sq. ft.) depending on the weeds to be controlled, as listed below. Remove existing weed growth before application.

Quali-Pro Oxadiazon 2G at 100 to 150 pounds (2.0 to 3.0 pounds of active ingredient) per acre is effective in controlling annual grasses and broad-leaf weeds such as:

Annual Broadleaf Weeds	
Bittercress	<i>Cardamine</i> spp.
Carpetweed	<i>Mollugo verticillata</i>
Common Groundsel	<i>Senecio vulgaris</i>
Common Purslane	<i>Portulaca oleracea</i>
*Eveningprimrose	<i>Oenothera</i> spp.
*Galinsoga	<i>Galinsoga</i> spp.
Golden Ragwort	<i>Senecio aureus</i>
Lambsquarters	<i>Chenopodium album</i>
Pennsylvania Smartweed	<i>Polygonum pensylvanicum</i>
Petty Spurge	<i>Euphorbia peplus</i>
Redroot Pigweed	<i>Amaranthus Retroflexus</i>
Speedwell	<i>Veronica</i> spp.
Spotted Catsear	<i>Hypochaeris radicata</i>
*Swinecress	<i>Coronopus didymus</i>
Yellow Woodsorrel	<i>Oxalis stricta</i>
Annual Grasses	
Annual Bluegrass	<i>Poa annua</i>
*Barnyardgrass	<i>Echinochloa crus-galli</i>
Crabgrass	<i>Digitaria</i> spp.
Fall Panicum	<i>Panicum dichotomiflorum</i>
Green Foxtail	<i>Setaria viridis</i>

*Use 150 pounds Quali-Pro Oxadiazon 2G (3.0 pounds of active ingredient) per acre for less susceptible weeds.

Note: For the control of liverwort (*Marchantia* spp.) in California, Goosegrass (*Eleusine Indica*) and Prostrate Spurge (*Euphorbia supina*) use 150 pounds Quali-Pro Oxadiazon 2G (3.0 pounds of active ingredient) per acre.

CROP TOLERANCE:

Quali-Pro Oxadiazon 2G is recommended for use on a wide variety of container and field grown woody ornamental shrubs, vines, and trees. Quali-Pro Oxadiazon 2G may be used at the recommended rates on the following ornamental species:

CONTAINER GROWN	
ESTABLISHED	
Species	Varieties
Agapanthus	Africanus
Andromeda	Japanese
Arborvitae	American, Eastern, Oriental, Aureanana, Globe, Pyramidal, Woodward
Arctostaphylos	Fred Oelher
Azalea	Hino Red, Red, Boudoir, Snowball, Karen, Kaempp, Stuartonian, Mollishybird, Sweetheart Supreme, Salmon Beauty, Caroline Gable, Louise, Stewartstonian, Rosebud
Barberry	Redleaf, Marshall Red, Darwinii, Irwinii, Purpleleaf
Boxwood	Common, Japanese
Callistemon	Bottle-brush
Camellia	Glen #40, Sasanqua, Colonel Fiery
Ceanothus Cedar	Atlas, Portoford (Allumii)
Cotoneaster	Paineyi, Bearberry, Coral Beauty, Greyleaf, Cranberry, Silverleaf, Peking, Bright Bead, Dammeri Skolgholm, Microphylla, Lowfast, Glaucophylla.
Cypress	Monterey
Dogwood	Red osier
Douglas Fir	Pseudotsugamenziesii
Eucalyptus	Sideroxylon, Nicholli

Euonymus	Greenlane, Microphylla Variegata, Emeraldgaety, Dwarf, Gold spot, Golden Green, Silver King, Wintercreeper Vegetus, Ermerald Cushion, Spreading
Forsythia	Lynwood gold
Gardenia	Amy Yashioka, August Beauty, Mystery
Ginko	Biloba
Heath Heather	Mrs. Maxwell
Hedera	English Ivy
Honeysuckle	Purpleleaf, Purpurea
Holly	Dwarf Okinowan, Convexa, Rotundifolia, Hetzi, Helleri, Burfordi, English, Hellus, Dwarf Yaupon, Nellie R. Stevens, Boxleafod, Green Island, Japanese
Jasmine	Yellow Star, Star
Juniper	Barharbor, Wiltoni, Torulosa, Pfitzeriana Compacti, Pfitzeriana Aurea, Torulosa Variegata, Pfitzeriana Glauca, Gulf Stream, Tam, Armstrongii, Shore Andora, San Jose, Blue Haven, Stricta, Jade River, Broadmoor, Meyeri, Rocket, Plumosa, Royal Beauty, Praecox, Greek, Apiculata Blackburn, Plumosa Compacta, Procumbens, Blue Pfizer
Justica	Brandegeana
Leucothoe	Catesbaei, Axillaris
Myrtle	Compacta Noellii
Magnolia	Dr. Merrilli, Samuel Somers, Grandiflora
Mahonia	Oregon Grape-Holly
Osmanthus	Fortunes
Periwinkle	Vinca Minor
Photinia	Frazerii
Pine	Slash, Japanese Black, Montana, Scotch, Digger, Shore, Thumbergii
Pittosporum	Variegated, Japanese
Podocarpus	Macrophylla
Privet	Japanese, Waxleaf, Glossy, Vicary, Golden Vicary, Curly Leaf
Pyracantha	Victory, Scarlet Firethorn, Wyattii, Graberri
Quercus Raphiolepis	India Hawthorn
Rhododendron	Cunningham, Blue Diamond, Snow, Christmas, Chionoides, 'PJM', Dr. Rutgers, Coral Bells, President Clay
Rose	Summer Snow, Bigleaf Winter Creeper, Charlotte Armstrong
Star Jasmine	Yellow Star Jasmine
Syzygium	Paniculatum
Viburnum	Sandankwa, Laurestinus, David
Xylosma	Congestum

NEWLY TRANSPLANTED

Species	Varieties
Andromeda	Japanese
Arborvitae	Glove, Woodward
Azalea	Hindo Red, Louise, Rosebud, Stewartstonian
Barberry	Marshall Red, Darwinii, Irwinii, Purpleleaf
Cedar	Portoford (allumii)
Cotoneaster	Cranberry, Peking, Coral Beauty, Bright Bead, Painei, Dammeri Skolgholm, Microphylla, Glaucohylla
Cypress	Monterey
Dogwood	Red Osier
Euonymus	Greenlane, Emerald Cushion, Spreading
Forsythia	Lynwood Gold
Holly	Rotundifolia, Helleri, English, Japanese
Juniper	Plumosa, Plumosa Compacta, Old Gold, Pfitzeriana Aurea, Torulosa, Armstrongii, Tam, Wiltoni, Procumbens

Leucothoe	Axillaris
Magnolia	Dr. Merrill
Mahonia	Oregon Grape-Holly
Pine	Slash
Privet	Japanese, Waxleaf, Glossy
Pyracantha	Graberii
Rhododendron	Snow, Blue Diamond, Chionoides, 'PJM', Dr. Rutgers
Rose	Summer Snow, Bigleaf Winter Creeper, Charlotte Armstrong

FIELD GROWN	
ESTABLISHED	
Species	Varieties
Arborvitae	Pyramidal
Ash	Green
Azalea	Hinodi geri
Barberry	Red, Mentor
Birch	Clump, Paper
Camellia	Tiffany, Marie Brace, Elizabeth Dowd, Tom Knudsen, Crimson Robe, R.D. Wheeler
Cotoneaster	Lowfast, Skolgholm
Crabapple	Flowing, Torringo
Dogwood	Redstemmed, Red, Flowering
Euonymus	Burning Bush, Dwarf Burning Bush, Jewell
Forsythia	Showy Boarder, Lynngold, Lynwood Gold, Goldenbell
Holly	Boxleafed, Green Island, Japanese
Honeysuckle	Zabeli
Juniper	Plumosa, Mint Julip, Arcardia, Variegata; Plumosa Compacta, Andora, Wiltoni, Hetzi
Lilac	Persian, Dwarf Korean
Maple	Amur
Oak	Red
Perwinkle	Vinca Minor
Pine	Austrian, Scotch, Norway, Stone, Japanese, Black, Loblolly, Slash, Eastern White
Pittosporum	Undulatum
Privet	California, Japanese
Rhododendron	Rosebud
Rose	Texas Centennial, Sutters Gold, Show Girl
Russian Olive Spruce	Black Hills, Norway, Serbian
Taxus	Taunton Spreading Yew, Hicks
Tsuga	Canadian Hemlock
Viburnum	European Cranberry Bush
Xylosma	
Ground Covers	
Carpobrotus	Edule
Osteo	
Spermum Sedum	Album
Algerian Ivy	
Vinca Major	
English Ivy	
Pachysandra	Terminalis
Ajuga	Reptans
Euonymus	Fortunei Radicans

NEWLY TRANSPLANTED	
Species	Varieties
Camellia	Tiffany, Marie Bracey, Elizabeth Dowd, R.D. Wheeler, Tom Knudsen, Crimson Robe
Dogwood	Redstemmed, Red, Flowering
Forsythia	Showy Border, Lynwood Gold, Goldenbell
Hedera	English Ivy
Holly	Green Island, Japanese
Juniper	Plumosa, Andora, Wiltoni, Hetzi
Oak	Red
Periwinkle	Vinca Minor
Pine	White Austrian, Eastern White
Privet	California, Japanese
Rhododendron	Rosebud
Russian Olive Spruce	Black Hills, Norway, Serbian
Taxus	Hicks
Tsuga	Canadian Hemlock

ADDITIONAL APPLICATIONS

Do not apply more than 300 lbs. Quali-Pro Oxadiazon 2G per acre per year (equivalent to 6 lbs. a.i./A/year)

Do not apply more than 150 lbs. Quali-Pro Oxadiazon 2G per acre (equivalent to 3 lbs. a.i./A) in a single application.

For continued weed control, additional application of Quali-Pro Oxadiazon 2G may be made 60 to 120 days after the previous application in the following situations:

2 REPEAT TREATMENTS					
Container Grown Established		Container Grown Newly Transplanted		Field Grown Established and Newly Transplanted	
Species	Varieties	Species	Varieties	Species	Varieties
Abelia	Grandiflora	Common Boxwood		Holly	Nellie R. Stevens Dwarf Burford Burford
Common boxwood		English Ivy			
English Ivy		Juniper	Tam		
Holly	Helleri	Pine	Slash		
Juniper	Tam	Rhododendron	Obtusa, Kurume Formosa		
Pine	Slash				
Rhododendron	Obtusa Kurume Formosa Kurume Snow				
3 REPEAT TREATMENTS					
Container Grown Established					
Species			Varieties		
Abelia			Grandiflora		
Holly			Helleri, Rotundifolia		
Rhododendron			Kurume Formosa, Kurume Snow		
Greenhouse Grown Established					
Species			Varieties		
Rose			Sonia, Forever Yours, Coed, Golden Wave		
Forest Nurseries Seedlings- At least 4 weeks old					
Species			Varieties		
Pine			Loblolly, Slash, Eastern White		

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in dry area. Do not store near other pesticides, or seeds.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Nonrefillable Container (flexible-bag-all weights): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable Container (rigid-fifty lbs. or less): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Nonrefillable Container (rigid-greater than fifty lbs.): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Refillable Container: Refillable container. Refill this container with oxadiazon only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

LIMITATION OF WARRANTY AND LIABILITY

Read the entire directions for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following **CONDITIONS, DISCLAIMER OF WARRANTIES, and LIMITATIONS OF LIABILITY.**

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Makhteshim Agan of North America, Inc. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Makhteshim Agan of North America, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Makhteshim Agan of North America, Inc.'s election, the replacement of product.

Quali-Pro® is a registered trademark of Makhteshim Agan of North America, Inc.