



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

September 8, 2023

Haley Wade
Lead Regulatory Reviewer
Control Solutions, Inc.
5903 Genoa-Red Bluff
Pasadena, TX 77507-1041

Subject: Registration Review Label Mitigation for Novaluron, Updating of Annual Bluegrass Weevil Larvae Application Instructions & Addition of Optional Marketing Statements
Product Name: CSI 15-185T
EPA Registration Number: 53883-480
Decision Numbers: 581294, 588206 & 593291

Dear Haley Wade:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all of the information submitted with your application to support the Registration Review of the above referenced product in connection with the Novaluron Final and/or Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 § CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

A copy of your label stamped "Accepted" is enclosed. Products shipped after 12 months from the date of this amendment must bear the new revised label. Your release for shipment of the product bearing the amended label constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

Page 2 of 2
EPA Reg. No. 53883-480
Decision No. 581294, 588206 & 593291


If you have any questions, please contact Carmen Rodia via e-mail at Rodia.Carmen@epa.gov.

Sincerely,



Tamica L. Cain, Product Manager 10
Invertebrate-Vertebrate Branch 2
Registration Division (7505T)

Enclosure: Master Label Stamped "Notification", September 8, 2023



{Text in braces and italics are notes to the label editor.}

[Text in brackets alone are optional statements or alternative statements.]

NOVALURON	GROUP	15	INSECTICIDE
-----------	-------	-----------	-------------

CSI 15-185T

**[ABN: Quali-Pro Suprado SC; Quali-Pro Suprado; Suprado; Suprado SC;]
 [Optional generic descriptor(s): [Insecticide] [Larvicide] [Insect Growth Regulator]]**

[An insect growth regulator for control of listed immature pests on turfgrass]

OR

[For control of listed insect pests in or on golf courses; sports and recreational turf; commercial sod farms; and lawns on and around residential, commercial and institutional properties]

ACTIVE INGREDIENT:

*Novaluron 10.00%

OTHER INGREDIENTS: 90.00%

TOTAL: 100.00%

*Contains 0.89 lb. of novaluron per gallon of formulated product.



Control Solutions, Inc.
 5903 Genoa Red Bluff
 Pasadena, TX 77507

EPA Reg. No: 53883-480
 EPA Est. No: _____
 Net Contents: _____
{Lot number will be printed on container}

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
 (If you do not understand the label, find someone to explain it to you in detail.)

[[See inside label booklet for [additional] First Aid, Precautionary Statements, Directions for Use, and Storage and Disposal.] [See other panels for additional precautionary information.] [See below for additional Precautionary Statements]]

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information on this pesticide product (including health concerns or medical emergencies), you may also contact SafetyCall® International at 1-844-685-9173 , twenty-four (24) hours per day, seven (7) days per week.	

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants; and
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change clothing.

ENVIRONMENTAL HAZARDS

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water.

Surface Water Advisory: This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water.

This product is classified as having a medium potential for reaching both surface water and aquatic sediment via runoff for several weeks after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of novaluron from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

PHYSICAL AND CHEMICAL HAZARDS

DO NOT mix or allow coming in contact with Oxidizing agents. Hazardous Chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS:

- This product is for outdoor use only.
- **DO NOT** apply through any irrigation system.
- **DO NOT** apply this product in a way that will contact people or pets.
- **DO NOT** allow children or pets to enter treated areas until applications have been watered in and areas have dried.
- **DO NOT** spray the product into fish pools, ponds, streams or lakes.
- **DO NOT** apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.
- **DO NOT** apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and crevice treatment.

- **DO NOT** allow the product to enter any drain during or after application. *No permita la entrada a desagües internos o externos.*
- Follow proper disposal procedures on this label. *Siga las indicaciones del etiquetado para el desecho apropiado del producto.*



- For turfgrass applications, **DO NOT** apply more 4 quarts this product (0.89 lbs. of novaluron) per acre in a single application.
- **DO NOT** make applications at less than 30-day intervals.
- **DO NOT** exceed a total of 9.2 quarts this product (2.03 lbs. of novaluron) per acre per year.
- **DO NOT** make applications when turfgrass areas are waterlogged or the soil is saturated with water. Sufficient distribution of the active ingredient cannot be achieved under these conditions. The treated turf area must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile.
- To treat insects harbored in voids and cracks-and-crevices, applications must be made in such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants.
- **DO NOT** mow until after sufficient irrigation or rainfall has occurred so that uniformity of application will not be affected.
- **DO NOT** apply or irrigate after application to the point of runoff.
- **DO NOT** make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours).
- Rainfall within 24 hours after application may cause unintended runoff of pesticide application.
- Treat surfaces to ensure thorough coverage but avoid runoff.
- **DO NOT** apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur, unless for labeled sewer or drain or water treatment.
- **DO NOT** allow this product to enter any drain during or after application unless for labeled drain treatment.
- For any requirements specific to your State or tribe, consult the agency responsible for pesticide regulation.

READ ALL DIRECTIONS COMPLETELY BEFORE USE.

MANDATORY SPRAY DRIFT MANAGEMENT

Ground Boom Applications:

- **DO NOT** release spray at a height greater than 4 feet above the ground or crop canopy.
- Applicators must select nozzle and pressure that deliver medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- **DO NOT** apply when wind speeds exceed 15 mph at the application site.
- **DO NOT** apply during temperature inversions.

Boomless Ground Applications:

- Applicators must select nozzle and pressure that deliver medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- **DO NOT** apply when wind speeds exceed 15 mph at the application site.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

BOOM HEIGHT - Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

SPRAY DRIFT ADVISORIES

Boomless Ground Applications:

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications:

- Take precautions to minimize spray drift.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this

label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls over long-sleeved shirt and long pants;
- Chemical-resistant gloves made of any waterproof material; and
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. The area treated must be vacated by unprotected persons.

Do not enter or allow others to enter until sprays have dried.

Do not treat areas while unprotected humans or domestic animals are present in the treatment area. Because certain States may require more restrictive reentry intervals, consult your State Department of Agriculture for further information.

PRODUCT INFORMATION

NOVALURON, the active ingredient in this product, is an **insect growth regulator (IGR)** that controls immature stages of listed insect pests. The primary mode of action is to disrupt cuticle formation and deposition when immature insects change from one developmental stage to another, resulting in death at molting. Due to this mode of action, this product has no curative effect on adults that no longer molt. Novaluron has the potential to reduce the number of offspring produced by adults in some species. Consult the "Application Instructions" section of this label for timing instructions on specific pests.

This product should not be expected to provide a rapid, knock-down effect. IGRs such as this product do not kill immature susceptible insects until the next time they molt. Due to this, it may be necessary to wait several days or weeks to accurately evaluate this product's efficacy, depending on target species life cycle and growth stages present.

Water Protection Statements:

- **DO NOT** spray the product into fish pools, ponds, streams, or lakes unless for labeled water treatment.
- **DO NOT** apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur, unless for labeled sewer or drain or water treatment.
- **DO NOT** allow the product to enter any drain during or after application unless for labeled drain treatment.
- **DO NOT** apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment.
- **DO NOT** apply or irrigate to the point of runoff.

Rain Related Statements:

- **DO NOT** make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours).
- Rainfall within 24 hours after application may cause unintended runoff of pesticide application.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, this product contains a Group 15 insecticide. Any insect population may contain individuals naturally resistant to this product and other Group 15 insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of this product or other Group 15 insecticides within a growing season, or among growing seasons, with different Groups that control the same pests.
- Use tank mixtures with an insecticide from a different group that is equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for IRM.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The IRM benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an IRM benefit for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisor for any additional IRM and/or IPM recommendations for specific site and pest problems in your area.
- For further information or to report suspected contact your local retailer or your local university extension specialist.

TANK MIXING: This product may also be used in tank mixtures. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Test compatibility of the intended tank mixture before adding this product to the spray or tank mix. Add proportionate amounts of each ingredient to a pint or quart jar, cap, shake, and let set 15 minutes. Formation of precipitates that do not readily redisperse indicates an incompatible mixture that cannot be used. Add other products in the following order:

1. Water-soluble bags,
2. Wettable powders,
3. Dry flowables,
4. Liquid flowables,
5. Emulsifiable concentrates, and
6. Soluble materials such as fertilizers.

If tank mixing with other products, especially soluble fertilizers containing nitrogen or boron, allow products packed in water soluble bags to dissolve first. Read and follow label directions of all products used in the tank mix

before using. When tank mixing, always read all individual manufacturers' labels. In interpreting all labels for the tank mix, the most restrictive situations must apply. Tank mixtures are permitted only in those states where the tank mix partner is registered.

TURFGRASS APPLICATIONS

This product may be used to control listed soil- and thatch-inhabiting insect pests on turfgrass including residential lawns, business and office complexes, shopping complexes, multi-family residential complexes, golf courses, airports, cemeteries, parks, playgrounds, athletic fields and sod farms.

For optimum control of listed pests, applications should be timed as described for each species in the table below. Application timing can be based on historical monitoring of the site, previous records or experiences, current season adult trapping, soap flushes, or consult your State Extension Turf Specialist for specific information concerning timing in your region.

Apply this product in sufficient water (minimum 2 gallons per 1,000 ft²), depending on target pest, to provide adequate canopy and thatch penetration. Low volume applications (2 gallons per 1,000 ft²) must be watered in immediately with 0.25" – 0.5" of irrigation to move the active ingredient into the thatch and soil unless directed otherwise below for a specific pest. The use of accurately calibrated equipment normally used for the application of turfgrass insecticides is recommended. Use equipment which will produce a uniform, coarse droplet spray, using a low pressure setting to eliminate off target drift. Check calibration periodically to ensure that equipment is working properly.

TURF APPLICATION RATES				
Target Pests	Quarts/Acre	Fl. Oz./1,000 ft ²	lbs. A.I. per acre	Application Instructions
Annual bluegrass weevil larvae [<i>Listronotus maculicollis</i>]	3 – 4	2.2 - 3	0.67 – 0.89	<p>Following a 24-hour drying period, irrigate with a minimum of 0.1".</p> <p>Preventive Applications: To avoid turf damage, applications should be timed to adult peak migration period, early-instar stage (inside the stem), or at peak 3rd-instar emergence timing before visible damage typically develops.</p> <p>This product can also be used on overlapping growth stages as a curative approach in second-generation populations occurring later in summer months.</p> <p>Repeat if desired every 30 days to target larvae of second and third generations</p>
Billbug larvae Bluegrass billbug [, <i>Sphenophorus parvulus</i>]	3 – 4	2.2 - 3	0.67 – 0.89	Time application to target larvae when they leave the

Hunting billbug[, <i>Sphenophorus venatus vestitus</i>]				grass stems to feed on turf crowns and roots
Turf caterpillars Armyworms, Cutworms, Sod Webworms	1 – 2	0.7 – 1.5	0.22 – 0.45	If site data suggests a history of caterpillars, this product can be used as a preventative treatment. Repeat applications as necessary for subsequent generations.
White grubs Northern & Southern masked chafers[, <i>Cyclocephala borealis</i> , <i>C. immaculata</i> , and/or <i>C. lurida</i>]; Asiatic garden beetle[, <i>Maladera castanea</i>]; European chafer[, <i>Rhizotrogus majalis</i>]; Green June beetle[, <i>Cotinis nitida</i>]; May or June beetle[, <i>Phyllophaga spp.</i>]; Japanese beetle[, <i>Popillia japonica</i>]; Oriental beetle[, <i>Anomala orientalis</i>]	3 – 4	2.2 - 3	0.67 – 0.89	Preventive Applications: Make applications through peak egg hatch of target species. Use the upper end of the rate range if application is made 14 days or more prior to peak adult flight and/or egg laying. Post egg hatch application: After egg hatch of the target species has occurred or there is obvious turf damage from the current generation, use the upper end of the rate range.
Chinch bugs Southern chinch bug[, <i>Blissus insularis</i>]; Hairy chinch bug[, <i>Blissus leucopterus</i>]	3 - 4	2.2 - 3	0.67 – 0.89	This product controls immature stages of chinch bugs.
Black turfgrass ataeinus larvae [, <i>Ataeinus spretulus</i>]	3 - 4	2.2 - 3	0.67 – 0.89	Preventive Applications: Apply when early stages of first generation grubs are present. Higher rate may be required for late-stage grubs causing visible damage. Target grub stages of second generation in regions where they occur.

Turf Application Dilution Chart

	Turf Application Rates	Quarts of this product diluted to these volumes of finished spray
--	-------------------------------	--

Application Volume gal./1,000 ft ²	Fl. Oz./1,000 ft ²	Quarts/Acre	lbs. A.I. per acre	1 gallon	5 gallons	10 gallons	100 gallons
2	0.7	1	0.22	0.5	3	5	50
	1.5	2	0.45	1	5	10	100
	2.2	3	0.67	1.5	8	15	150
	3.0	4	0.89	2	10	20	200
3	0.7	1	0.22	0.3	1.7	3.3	33.3
	1.5	2	0.45	0.7	3.3	6.7	66.7
	2.2	3	0.67	1	5	10	100
	3.0	4	0.89	1.3	6.7	13	133
4	0.7	1	0.22	0.25	1.25	2.5	25
	1.5	2	0.45	0.5	2.5	5	50
	2.2	3	0.67	0.75	3.75	7.5	75
	3.0	4	0.89	1	5.0	10	100
5	0.7	1	0.22	0.2	1	2	20
	1.5	2	0.45	0.4	2	4	40
	2.2	3	0.67	0.6	3	6	60
	3.0	4	0.89	0.8	4	8	80

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Keep this product in its tightly closed original container. Store in a cool, dry (preferably locked) area that is inaccessible to children and animals, in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. **DO NOT** store this product above 100°F for extended periods of time. Storage below 32°F may cause freezing. If container is damaged or if a spill has occurred, use product immediately, or dispose of product as indicated below.

PESTICIDE DISPOSAL: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility. **DO NOT** pour or dispose of down the drain or sewer. Call your local solid waste agency for local disposal options.

CONTAINER HANDLING: Nonrefillable container. DO NOT reuse or refill this container! Empty container by using the product according to the label directions. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows:

[(<5 gal)] Empty the remaining contents into application equipment or a mix tank and drain for 10 sec after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 sec. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure 2 more times.]

[(>5 gal)] Empty remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 sec. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure 2 more times.]

Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by other means allowed by Federal, state or local authorities.

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

WARRANTY STATEMENT

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Control Solutions, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Control Solutions, Inc. and Seller harmless for any claims relating to such factors.

To the extent consistent with applicable law, Control Solutions, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Control Solutions, Inc., and Buyer and User assume the risk of any such use. CONTROL SOLUTIONS, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, IT IS **CONTROL SOLUTIONS INC., AND SELLERS INTENTION THAT** in no event shall Control Solutions or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF CONTROL SOLUTIONS, INC., AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF CONTROL SOLUTIONS, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

To the extent consistent with applicable law, Control Solutions, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Control Solutions, Inc.

Optional Marketing Statements:

- Disrupts Insect Development: as a third-generation chitin inhibitor, this product interferes with insect development via transovarian disruption, resulting in reduced egg lay and hatch.
- IPM Friendly: unique mode of action reduces the potential for environmental exposure when compared to many current standard materials.
- Resistance Management: alternative mode of action provides an excellent resistance management tool for long lasting protection.
- Insect Growth Regulator (IGR)
- Controls [1st and 2nd] [early] [late] instar larva[e]
- Breaks the life-cycle of listed insect pests by reducing egg viability.
- Stops damage before it starts!
- Early application for immediate protection, followed by additional sprays to keep insects in check.
- The only IGR for [grub] [caterpillar] [Annual bluegrass weevil] control.
- Preventative Grub Control
- Preventative Control of Grubs
- Controls Grubs
- Residual Grub Control

- Preventative Control of Listed Insect Pests
- Reduces [egg lay and hatch] [egg viability] of annual bluegrass weevil populations that can cause turf loss.
- Effective against pyrethroid-resistant annual bluegrass weevil populations by reducing [egg lay and hatch] [egg viability]
- Effective against pyrethroid-resistant annual bluegrass weevil populations by reducing fecundity
- Reduces fecundity of pyrethroid-resistant annual bluegrass weevil populations
- A new turfgrass mode of action for improved resistance management
- Patent Pending
- Shake well

Optional Graphic(s):

