



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Registration Division (7505P)
 1200 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

EPA Reg. Number:

53883-489

Date of Issuance:

9/3/21

NOTICE OF PESTICIDE:

Registration
 Reregistration
 (under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

CSI 19-313H Glufosinate +
 Imazapyr

Name and Address of Registrant (include ZIP Code):

Control Solutions, Inc.
 5903 Genoa-Red Bluff
 Pasadena, TX 77507-1041

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. The data requirements for storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) are not satisfied. A one year study is required to satisfy these data requirements. You have 18 months from the date of registration to provide these data.

Signature of Approving Official:

Heather E McFarley

Heather McFarley, Acting Product Manager 24
 Fungicide Herbicide Branch, Registration Division (7505P)

Date:

9/3/21

3. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 53883-489.”
4. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 07/01/2020
- Alternate CSF 1 dated 07/01/2020
- Alternate CSF 2 dated 07/01/2020
- Alternate CSF 3 dated 07/01/2020
- Alternate CSF 4 dated 07/01/2020
- Alternate CSF 5 dated 07/01/2020
- Alternate CSF 6 dated 07/01/2020
- Alternate CSF 7 dated 07/01/2020
- Alternate CSF 8 dated 07/01/2020
- Alternate CSF 9 dated 07/01/2020
- Alternate CSF 10 dated 07/01/2020
- Alternate CSF 11 dated 07/01/2020

If you have any questions, please contact Manjula Unnikrishnan by phone at 703-347-8520, or via email at unnikrishnan.manjula@epa.gov .

Enclosure:

- Stamped label

[Text in brackets and italics are notes to the label editor.]

[Text in brackets alone are optional statements or alternative statements.]

Glufosinate-ammonium	Group	10	Herbicide
Imazapyr	Group	2	Herbicide

CSI 19-313H Glufosinate + Imazapyr

Alternate brand name(s): [Eraser Max GFSN], [Eraser Pro GFSN], [Eraser EXtra], [Eraser Xtra], [Eraser Max GFSN 5.78], [Eraser Pro GFSN 5.78], [Eraser Extra 5.78], [Eraser Xtra 5.78] [Eraser Extra 5.78 GFSN]

Optional generic descriptor(s): [Farm and Ranch][Farm & Ranch]

ACTIVE INGREDIENT:

Glufosinate-ammonium 5.78%

Imazapyr..... 0.39%

OTHER INGREDIENTS: 93.83%

TOTAL: 100.00%

Contains 0.47 lbs. of glufosinate-ammonium per gallon.

Contains 0.032 lbs. of imazapyr per gallon.



ADAMA

Consumer & Professional Solutions

Manufactured for:

Control Solutions, Inc.
5903 Genoa Red Bluff
Pasadena, TX 77507

EPA Reg. No: 53883-XXX

EPA Est. No: _____

Net Contents: _____


[Store and transport in an upright position.]

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you **DO NOT** understand the label, find someone to explain it to you in detail.)

[Alternate text – all or in part: See side panel(s) for Precautionary Statements, Directions for Use and Storage and Disposal.]

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency medical treatment information call: SafetyCall® (866) 897-8050.	



PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Wear protective eyewear. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to plants. Drift and run-off may be hazardous to plants in water adjacent to treated areas. To protect the environment, **DO NOT** allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow pesticide off the treatment area. Rinsing application equipment over the treated area will help to avoid run off to water bodies or drainage systems. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate

What does the product do?

[Gluf 5.78 – Imaz 0.39] [This product] kills and prevents regrowth of unwanted weeds, grasses and plants for up to one year. Use this product to kill unwanted vegetation on driveways, walkways, parking areas, sidewalks, patios, gravel areas, rock landscapes, under fences and around curbs. Use in areas where no vegetation growth is desired for up to [1] [one] year. [Product name] [This product] starts working as soon as it contacts the plant and weeds begin to turn yellow and wilt within 3-5 days. Complete kill occurs in 1 to 2 weeks.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All handlers must :

- Wear long-sleeved shirt, long pants, shoes and socks
- Wear protective eyewear

USER SAFETY REQUIREMENTS

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

PRODUCT INFORMATION

This product is a nonselective water-soluble herbicide for application as a foliar spray for the control of a broad spectrum of emerged annual and perennial grass and broadleaf weeds. This product will also control certain woody species. Plants that have not yet emerged at the time of application will not be controlled. THOROUGH SPRAY COVERAGE IS IMPORTANT. Visual effects and control from application of This product occur within 2 to 4 days after application under good growing conditions.

This product is nonselective and toxic to plants. Avoid all contact with foliage or green tissue of desirable vegetation. Avoid direct spray contact with green, thin, or uncalloused bark of desirable vegetation or plant injury may result. Avoid spraying in areas where desirable plant roots may be present. **DO NOT** apply closer than twice the distance from the trunk to the drip line of desirable trees and shrubs. Apply to locations only where no vegetation growth is desired for up to one year.

This product works best when weeds are actively growing. Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures. Weeds under stress or in dense populations will require application at the highest rate specified. Refer to the How to Apply section of this label.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

READ ALL DIRECTIONS COMPLETELY BEFORE USE.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

DO NOT use this product until you have read the entire label. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

NON AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. The application for trimming and edging, industrial, recreational and public areas, and farmsteads are not within the scope of the WPS.

WEED RESISTANCE MANAGEMENT

For resistance management, This product contains Group 10 and Group 2 herbicides. Any weed population may contain or develop plants naturally resistant to This product and other Group 10 and 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of This product or other Group 10 and Group 2 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide used and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g. higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout before and after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method including hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.

- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance management and/or integrated weed-management directions for specific crops and weed biotypes.
- Report any incidence of non-performance of this product against a particular weed species to Control Solutions, Inc., your local retailer or your local extension specialist. If resistance is suspected, treat weed escapes with a herbicide having a different mechanism of action and/or use non-chemical means to remove escapes, as practical, with the goal of preventing further seed production.

BEST MANAGEMENT PRACTICES

Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is advised. A diversified weed management program may include the use of multiple herbicides with different modes of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices. Research has demonstrated that using the labeled rate and directions for use is important to delay the selection for resistant weeds. Scouting after herbicide application is important because it can facilitate the early identification of weed shifts and/or weed resistance and thus provide direction on future weed management practices. One of the best ways to contain resistant weed populations is to implement measures to avoid allowing weeds to reproduce by seed or to proliferate vegetatively. Cleaning equipment between sites and avoiding movement of plant material between sites will greatly aid in reducing the spread of resistant weed seed.

MANDATORY SPRAY DRIFT

- **DO NOT** apply where wind speeds 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.
- For ground applications, applicators must select nozzle and pressure that deliver medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- Spray at the appropriate boom height based on nozzle selection and nozzle spacing, but **DO NOT** exceed a boom height of 24 inches above target pest or crop canopy. Set boom to lowest effective height over the target pest or crop canopy based on equipment manufacturer's directions. Automated boom height controllers are directed with large booms to better maintain optimum nozzle to canopy height. Excessive boom height will increase the potential for spray drift.
- For non-crop vegetation management ground applications, apply with the nozzle height no more than 4 feet above the ground or target vegetation, unless necessitated by the application equipment. Examples would include roadside, railroad, utility rights of way, forestry and other industrial vegetation management applications where safety or natural barriers obstruct application.

SPRAY DRIFT ADVISORIES

POLLINATOR ADVISORY: This product contains a herbicide. Follow all label directions and precautions to minimize potential off-target exposure in order to prevent effects to non-target plants adjacent to the treated site which may serve as habitat or forage for pollinators.

Spray Drift Management: The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors where making application decisions.

- **Importance of Droplet Size:** The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!** See Wind, Temperature and Humidity, and Temperature Inversions sections of this label.

- Techniques for Controlling Droplet Size:
 - Volume – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rates flow produce larger droplets.
 - Pressure – Use the lower spray pressures specified for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.
 - Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.
- Wind: Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. AVOID APPLICATIONS DURING GUSTY OR WINDLESS CONDITIONS. Note: Local terrain can influence wind patterns. Every applicator needs to be familiar with local wind patterns and how they affect spray drift.
- Temperature and Humidity: When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.
- Temperature Inversions: Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.
- Shield Sprayers: Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

USE RESTRICTIONS:

- **DO NOT** apply this product by aerial application.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply directly to or allow drift to contact desirable green tissue or green, thin or uncalloused bark of desirable vegetation.
- **DO NOT** apply this product to residential areas as other than a spot or directed spray.
- **DO NOT** apply near bodies of water including, fish pools, ponds, lakes or streams.
- **DO NOT** apply over the roots systems of desirable plants. For trees and shrubs **DO NOT** apply closer than twice the distance from the trunk to the drip line as roots may be within this area.
- **DO NOT** apply to slopes in landscape as movement on soil surfaces may damage desirable plants down slope.
- **DO NOT** apply next to a fence if desirable vegetation is growing on the other side, or if future planting is intended.
- **DO NOT** use before planting lawns, fruits, vegetables, flowers, or other plants because this product remains active in soil for up to one year. In soils with heavy clay, weed control may exceed one year.
- **DO NOT** use on desirable lawns.
- **DO NOT** apply to powdery, dry or light sandy soils when the soil may be blown by wind. If necessary soil surface must be settled by rain or irrigation.
- **DO NOT** apply more than 9.37 fl. oz. of product (0.34 lbs. glufosinate-ammonium, 0.002 lbs. imazapyr) per 1000 sq. ft. in any single application.
- **DO NOT** apply more than 9.37 fl. oz. of product (0.34 lbs. glufosinate-ammonium, 0.002 lbs. imazapyr) per 1,000 sq. ft. per year.
- **DO NOT** apply more than 408.5 fl. oz. (12.77 qts.) of product (1.5 lbs. glufosinate-ammonium, 0.10 lbs. imazapyr) per acre in any single application.

- **DO NOT** apply more than 408.5 fl. oz. (12.77 qts.) of product (1.5 lbs. glufosinate-ammonium, 0.10 lbs. imazapyr) per acre per year.
- **DO NOT** apply more than 1 applications per year when using the maximum single application rate (excluding spot treatments).
- **In the State of New York only:** Not for use in Nassau and Suffolk Counties

WHERE TO USE: Use on non-crops areas: Industrial sites, petroleum tank farms, lumberyards, manufacturing sites, storage areas, airports, roadsides, curbs, sidewalks, driveways, parking areas, brick walks, gravel paths, patios, vacant lots, paved areas (including crack and crevice), warehouse areas, urban areas, recreational areas, tennis courts, fencerows, along fences of paddocks, corrals, feedlots, and livestock holding areas, hedgerows, dry ditchbanks, around domestic dwellings, around farm buildings and barns, shelter beds, farm yards, farm premises, outdoor buildings, structures and shelters including beef barns, hog barns, dairy barns, stables, dog kennels and poultry houses, and for establishment and maintenance of wildlife openings.

WHEN TO APPLY: This product has foliar and root activity. Apply to bare ground or when weeds are present. Best results are obtained when weeds are actively growing. If weeds are taller than six inches, mow or cut down weeds and remove debris before treatment, and allow for 2 weeks of new growth. If woody vines or brush is mowed or cut, wait until full leaf formation before treatment. Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures. Weeds under stress or in dense populations will require application of the highest rate. Refer to the How to Apply section of this label.

HOW TO MIX: This product must be mixed with water to make a finished spray solution as follows:

1. Fill the spray tank with the required amount of water.
2. Add the proper amount of this product, then mix thoroughly.

HOW TO APPLY

Equipment: Use hand operated spray equipment including backpack sprayers, pump-up sprayers or sprinkling cans, or ATV/Tractor mounted sprayers.

Spot Application Use Rates: This product may be used as a spot or directed spray application using 4 to 9.37 fl. oz. (0.015 to 0.034 lbs. glufosinate-ammonium, 0.0010 to 0.0023 lbs. imazapyr) per gallon of water. Spray undesirable vegetation foliage on a spray-to-wet basis. Avoid product runoff to nearby desirable plants. Ensure uniform and complete coverage. Use a coarse spray. **DO NOT** spray during windy conditions. Backpack, pump-up, and hydraulic sprayers may be used. Thoroughly clean the sprayer following use.

Broadcast or Boom Applications: Apply 4 to 12.77 qt. per acre (0.47 to 1.5 lbs. glufosinate-ammonium, 0.032 to 0.10 lbs imazapyr) depending upon the weed and stage of growth. **DO NOT** spray during windy conditions. Backpack, pump-up, and hydraulic sprayers may be used. Thoroughly clean the sprayer following use.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Store upright at room temperature. Avoid exposure to extreme temperatures. Keep this product in its tightly closed original container. Store in a cool, dry (preferably locked) area that is inaccessible to children and animals. In case of spillage or leakages, soak up with an absorbent material including sand, sawdust, earth, Fuller's earth, etc. Dispose of with chemical waste.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spilled bait, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance.

CONTAINER HANDLING: If empty - Nonrefillable container. **DO NOT** reuse or refill this container. Offer empty container for recycling if available or puncture and dispose of container in a sanitary landfill,

or by other procedures approved by State and local authorities. Clean container promptly after emptying. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

If partly filled: Call your local solid waste agency for disposal instructions. Unless otherwise instructed, place in trash. Never pour unused product down the drain or on the ground.

[Commercial Language]

For Bulk containers: (Refillable Container) Refill this container with pesticides only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the re-filler. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or re-circulate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

WARRANTY STATEMENT

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Control Solutions, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Control Solutions, Inc. and Seller harmless for any claims relating to such factors.

To the extent consistent with applicable law, Control Solutions, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Control Solutions, Inc., and Buyer and User assume the risk of any such use. CONTROL SOLUTIONS, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, IT IS **CONTROL SOLUTIONS INC., AND SELLERS INTENTION THAT** in no event shall Control Solutions or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF CONTROL SOLUTIONS, INC., AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF CONTROL SOLUTIONS, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

To the extent consistent with applicable law, Control Solutions, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Control Solutions, Inc.

[In lieu of warranty statement, the following text may be used:]

Notice – To the extent consistent with applicable law, Buyer assumes all responsibility for safety and use not in accordance with directions.

Optional Marketing Statements:

- Make only what you need
- Mix into tank sprayer
- Multipurpose grass and broadleaf weed control
- Next day [this weekend] results: Begins killing on contact, visible results in 24 hours.
- No odor
- No root. No weed. No problem
- Fast acting formula
- Non-Selective: Kills weeds & grasses
- Once it enters the plant, it won't wash off
- One application kills over 200 types of weeds and grasses, roots and all (maximum effectiveness)
- Proven (trusted) performance
- Provides maximum control: Kills weeds to the root so they don't come back
- Quick-action formula lets you see results fast
- RainFast in hours - enters the plant on contact
- Same effective results even if it rains within hours
- Two [2] applications control (kills) most weeds when used as broadcast or boom
- Three [3] applications control (kills) most weeds when used as spot-sprays
- Spray weeds until thoroughly wet
- Starts to work immediately
- Starts working on contact
- Starts working in minutes
- [Use] For Brush Control
- [Use] For Weed Control: Use along fences, paths, patios, sidewalks and driveways, buildings, and in brick and gravel walkways.
- [Use] For Spot Spraying: Use in and around fences, driveways and walkways
- [Use] For Stumps
- Use as a spot treatment along sidewalks, fences, driveways, and patios
- Use along fences; in cracks of walks, drives and patios
- Visible effects are gradual wilting and yellowing advancing to complete browning and root destruction.
- [Provides] Visible [see] results in [hours] [1 day]
- Visible Results in 24 Hours
- See results in 24 hours
- Water based formula
- Weed Eliminator
- Weed Killer
- Kills to the root!
- Kills to the root so weeds won't grow back
- Kills weeds in around fences, driveways, walkways, in and around patios, paths, sidewalk cracks
- See results in 1-2 days
- Contains See-Spray [lets you see where you spray]
- Kills tough weeds, roots and all
- [This product] will kill almost all unwanted vegetation, including annual and perennial weeds, grasses, and other unwanted plants
- [2 1/2 gallons] makes up to [106 1/2] [80] [52.5] [40] gallons. *{Note to reviewer: Mixture volume on label to vary based on equivalent use rate for different packaging sizes}*
- [2 1/2 gallons] covers up to [106,500 sq. ft.] [80,000 sq. ft] [52,500 sq. ft] [40,000 sq. ft.] [16,000

sq. ft] *{Note to reviewer: Area of coverage on label to vary based on equivalent use rate for different packaging sizes}*

- Kills the Root
- Kills weeds, grasses and brush
- Use around sidewalks and driveways.
- [One gallon] makes up to [42 1/2] [32] [21] [16] gallons. *{Note to reviewer: Mixture volume on label to vary based on equivalent use rate for different packaging sizes}*
- [One gallon] covers up to [42,500 sq. ft.] [32,000 sq. ft] [21,000 sq. ft.] [16,000 sq. ft] *{Note to reviewer: Area of coverage on label to vary based on equivalent use rate for different packaging sizes}*
- [1/2 gallon] makes up to [21 1/4] [16] [10.5] [8] gallons. *{Note to reviewer: Mixture volume on label to vary based on equivalent use rate for different packaging sizes}*
- [1/2 gallon] covers up to [21,250 sq. ft.] [16,000 sq. ft] [10,500 sq. ft.] [8,000 sq. ft] *{Note to reviewer: Area of coverage on label to vary based on equivalent use rate for different packaging sizes}*
- [One quart] makes up to [10 1/2] [10] [5] [4] gallons. *{Note to reviewer: Mixture volume on label to vary based on equivalent use rate for different packaging sizes}*
- [One quart] covers up to [10,500 sq. ft.] [10,000 sq. ft] [5,000 sq. ft.] [4,000 sq. ft] *{Note to reviewer: Area of coverage on label to vary based on equivalent use rate for different packaging sizes}*
- [One pint] makes up to [5] [3] [2] gallons. *{Note to reviewer: Mixture volume on label to vary based on equivalent use rate for different packaging sizes}*
- [One pint] covers up to [5000 sq. ft.] [3,000 sq. ft] [2,000 sq. ft.] *{Note to reviewer: Area of coverage on label to vary based on equivalent use rate for different packaging sizes}*
- Makes up to [42 1/2] [42] [10 1/2] [10] [5] gallons

The following claims used for six (6) months from the first date of shipment to a customer in accordance with Federal and State Registrations.

- New Formula
- New