



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

April 24, 2025

Shannon Cooley
Regulatory Specialist
Control Solutions, Inc.
5903 Genoa-Red Bluff
Pasadena, TX 77507-1041

Subject: Notification per PRN 98-10 – Adding logo/graphics to the label.
Product Name: CSI 19-304 Fungicide
EPA Registration Number: 53883-502
Application Date: 05/10/2024
Case Number: 611865

Dear Ms. Cooley:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped “NOTIFICATION” and placed in our records.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you have any questions, please contact Jamey Shuler by phone at (202) 566-2898, or via email at Shuler.Jamey@epa.gov.

Sincerely,

Kable Bo Davis

Kable Bo Davis, Senior Advisor
Office of Pesticide Programs
Registration Division; Immediate Office

[Text in brackets and italics are notes to the label editor.]

[Text in brackets alone are optional statements or alternative statements.]

[Text separated by "/" indicates an either/or choice.]

NOTIFICATION

53883-502

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

04/24/2025

Fluazinam	GROUP	29	FUNGICIDE
Tebuconazole	GROUP	3	FUNGICIDE
Azoxystrobin	GROUP	11	FUNGICIDE

CSI 19-304 Fungicide

[Fungicide] [Flowable Fungicide] [3-Way Fungicide]

[Broad Spectrum fungicide for control of turf diseases in golf course turf.]

[ABNs: Quali-Pro® Elluvia, Elluvia, Quali-Pro® Kanati, Kanati, Quali-Pro® Synch, Synch, Quali-Pro® Amira, Amira]

A combination of contact and systemic fungicides for broad spectrum control of turfgrass diseases on golf courses only.

ACTIVE INGREDIENTS

Fluazinam: [3-chloro-N-[3-chloro-2,6-dinitro-4trifluoromethyl)-phenyl]- 5-trifluoromethyl-2- pyridinamine] 19.5%

Tebuconazole: [alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl) -1H-1,2,4-triazole-1-ethanol] 6.8%

Azoxystrobin: [methyl (E)-2-[2-[6-(2-cyanophenoxy)-pyrimidin-4-yloxy] phenyl]-3-methoxyacrylate] 6.2%

OTHER INGREDIENTS 67.5%

TOTAL.....100.00%

Contains 1.88 lbs. Fluazinam, 0.65 lbs. Tebuconazole, and 0.59 lbs. Azoxystrobin per gallon.



ADAMA

Consumer &
Professional
Solutions

Manufactured for:
Control Solutions, Inc.
5903 Genoa Red Bluff
Pasadena, TX 77507

EPA Reg. No: 53883-502

EPA Est. No: _____

Net Contents: _____

KEEP OUT OF REACH OF CHILDREN

CAUTION

[Alternate text – all or in part: See [side] [back] panel(s) for First Aid, Precautionary Statements, Directions for Use and Storage and Disposal.]

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• DO NOT induce vomiting unless told to do so by the poison control center or doctor.• DO NOT give anything by mouth to an unconscious person.
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER <p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact SafetyCall® International for emergency medical treatment at (866) 897-8050 or the Poison Control Hotline at 1(800)222-1222</p> <p>For general information on this product, contact the National Pesticides Information Center (NPIC) at 1-800-858-7378, Monday through Friday, 8 AM to 12 PM PST, or at http://npic.orst.edu.</p>	

[Note to label editor: Table format will be used if space allows for the first aid section; otherwise, paragraph format will be used.]

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Causes moderate eye irritation. Wear appropriate protective eyewear. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants coverall/double layer
- Chemical resistance gloves
- Shoes plus socks
- Protective eyewear

User Safety Requirements

Follow manufacturer’s instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product’s concentrate. Do not reuse.

Engineering Controls

When handlers use closed systems, or enclosed cabs, in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for “applicators and other handlers” and have such PPE immediately available for use in an emergency, including a spill or equipment break down. **DO NOT** allow contact between contaminated sprayer parts and unprotected skin. Ensure sprayer is washed down daily.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS:

This product is toxic to mammals, fish and aquatic invertebrates. **DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high-water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate. **DO NOT** apply when weather conditions favor drift from treated areas. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas.

GROUND WATER ADVISORY

Azoxystrobin can be persistent for several months or longer. Azoxystrobin has degradation products which have properties similar to chemicals which are known to leach through soil to ground water under certain conditions as a result of agricultural use. Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this fungicide product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

SURFACE WATER ADVISORY

Leave an uncultivated boundary within 25 feet of permanent water bodies (lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, and estuaries) to allow growth of a vegetative filter strip. This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Failure to follow the use directions and precautions on this label may result in turf injury, poor disease control, and/or illegal residues. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Turf grasses on golf courses are not within the scope of the Worker Protection Standard. **DO NOT enter treated areas until sprays have dried.**

PRODUCT INFORMATION

This product is a broad-spectrum fungicide for use to control diseases of turf grasses on golf courses only.

This product may be applied with all types of spray equipment used for ground applications on golf courses. Apply this product in enough water to obtain adequate uniform coverage of the turf (minimum spray volume of 87 gallons per acre or minimum of 2 gallons per 1000 ft²).

PRECAUTIONARY STATEMENTS

DO NOT apply consecutive applications during or just after dormancy break.

DO NOT apply to bermudagrass greens when temperatures will exceed 85°F.

DO NOT make consecutive applications of this product on greens.

USE RESTRICTIONS

- For use on golf course turf only.
- Not for use in California.
- Not for use in New York State.
- **DO NOT** spray this product where spray drift may reach apple trees.
- **DO NOT** use spray equipment which has been previously used to apply this product to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.
- **DO NOT** apply this product to home lawns (turf) or any other residential turf.
- **DO NOT** use on home lawns and other residential turf sites including apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e., elementary, middle and high school), campgrounds, churches, and theme parks.
- **DO NOT** apply more than 2 sequential applications for Pythium spp. Control.
- For all other diseases when Pythium spp. Is not present, do not apply more than 3 sequential applications.
- Not for use on turf being grown for sale or commercial use as sod.
- **DO NOT** use treated grass or turf clippings for animal feed.
- **DO NOT** apply this product by aerial application.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply this product when the wind direction is toward aquatic areas to minimize drift to these areas.
- **DO NOT** exceed 572 fl. oz. of this product (8.4 lbs. fluazinam, 2.9 lbs. tebuconazole, 2.6 lbs. azoxystrobin) per acre per year.
- Maximum single application rate is 1.16 fl. oz. (0.017 lbs. fluazinam, 0.0059 lbs. tebuconazole, 0.0053 lbs. azoxystrobin)/1000 sq. ft.
- **DO NOT** apply more than six (6) applications per year
- **DO NOT** apply at less than 14-day intervals.
- Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather-related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions. Where states have more stringent regulations than this label, they must be observed. **DO NOT** allow spray mixture to stand overnight or for prolonged periods in the spray tank.
- **DO NOT** apply on top of snow.
- **DO NOT** apply this product within 25 feet of permanent water bodies (lakes, reservoirs, rivers, permanent streams, marshes or natural ponds and estuaries)

INTEGRATED PEST MANAGEMENT

This product is an excellent disease control agent when used according to label directions for control of a broad spectrum of listed plant diseases. This product is for use in programs which are compatible with the principles of Integrated Pest Management (IPM), which may include the use of disease resistant crop varieties, cultural practices, biological control agents, pest scouting and disease forecasting systems aimed at preventing economic pest damage and reducing unnecessary applications of pesticides. Practices known to reduce disease development should be followed. Consult your state cooperative extension service or local agricultural authorities for additional IPM strategies established in your area. This product may be used in State Agricultural Extension advisory (disease forecasting) programs that specify application timing based on environmental factors which favor disease development.

FUNGICIDE RESISTANCE MANAGEMENT

For resistance management, note that this product contains fungicides from FRAC Group 3 (tebuconazole), Group 11 (azoxystrobin), and Group 29 (fluazinam). Any fungal population may contain individuals naturally resistant to this product and other Group 3, Group 11, or Group 29 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies must be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 3, Group 11, or Group 29 fungicides within a growing season sequence with different groups including one from the dicarboximide (Group 2) or benzimidazole class (Group 1) that control the same pathogens. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, use of historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, and disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact your local retailer or your university extension specialist.

MANDATORY SPRAY DRIFT

Ground Applications

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy.
- For all applications, applicators are required to use a medium or coarser spray droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

- THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
- BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.
- IMPORTANCE OF DROPLET SIZE
An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.
Controlling Droplet Size - Ground Boom
 - Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
 - Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
 - Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.
- BOOM HEIGHT - Ground Boom
Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.
- SHIELDED SPRAYERS
Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.
- TEMPERATURE AND HUMIDITY
When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.
- TEMPERATURE INVERSIONS
Drift potential is high during a temperature inversion. Temperature inversions are characterized

by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

- WIND Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

- Boom-less Ground Applications:
Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.
- Handheld Technology Applications:
Take precautions to minimize spray drift.

PLANT SENSITIVITY

NOTE TO USER: Although this product has been evaluated on several turf species with no indication of phytotoxicity, neither the manufacturer nor seller has determined whether or not THIS PRODUCT can be used safely on all turf species. The professional user need to determine if this product can be used safely prior to commercial use by testing on the type of turf to be treated at specified rates for phytotoxicity.

TANK MIXING

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. This product is physically compatible with many products specified for control of diseases, liquid fertilizers, growth regulators, insecticides and biological control products. If tank mixes are desired, read and follow all directions, precautions, and limitations on labeling of all products used. This product must not be mixed with any product which prohibits such mixing. It is the applicator's responsibility to ensure that the companion product is EPA approved for use on the intended site. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are registered.

This product is compatible with the products mentioned above, provided sufficient free water is available for dispersion of all the tank mix products. However, the physical compatibility of this product with tank mix partners needs to be evaluated before use. A jar test should be conducted with the intended tank-mix products prior to preparation of large volumes. Use the following procedure:

Jar Compatibility Test:

1. Pour the specified proportions of the products into a suitable container of water.
2. Mix thoroughly.

3. Allow to stand for 5 minutes. If the combination remains mixed or can be re-mixed readily, it is considered physically compatible. Any physical incompatibility in the jar test indicates that this product must not be used in the tank mix. Consult compatibility charts or your local or state agricultural authorities for additional compatibility information.

Mixing Order:

- Materials should be thoroughly dispersed and mixed before adding additional tank mix components.
- Maintain constant agitation during mixing and application.
- Fill tank with water to half of the desired final spray volume.
- Begin agitation prior to adding tank mix partners in the following order:
 1. Products packaged in water-soluble bags
 2. Water-dispersible products (dry flowables, wettable powders, suspension concentrates)
 3. Liquid flowables
 4. Water-soluble products
 5. Emulsifiable concentrates
- Maintain agitation and add remaining water to bring to final volume.

TURFGRASS DISEASE CONTROL

This product can be used for golf course applications on warm season and cool season grasses or their mixtures. This product is not phytotoxic when used in accordance to the label.

Note: Bermudagrass can be sensitive to this product under certain conditions.

This product can be used for the prevention and control of the diseases listed below. For optimum disease control, apply this product preventatively when conditions are favorable for disease development. Apply uniformly with properly calibrated low volume, handheld, backpack, mechanical or motorized ground broadcast spray equipment.

Apply in sufficient water (1 to 4 gallons of water per 1000 ft²) to provide thorough coverage of targeted turf plant parts. Higher spray volumes are more typically needed on higher cuts of turf. For foliar diseases, allow foliage to dry thoroughly before irrigation.

[Text in brackets and italics are notes to the label editor.]
[Text in brackets alone are optional statements or alternative statements.]
[Text separated by “/” indicates an either/or choice.]

USE RATE		
Disease	Rate of this product fl. oz. / 1000 ft ²	Notes
Anthracnose <i>Colletotrichum cereale</i>	1.16 (0.017 lbs. Fluazinam, 0.0059 lbs. Tebuconazole, 0.0053 lbs. Azoxystrobin) (21 day interval)	Apply preventatively when conditions favor disease development. Allow spray to dry on leaves without watering in.
Brown Patch, Large Patch <i>Rhizoctonia solani</i>	1.16 (0.017 lbs. Fluazinam, 0.0059 lbs. Tebuconazole, 0.0053 lbs. Azoxystrobin) (21 day interval)	Apply preventatively when conditions favor disease development.
Dollar Spot , <i>Clarireedia</i> spp. (formerly <i>Sclerotinia</i> <i>homeocarpa</i>)	1.16 (0.017 lbs. Fluazinam, 0.0059 lbs. Tebuconazole, 0.0053 lbs. Azoxystrobin) (21 day interval)	Apply preventatively when conditions favor disease development.
Gray Snow Mold <i>Typhula</i> spp., Pink Snow Mold <i>Microdochium nivale</i>	1.16 (0.017 lbs. Fluazinam, 0.0059 lbs. Tebuconazole, 0.0053 lbs. Azoxystrobin) (Late Fall)	Apply just before snow cover.
Leaf spot, Melting Out, Brown Blight <i>Drechslera</i> spp. (including <i>D. poae</i> , <i>D. siccans</i>)]	1.16 (0.017 lbs. Fluazinam, 0.0059 lbs. Tebuconazole, 0.0053 lbs. Azoxystrobin) (21 day interval)	[Apply preventatively when conditions favor disease development.
Microdochium Patch , <i>Microdochium nivale</i>	1.16 (0.017 lbs. Fluazinam, 0.0059 lbs. Tebuconazole, 0.0053 lbs. Azoxystrobin) (21 day interval)	Apply in the fall or early spring when conditions are favorable for disease development.
Stem rust <i>Puccinia</i> <i>graminis</i> var. <i>graminicola</i>	1.16 (0.017 lbs. Fluazinam, 0.0059 lbs. Tebuconazole, 0.0053 lbs. Azoxystrobin) (14 day interval)	Apply preventatively when conditions favor disease development (cool and cloudy weather).
Summer Patch <i>Magnaporthe poae</i>]	1.16 (0.017 lbs. Fluazinam, 0.0059 lbs. Tebuconazole, 0.0053 lbs. Azoxystrobin)	Apply preventatively when conditions favor disease development.

[Text in brackets and italics are notes to the label editor.]

[Text in brackets alone are optional statements or alternative statements.]

[Text separated by “/” indicates an either/or choice.]

	(21 day interval)	
Take-All Patch <i>Gaeumannomyces</i>	1.16 (0.017 lbs. Fluazinam, 0.0059 lbs. Tebuconazole, 0.0053 lbs. Azoxystrobin) (21 day interval)	Water in immediately with 0.125-0.25 inch of irrigation.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in dry area. **DO NOT** store in a manner where cross-contamination with other pesticides, fertilizers, food or feed could occur. If spilled during storage or handling, absorb with sand or other inert material and dispose of absorbent in accordance with the Pesticide Disposal instructions listed below.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: [Nonrefillable Plastic Containers 5 Gallons or Less]

Nonrefillable plastic container. DO NOT reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration.

[Nonrefillable plastic containers larger than 5 gallons]

Nonrefillable plastic container. DO NOT reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

LIMITATION OF WARRANTY AND LIABILITY

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Control Solutions, Inc. or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Control Solutions, Inc. and Seller harmless for any claims relating to such factors.

Control Solutions, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Control Solutions, Inc., and Buyer and User assume the risk of any such use. To the extent permitted by applicable law, **CONTROL SOLUTIONS, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

IT IS CONTROL SOLUTIONS INC., AND SELLERS INTENTION THAT in no event shall Control Solutions or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF CONTROL SOLUTIONS, INC., AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF CONTROL SOLUTIONS, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Control Solutions, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Control Solutions, Inc.

{Optional Marketing Claims}

- Combination Chemistry
- 3 Active Ingredients
- Multiple Modes of Action
- Proprietary Formulation
- Broad Spectrum Disease Control
- Cures & Prevents common Turf Diseases
- Controls Brown Patch, Dollar Spot, rusts, Summer Patch, Anthracnose and other listed diseases.
- Easy to mix formulation

{Optional Marketing Graphics}

