

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Virbac, Inc.  
P.O. Box 16598  
Ft. Worth, TX 76162

No 249264  
78/2

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.

2. Make the labeling changes listed below before you release the product for shipment:

**BEST AVAILABLE COPY**

- a. Add the phrase, "EPA Registration No. 54022-a."
- b. Change the Note to Physician to read:  
Note to Physician: Diazinon is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PP is also antidotal and may be administered, but only in conjunction with atropine.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE

July 11, 1978

Richards

**PROTECTION™ Plus 150 Flea and Tick Collar for Dogs  
with EFA\***

\*Slow release coat conditioner helps enhance texture, manageability and luster.

**Kills fleas for 5 months  
Kills ticks for 5 months**

(Color of collar visible through affixed blister package)

**ACTIVE INGREDIENT:**

Diazinon [O,O-diethyl O-(2-isopropyl-6-  
methyl-4-pyrimidinyl)

phosphorothioate] ..... 11.0 %

**INERT INGREDIENTS\*** ..... 89.0 %

**TOTAL** ..... 100.0%

\*Includes 5% unsaturated fatty acids as a grooming aid.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION:** See back panel for additional precautionary statements and directions.

For neck size up to 22".

**FRANCODEX**

**CONTAINS 1 FLEA AND TICK COLLAR**

**NET WT. 1.3 OZ. (36.0 g)**

54022-I

54022-8

## **BLISTER PACKAGE**

### **BACK PANEL (CARDBOARD BACKING)**

#### **PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar.

**Note to Physicians/Veterinarians:** This product contains a cholinesterase inhibitor. Atropine sulfate and 2-PAM are antidotal only if signs of cholinesterase inhibition are present.

Do not use other pesticides on dog while collar is worn. Some dogs and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing dogs.

**Adjustable - fits mature dogs, 22" neck or smaller**

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this collar in a manner inconsistent with its labeling.

Place Protection™ Plus 150 collar around dog's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and dog's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days and will continue killing and controlling ticks for up to 150 days. For unusually large breeds, collar aids in tick control in the head and neck area for up to 2 months. Fleas and ticks present in the animal's environment that may reappear on the dog will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.

This Protection™ Plus 150 collar contains Essential Fatty Acids necessary for healthy skin and hair. Unsaturated fatty acids released from the collar are rapidly absorbed to help maintain a lustrous, manageable coat.

#### **DISPOSAL**

Do not reuse empty blister. Wrap blister and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

EPA Reg. No. 54022-xx-xxxxx

EPA Est. 15577-FR-1

Manufactured for Francodex Laboratories, Inc.  
Subsidiary of Virbac, Inc.  
Fort Worth, Texas 76118  
Made in France

**FOIL POUCH/CARTON PACKAGE**

**FRONT PANEL (CARTON)**

**PROTECTION™ Plus 150 Flea and Tick Collar for Dogs  
with EFA\*  
(color)**

\*Slow release coat conditioner helps enhance texture, manageability and luster.

**Kills fleas for 5 months  
Kills ticks for 5 months**

For neck size up to 22".

**ACTIVE INGREDIENT:**

Diazinon [O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl)  
phosphorothioate]

.....	11.0 %
<b>INERT INGREDIENTS*</b> .....	<b>89.0 %</b>
<b>TOTAL</b> .....	<b>100.0%</b>

\*Includes 5% unsaturated fatty acids as a grooming aid.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION:** See back panel for additional precautionary statements and directions.

**FRANCODEX**

**CONTAINS 1 FLEA AND TICK COLLAR  
NET WT. 1.3 OZ. (36.0 g)**

**FOIL POUCH/CARTON PACKAGE**

**BACK PANEL(CARTON)**

**PROTECTION™ Plus 150 Flea and Tick Collar for Dogs with EFA (Color)  
Adjustable - fits mature dogs, 22" neck or smaller.**

**PRECAUTIONARY STATEMENTS  
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar.**

**Note to Physicians/Veterinarians: This product contains a cholinesterase inhibitor. Atropine sulfate and 2-PAM are antidotal only if signs of cholinesterase inhibition are present.**

**Do not use other pesticides on dog while collar is worn. Some dogs and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing dogs.**

**DIRECTIONS FOR USE**

**It is a violation of Federal law to use this collar in a manner inconsistent with its labeling.**

**Place Protection™ Plus 150 collar around the dog's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and dog's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days and will continue killing and controlling ticks for up to 150 days. For unusually large breeds, collar aids in tick control in the head and neck area for up to 2 months. Fleas and ticks present in the animal's environment that may reappear on the dog will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.**

**This Protection™ Plus 150 collar contains Essential Fatty Acids necessary for healthy skin and hair. Unsaturated fatty acids released from the collar are rapidly absorbed to help maintain a lustrous, manageable coat.**

**DISPOSAL**

**Do not reuse empty pouch or box. Wrap pouch or box and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.**

**EPA Reg. No. 54022-xx-xxxxx**

**EPA Est. 15577-FR-1**

**Manufactured for Francodex Laboratories, Inc.**

**Subsidiary of Virbac, Inc.**

**Fort Worth, Texas 76118**

**Made in France**

FOIL POUCH/CARTON PACKAGE

POUCH LABEL

**PROTECTION™ Plus 150 Flea and Tick Collar for Dogs  
with EFA**

**Do not open until ready to use**

**CAUTION**

**KEEP OUT OF THE REACH OF CHILDREN**

**SEE BACK PANEL OF OUTER CARTON FOR  
ADDITIONAL PRECAUTIONARY STATEMENTS**

**ACTIVE INGREDIENT:**

Diazinon [O,O-diethyl O-(2-isopropyl-6-  
methyl-4-pyrimidinyl)

phosphorothioate] ..... 11.0 %

INERT INGREDIENTS\* ..... 89.0 %

TOTAL ..... 100.0%

\*Includes 5% unsaturated fatty acids as a grooming aid

EPA Reg. No. 54022-xx-xxxxx

EPA Est. 15577-FR-1

Manufactured for Francodex Laboratories, Inc.  
Subsidiary of Virbac, Inc.  
Fort Worth, Texas 76118  
Made in France